













# **INTERMEDIATE TRIGONOMETRY**

## **OUR COLLEGE PUBLICATIONS**

**By Das & Mukherjee**

1. **INTERMEDIATE STATICS** (*14th Edition*)
2. **INTERMEDIATE DYNAMICS** (*14th Edition*)
3. **HIGHER TRIGONOMETRY** (*19th Edition*)
4. **INTEGRAL CALCULUS** (*21st Edition*)
5. **DIFFERENTIAL CALCULUS** (*17th Edition*)
6. **A SHORT COURSE OF COMPLEX VARIABLES  
& HIGHER TRIGONOMETRY** (*3rd Edition*)
7. **ANALYTICAL DYNAMICS  
OF A PARTICLE** (*3rd Edition*)
8. **ELEMENTS OF CO-ORDINATE AND SOLID  
GEOMETRY** (*4th Edition*)  
**By Ganguli & Mukherjee**
9. **INTERMEDIATE ALGEBRA** (*23rd Edition*)
10. **PRE-UNIVERSITY ALGEBRA** (*5th Edition*)  
**By Drs. J. G. Chakravarti & P. R. Ghosh**
11. **HIGHER ALGEBRA**
12. **VECTOR ANALYSIS** (*Pass & Hons.*)  
**By S. Mukherjee & N. Das**
13. **KEY TO INTER./P. U. TRIGONOMETRY**
14. **KEY TO STATICS & DYNAMICS**
15. **KEY TO HIGHER TRIGONOMETRY**
16. **KEY TO DIFFERENTIAL CALCULUS**
17. **KEY TO COMPLEX VARIABLES & HIGHER TRIG.**  
**By P. K. Das**
18. **KEY TO ANALYTICAL DYNAMICS OF A PARTICLE**  
**By Pain & Pain**
19. **KEY TO HIGHER ALGEBRA**
20. **KEY TO INTERMEDIATE ALGEBRA**
21. **KEY TO PRE-UNIVERSITY ALGEBRA**
22. **KEY TO INTEGRAL CALCULUS**

# INTERMEDIATE TRIGONOMETRY

BY

**B. C. DAS, M. Sc.**

PROFESSOR OF MATHEMATICS, PRESIDENCY COLLEGE,  
CALCUTTA : (RETD.),  
EX-LECTURER IN APPLIED MATHEMATICS, CALCUTTA UNIVERSITY

AND

**B. N. MUKHERJEE, M. A.**

*Premchand Roychand Scholar*

PROFESSOR OF MATHEMATICS,  
SCOTTISH CHURCH COLLEGE, CALCUTTA (RETD.)

**U. N. DHUR & SONS PRIVATE LTD.**

**BOOK-SELLERS & PUBLISHERS**

**15, BANKIM CHATTERJEE STREET, CALCUTTA 12**

*Rs. 7 50 only*

*Published by*  
**DWIJENDRANATH DHUB, LL.B.**  
**For U. N. DHUB & SONS PRIVATE LTD.,**  
**15, Bankim Chatterjee Street, Calcutta 12**

**25TH EDITION—1948**

*Printed by*  
**DESH BASU, B. A.,**  
**THE S. S. BASU PRINTING WORKS,**  
**11, Gossain Lane, Calcutta 6**

## PREFACE TO THE FIRST EDITION

**T**HIS book, as its name indicates, is meant to be a textbook for the Intermediate students of Indian Universities, especially the University of Calcutta. Regarding the subject-matter, we have tried to make the exposition clear and concise, without going into unnecessary details. A good number of examples have been worked out by way of illustrations, and examples set have been carefully selected.

Important formulæ and results have been given at the beginning of the book for reference. Calcutta University questions of recent years are given at the end, to give the students an idea of the standard of the examination.

It is hoped that the book will meet the requirements of those for whom it is intended and we shall deem our labours amply rewarded if the students find the book useful to them.

The book had to be hurried through to the press practically within the period of a fortnight, and we must thank the authorities and officers of the K. P. Basu Printing Works, Calcutta, who, in spite of their various preoccupations had the kindness to complete the printing in such a short period of time.

Any criticism, correction and suggestion towards improvement will be thankfully received.

B. C. D.  
B. N. M.



# CONTENTS

CHAP.		PAGE
I.	Measurement of angles ... ..	1
II.	Trigonometrical ratios ... ..	15
III.	Trigonometrical ratios of some standard angles ... ..	29
IV.	Trigonometrical ratios of angles associated with a given angle $\theta$ ... ..	37
V.	Simple practical applications of Trigonometry ... ..	52
VI.	Compound angles ... ..	60
VII.	Transformations of Products and Sums ... ..	71
VIII.	Multiple angles ... ..	76
IX.	Sub-multiple angles ... ..	82
X.	Trigonometrical Identities ... ..	88
XI.	Trigonometrical Equations and general values ... ..	98
XII.	Inverse Circular Functions ... ..	112
	<i>Miscellaneous Examples I</i> ... ..	122
XIII.	Logarithms ... ..	124
XIV.	Properties of triangles ... ..	145
XV.	Solution of triangles ... ..	169
	<i>Miscellaneous Examples II</i> ... ..	186
XVI.	Graphs of Trigonometrical Functions ... ..	189



CHAP.		PAGE
XVII.	Miscellaneous Theorems and Examples ...	215
	SEC. A—Heights and Distances ...	215
	SEC. B—Summation of Series ...	225
	SEC. C—Elimination ...	233
	Appendix ...	238
	<i>Miscellaneous Examples III</i> ...	251
	Answers ...	258
	University Papers ...	268
	Mathematical Tables	i-xxii

---

**TRIGONOMETRY SYLLABUS  
OF THE CALCUTTA UNIVERSITY  
FOR  
I. A. & I. Sc. EXAMINATIONS**

Measurement of angles.

Trigonometrical ratios.

Applications of algebraic signs ; angles of any magnitude.

Graphs of Trigonometrical ratios.

Elementary Trigonometrical formulæ and their applications.

Logarithmic sines, cosines etc.

Relations between the sides and angles of a triangle.

Practical solutions of triangles with applications.

Elementary cases of Inverse Functions.

**GREEK LETTERS USED IN THE BOOK**

$\alpha$ (Alpha)	$\theta$ (Theta)
$\beta$ (Beta)	$\pi$ (Pai)
$\gamma$ (Gamma)	$\phi$ (Phai)
$\delta$ (Delta)	$\psi$ (Psi).

$\Delta$  (Delta)

**Note.** The notation *C. U.* used at the end of any example means that the example was set in the Intermediate Examination of the Calcutta University.



## IMPORTANT FORMULÆ AND RESULTS

I. A radian =  $57^{\circ} 17' 44''$  nearly.

1 degree =  $\cdot 01745$  radians nearly.

2 right angles =  $180^{\circ} = \pi$  radians.

$\pi = \frac{22}{7} = 3\cdot 1416$  approximately.

Radian measure of an angle at the centre of a circle  

$$= \frac{\text{subtending arc}}{\text{radius}}$$

$$\text{II. } \left. \begin{array}{l} \sin^2 \theta + \cos^2 \theta = 1 ; \\ \sec^2 \theta = 1 + \tan^2 \theta ; \\ \operatorname{cosec}^2 \theta = 1 + \cot^2 \theta , \end{array} \right\} \begin{array}{l} \frac{\sin \theta}{\cos \theta} = \tan \theta . \\ \frac{\cos \theta}{\sin \theta} = \cot \theta . \end{array}$$

$$\begin{array}{lll} \text{II. } \sin 0^{\circ} = 0 ; & \cos 0^{\circ} = 1 ; & \tan 0^{\circ} = 0 . \\ \sin 30^{\circ} = \frac{1}{2} ; & \cos 30^{\circ} = \frac{\sqrt{3}}{2} ; & \tan 30^{\circ} = \frac{1}{\sqrt{3}} \\ \sin 45^{\circ} = \frac{1}{\sqrt{2}} ; & \cos 45^{\circ} = \frac{1}{\sqrt{2}} ; & \tan 45^{\circ} = 1 . \\ \sin 60^{\circ} = \frac{\sqrt{3}}{2} ; & \cos 60^{\circ} = \frac{1}{2} ; & \tan 60^{\circ} = \sqrt{3} \\ \sin 90^{\circ} = 1 ; & \cos 90^{\circ} = 0 ; & \tan 90^{\circ} = \infty . \\ \sin 15^{\circ} = \frac{\sqrt{3}-1}{2\sqrt{2}} ; & \cos 15^{\circ} = \frac{\sqrt{3}+1}{2\sqrt{2}} ; & \tan 15^{\circ} = 2 - \sqrt{3} . \\ \sin 75^{\circ} = \frac{\sqrt{3}+1}{2\sqrt{2}} ; & \cos 75^{\circ} = \frac{\sqrt{3}-1}{2\sqrt{2}} ; & \tan 75^{\circ} = 2 + \sqrt{3} . \\ \sin 18^{\circ} = \frac{1}{2}(\sqrt{5}-1) ; & \cos 36^{\circ} = \frac{1}{2}(\sqrt{5}+1) . \\ \sin 120^{\circ} = \frac{\sqrt{3}}{2} ; & \cos 120^{\circ} = -\frac{1}{2} . \\ \sin 180^{\circ} = 0 ; & \cos 180^{\circ} = -1 ; & \tan 180^{\circ} = 0 . \\ \sin 270^{\circ} = -1 ; & \cos 270^{\circ} = 0 ; & \tan 270^{\circ} = \infty . \\ \sin 360^{\circ} = 0 ; & \cos 360^{\circ} = 1 ; & \tan 360^{\circ} = 0 . \end{array}$$

IV.  $\sin(-\theta) = -\sin \theta$ ;  $\cos(-\theta) = \cos \theta$ ;  $\tan(-\theta) = -\tan \theta$

$\sin(90^\circ - \theta) = \cos \theta$ ;  $\sin(90^\circ + \theta) = \cos \theta$ .


$\cos(90^\circ - \theta) = \sin \theta$ ;  $\cos(90^\circ + \theta) = -\sin \theta$ .

$\tan(90^\circ - \theta) = \cot \theta$ ;  $\tan(90^\circ + \theta) = -\cot \theta$ .

$\sin(180^\circ - \theta) = \sin \theta$ ;  $\sin(180^\circ + \theta) = -\sin \theta$ .

$\cos(180^\circ - \theta) = -\cos \theta$ ;  $\cos(180^\circ + \theta) = -\cos \theta$ .

$\tan(180^\circ - \theta) = -\tan \theta$ ;  $\tan(180^\circ + \theta) = \tan \theta$ .



V.  $\sin(A+B) = \sin A \cos B + \cos A \sin B$

$\sin(A-B) = \sin A \cos B - \cos A \sin B$

$\cos(A+B) = \cos A \cos B - \sin A \sin B$

$\cos(A-B) = \cos A \cos B + \sin A \sin B$ .

$\tan(A+B) = \frac{\tan A + \tan B}{1 - \tan A \tan B}$

$\tan(A-B) = \frac{\tan A - \tan B}{1 + \tan A \tan B}$

$\tan(A+B+C)$

$= \frac{\tan A + \tan B + \tan C - \tan A \tan B \tan C}{1 - \tan B \tan C - \tan C \tan A - \tan A \tan B}$ .

VI.  $2 \sin A \cos B = \sin(A+B) + \sin(A-B)$

$2 \cos A \sin B = \sin(A+B) - \sin(A-B)$

$2 \cos A \cos B = \cos(A+B) + \cos(A-B)$

$2 \sin A \sin B = \cos(A-B) - \cos(A+B)$ .

VII.  $\sin C + \sin D = 2 \sin \frac{C+D}{2} \cos \frac{C-D}{2}$

$\sin C - \sin D = 2 \cos \frac{C+D}{2} \sin \frac{C-D}{2}$

$$\cos C + \cos D = 2 \cos \frac{C+D}{2} \cos \frac{C-D}{2}$$

$$\cos C - \cos D = 2 \sin \frac{C+D}{2} \sin \frac{D-C}{2}$$

VIII.  $\sin 2A = 2 \sin A \cos A$

$$\cos 2A = \cos^2 A - \sin^2 A = 1 - 2 \sin^2 A = 2 \cos^2 A - 1$$

$$\tan 2A = \frac{2 \tan A}{1 - \tan^2 A}$$

$$\sin 2A = \frac{2 \tan A}{1 + \tan^2 A}; \cos 2A = \frac{1 - \tan^2 A}{1 + \tan^2 A}$$

$$\left. \begin{aligned} 1 - \cos 2A &= 2 \sin^2 A \\ 1 + \cos 2A &= 2 \cos^2 A \end{aligned} \right\}$$

$$\tan^2 A = \frac{1 - \cos 2A}{1 + \cos 2A}$$

IX.  $\sin 3A = 3 \sin A - 4 \sin^3 A,$

$$\cos 3A = 4 \cos^3 A - 3 \cos A,$$

$$\tan 3A = \frac{3 \tan A - \tan^3 A}{1 - 3 \tan^2 A}$$

X.  $\sin \theta = 2 \sin \frac{\theta}{2} \cos \frac{\theta}{2}$

$$\cos \theta = \cos^2 \frac{\theta}{2} - \sin^2 \frac{\theta}{2} = 2 \cos^2 \frac{\theta}{2} - 1 = 1 - 2 \sin^2 \frac{\theta}{2}$$

$$\tan \theta = \frac{2 \tan \frac{\theta}{2}}{1 - \tan^2 \frac{\theta}{2}}$$

$$\sin \theta = \frac{2 \tan \frac{\theta}{2}}{1 + \tan^2 \frac{\theta}{2}}; \cos \theta = \frac{1 - \tan^2 \frac{\theta}{2}}{1 + \tan^2 \frac{\theta}{2}}$$

$$1 - \cos \theta = 2 \sin^2 \frac{\theta}{2}$$

$$1 + \cos \theta = 2 \cos^2 \frac{\theta}{2}$$

$$\frac{1 - \cos \theta}{1 + \cos \theta} = \tan^2 \frac{\theta}{2}$$

**XI.** If  $\sin \theta = \sin a$ , then  $\theta = n\pi + (-1)^n a$ .

If  $\cos \theta = \cos a$ , then  $\theta = 2n\pi \pm a$ .

If  $\tan \theta = \tan a$ , then  $\theta = n\pi + a$ .

If  $\sin \theta = 0$ , or,  $\tan \theta = 0$ ,  $\theta = n\pi$ .

If  $\cos \theta = 0$ , or,  $\cot \theta = 0$ ,  $\theta = (2n+1)\frac{\pi}{2}$ .

If  $\sin \theta = 1$ ,  $\theta = (4m+1)\frac{\pi}{2}$ ; if  $\sin \theta = -1$ ,  $\theta = (4m-1)\frac{\pi}{2}$

If  $\cos \theta = 1$ ,  $\theta = 2m\pi$ ; if  $\cos \theta = -1$ ,  $\theta = (2m+1)\pi$ .

**XII.**  $\sin^{-1}x + \cos^{-1}x = \frac{1}{2}\pi$

$\tan^{-1}x + \cot^{-1}x = \frac{1}{2}\pi$

$\sec^{-1}x + \operatorname{cosec}^{-1}x = \frac{1}{2}\pi$

$\tan^{-1}x + \tan^{-1}y = \tan^{-1} \frac{x+y}{1-xy}$

$\tan^{-1}x - \tan^{-1}y = \tan^{-1} \frac{x-y}{1+xy}$

$\tan^{-1}x + \tan^{-1}y + \tan^{-1}z = \tan^{-1} \frac{x+y+z-xyz}{1-yz-zx-xy}$

$\sin^{-1}x \pm \sin^{-1}y = \sin^{-1}\{x\sqrt{1-y^2} \pm y\sqrt{1-x^2}\}$ .

$\cos^{-1}x \pm \cos^{-1}y = \cos^{-1}\{xy \mp \sqrt{1-x^2} \cdot \sqrt{1-y^2}\}$ .

**XIII.**  $\log_a mn = \log_a m + \log_a n$

$\log_a \frac{m}{n} = \log_a m - \log_a n$ ;  $\log_a m^n = n \log_a m$ ;

$\log_a m = \log_b m \times \log_a b$ ;  $\log_a 1 = 0$ ;  $\log_a a = 1$ .

**XIV.**  $\frac{a}{\sin A} = \frac{b}{\sin B} = \frac{c}{\sin C} = 2R$

$\cos A = \frac{b^2 + c^2 - a^2}{2bc}$ ;

$\cos B = \frac{c^2 + a^2 - b^2}{2ca}$ ;

$\cos C = \frac{a^2 + b^2 - c^2}{2ab}$ ;

$$a = b \cos C + c \cos B.$$

$$b = c \cos A + a \cos C.$$

$$c = a \cos B + b \cos A.$$

$$\sin \frac{A}{2} = \sqrt{\frac{(s-b)(s-c)}{bc}}$$

$$\cos \frac{A}{2} = \sqrt{\frac{s(s-a)}{bc}}$$

$$\tan \frac{A}{2} = \sqrt{\frac{(s-b)(s-c)}{s(s-a)}}$$

$$\sin A = \frac{2}{bc} \sqrt{s(s-a)(s-b)(s-c)} = \frac{2\Delta}{bc}$$

$$\sin B = \frac{2}{ca} \sqrt{s(s-a)(s-b)(s-c)} = \frac{2\Delta}{ca}$$

$$\sin C = \frac{2}{ab} \sqrt{s(s-a)(s-b)(s-c)} = \frac{2\Delta}{ab}.$$

$$\begin{aligned} \Delta &= \frac{1}{2}bc \sin A = \frac{1}{2}ca \sin B = \frac{1}{2}ab \sin C \\ &= \sqrt{s(s-a)(s-b)(s-c)}, \text{ where } 2s = a + b + c \\ &= \frac{abc}{4R}. \end{aligned}$$

$$R = \frac{a}{2 \sin A} = \frac{b}{2 \sin B} = \frac{c}{2 \sin C} = \frac{abc}{4\Delta}.$$

$$\begin{aligned} r &= \frac{\Delta}{s} = 4R \sin \frac{1}{2}A \sin \frac{1}{2}B \sin \frac{1}{2}C \\ &= (s-a) \tan \frac{1}{2}A = (s-b) \tan \frac{1}{2}B = (s-c) \tan \frac{1}{2}C. \end{aligned}$$

$$\begin{aligned} r_1 &= \frac{\Delta}{s-a} = 4R \sin \frac{1}{2}A \cos \frac{1}{2}B \cos \frac{1}{2}C \\ &= s \tan \frac{1}{2}A. \end{aligned}$$

$$\begin{aligned} r_2 &= \frac{\Delta}{s-b} = 4R \cos \frac{1}{2}A \sin \frac{1}{2}B \cos \frac{1}{2}C \\ &= s \tan \frac{1}{2}B. \end{aligned}$$

$$\begin{aligned} r_3 &= \frac{\Delta}{s-c} = 4R \cos \frac{1}{2}A \cos \frac{1}{2}B \sin \frac{1}{2}C \\ &= s \tan \frac{1}{2}C. \end{aligned}$$



### IMPORTANT RESULTS

1. If  $A + B + C = \pi$ , then

$$\checkmark (i) \sin A + \sin B + \sin C = 4 \cos \frac{1}{2}A \cos \frac{1}{2}B \cos \frac{1}{2}C.$$

$$\times (ii) \cos A + \cos B + \cos C = 1 + 4 \sin \frac{1}{2}A \sin \frac{1}{2}B \sin \frac{1}{2}C.$$

$$(iii) \tan A + \tan B + \tan C = \tan A \tan B \tan C.$$

$$(iv) \sin 2A + \sin 2B + \sin 2C = 4 \sin A \sin B \sin C.$$

$$(v) \cos 2A + \cos 2B + \cos 2C \\ = -4 \cos A \cos B \cos C - 1.$$

$$(vi) \cos^2 A + \cos^2 B + \cos^2 C + 2 \cos A \cos B \cos C = 1.$$

$$(vii) \cot B \cot C + \cot C \cot A + \cot A \cot B = 1.$$

$$(viii) \sin \frac{A}{2} + \sin \frac{B}{2} + \sin \frac{C}{2} \\ = 1 + 4 \sin \frac{B+C}{4} \sin \frac{C+A}{4} \sin \frac{A+B}{4}.$$

$$(ix) \cos \frac{A}{2} + \cos \frac{B}{2} + \cos \frac{C}{2} \\ = 4 \cos \frac{B+C}{4} \cos \frac{C+A}{4} \cos \frac{A+B}{4}.$$

$$(x) \tan \frac{B}{2} \tan \frac{C}{2} + \tan \frac{C}{2} \tan \frac{A}{2} + \tan \frac{A}{2} \tan \frac{B}{2} = 1.$$

$$(xi) \cot \frac{A}{2} + \cot \frac{B}{2} + \cot \frac{C}{2} = \cot \frac{A}{2} \cot \frac{B}{2} \cot \frac{C}{2}.$$

$$2. \lim_{\theta \rightarrow 0} \frac{\sin \theta}{\theta} = 1; \lim_{\theta \rightarrow 0} \cos \theta = 1, \lim_{\theta \rightarrow 0} \frac{\tan \theta}{\theta} = 1.$$

3. Area of a circle of radius  $r = \pi r^2$ .

Perimeter of a circle of radius  $r = 2\pi r$ .

---

# INTERMEDIATE TRIGONOMETRY



## CHAPTER I

### MEASUREMENT OF ANGLES

1. **TRIGONOMETRY**, as indicated by its very name, originally meant a subject which dealt with the methods of measurements of triangles. At present its scope has widened, and now it means a subject which deals with the measurements relating to any angle, not necessarily an angle of a triangle.


#### 2. Angles in Trigonometry.

In Geometry, angles are supposed to be formed by the intersection of two straight lines and are always restricted to lie between  $0^\circ$  and  $360^\circ$ , being acute, obtuse or reflex. Moreover, they are always positive, negative angles having no meaning. In Trigonometry however, the idea of an angle is much more general.

An angle in Trigonometry is supposed to be formed by the revolution of a straight line which starts from an initial position coinciding with one arm, and traces out the angle by its revolution about one extremity until it reaches the final position coinciding with the other arm.

For instance, the angle  $XOP$  is formed by the revolution of a line which starts from the initial position  $OX$ , and revolving in the anti-clockwise direction, traces out the angle  $XOP$  which is acute. The same line again, starting from  $OX$  and revolving in the anti-clockwise direction may make a complete revolution and further move up to the position  $OQ$ . The angle formed in this case is more than

five right angles. Now revolutions may be clockwise or anti-clockwise. *It is conventional to consider angles formed by the anti-clockwise revolution of the revolving line to be positive.* Angles formed by clockwise revolutions of the



revolving line will then be considered *negative angles*. For example, the angle  $XOR$  measured in the clockwise direction from the initial position  $OX$  is a negative angle.

*Thus, angles in Trigonometry may be of any magnitude and may be positive as well as negative.*

$OX$  being the initial position of the revolving line, produce  $XO$  to  $X'$ , and let  $YOY'$  be the perpendicular line. The whole plane is thus divided into four quadrants, the first being  $XOY$ , the second  $YOX'$ , the third  $X'OY'$ , and the fourth  $Y'OX$ . If we contemplate an angle say  $+920^\circ$  to be traced out by the revolving line, the line must have completed two complete revolutions, thereby describing  $2 \times 360^\circ = 720^\circ$ , and have further traced out an angle  $200^\circ$ , so that the final position of the revolving line is in the third quadrant. Similarly, if we consider an angle  $-1354^\circ$ , the final position of the revolving line is in the first quadrant, for  $-1354^\circ = -360^\circ \times 3 - 274^\circ$ .

It should be noted that if two angles differ by complete multiples of  $360^\circ$ , the starting line being the same, the final

positions of the revolving line will be coincident for the two angles. For example, the angles  $255^\circ$  and  $-105^\circ$  will have the final positions of the revolving line same, if both start from the same initial position.

### 3. Units of measurement of angles.

We should now define the different systems of units used for the measurement of angles. In defining a unit however, a standard angle, which has no reference to any particular system of unit, should form the basis, and such a standard angle is a right angle. A right angle is defined in books on Geometry to be an angle which any straight line standing on another makes with it, when the two adjacent angles formed are equal to one another. A right angle is always the same everywhere, and it thus forms a suitable basis to start with, in defining the different systems of units of measurement of angles.

There are three systems of units used in Trigonometry for measurement of angles, *viz.*

- (i) Sexagesimal unit.
- (ii) Centesimal unit.
- (iii) Circular unit.

**Sexagesimal\* System.** In this system, a right angle is divided into 90 equal parts, each being called a *degree*. A degree is again divided into 60 sexagesimal *minutes*, and each minute is further subdivided into 60 sexagesimal *seconds*, so that

$$\begin{aligned} 1 \text{ rt. angle} &= 90^\circ \text{ (degrees)} \\ 1^\circ &= 60' \text{ (sexagesimal minutes)} \\ 1' &= 60'' \text{ (sexagesimal seconds)} \end{aligned}$$

\* So called, since the subdivisions are mostly by sixtieth parts. It is also called the *Common* or the *English System*.

**Centesimal † System.** In this system, the subdivisions of a right angle are as follows :

$$1 \text{ rt. angle} = 100^g \text{ (grades)}$$

$$1^g = 100' \text{ (centesimal minutes)}$$

$$1' = 100'' \text{ (centesimal seconds)}$$

**Note.** It may be noted that  $1'$  (centesimal minute) is not the same as  $1'$  (sexagesimal minute), the former being  $\frac{1}{100 \times 100}$  of a right angle and the latter being  $\frac{1}{90 \times 60}$  of a right angle, so that the first is  $\frac{3}{8}$ th part of the second. Similarly,  $1''$  is less than  $1''$ , being only  $\frac{3}{8}$ th part of it.

The connection between the two systems of units may be effected through a right angle, remembering that  $1 \text{ right angle} = 90^\circ = 100^g$ , so that  $9^\circ = 10^g$ . Any angle in the first system may be reduced to degrees, and then multiplied by  $\frac{1}{9}$  will be reduced to grades. Similarly, an angle in the second system may be changed to the first.

We shall presently deal with the third system, namely the circular system.

**4. Theorem.** *In all circles, the circumference bears a constant ratio to its diameter.*


Take any two circles of any radii, and place them with a common centre  $O$ . In one, let  $ABCD\dots$  be an inscribed regular polygon of  $n$  sides. Let  $A', B', C', \dots$  be the points of intersection of the radii  $OA, OB, OC, \dots$  with the other circle. It is easily seen that  $A'B'C'\dots$  is also a regular polygon of  $n$  sides, inscribed in the second circle. Now  $OA = OB$ , as also  $OA' = OB'$ , so that in the triangles  $OAB, OA'B'$ ,

† So called, because the subdivisions are by hundredths. It is also called the *French System*.

$OA : OA' = OB : OB'$ , and angle  $O$  is common. The two triangles are therefore similar. Hence  $AB : A'B' = OA : OA'$ .

Thus,

$$\frac{\text{perimeter of polygon } ABCD\dots}{\text{perimeter of polygon } A'B'C'D'\dots} = \frac{n \cdot AB}{n \cdot A'B'} = \frac{OA}{OA'}.$$



This being true, whatever the number of sides  $n$  may be, making  $n$  infinitely large, the perimeters of the polygons can be made practically coincident with the circumferences of the corresponding circles, and thus we deduce that

$$\frac{\text{circumference of the circle } ABCD\dots}{\text{circumference of the circle } A'B'C'D'\dots} = \frac{OA}{OA'}.$$

$$\text{i.e.} = \frac{\text{radius of circle } ABC\dots}{\text{radius of circle } A'B'C'\dots}.$$

Thus circumference of any circle : its radius is the same for all circles. As diameter is twice the radius, we deduce that the circumference of any circle bears a constant ratio to its diameter.

This constant ratio is denoted by the Greek letter  $\pi$ . Its actual value has been determined by methods which are outside the scope of the present book, by some mathematicians

to more than 500 places of decimals. An approximate value commonly used is  $\frac{22}{7}$ . A more accurate value is  $\frac{355}{113}$ .

Expressed in decimal, the value is nearly  $3\cdot14159\dots$

Hence, if  $r$  be the radius of a circle,  $d$  its diameter,

$$\text{the circumference} = \pi d = 2\pi r,$$


where  $\pi = 3\cdot14159\dots = \frac{22}{7}$  roughly.

## 5. Circular Unit or Radian Measure.

In any circle, if we take an arc whose length is equal to the radius of the circle, the angle which this arc subtends at the centre is called a *radian*, and is written as  $1^c$ .

We shall now show that with reference to whichever circle it may be defined, a radian is a constant angle, and hence it may be used as a suitable unit for measurement of angles, which is known as the circular unit.

**Theorem I.** *A radian is a constant angle.*



Let  $AB$  be an arc of any circle with centre  $O$ , whose length is equal to its radius  $OA$ . By definition,  $\angle AOB = 1$  radian. Since angles at the centre of a circle are proportional to the arcs which subtend them, and the whole angle

round  $O$  subtended by the complete circumference being known from Geometry to be 4 right angles, we get

$$\frac{\angle AOB}{4 \text{ right angles}} = \frac{\text{arc } AB}{\text{whole circumference}} = \frac{\text{radius}}{\text{circumference}},$$

$$\text{i.e., } \frac{1 \text{ radian}}{4 \text{ rt. } \angle} = \frac{r}{2\pi r} = \frac{1}{2\pi}, \text{ } r \text{ being the radius.}$$

$$\text{Hence, } 1 \text{ radian} = \frac{2}{\pi} \text{ rt. angle.}$$

$\therefore$  a radian is a constant angle. ( $\pi$  being constant)

**Note.** We thus see that whatever be the radius of the circle with reference to which a radian is defined, its magnitude is the same.

From above,  $\pi$  radians =  $180^\circ$ .

$$\begin{aligned} \therefore 1 \text{ radian} &= \frac{180^\circ}{\pi} = \frac{180}{3.14159} = 57.29577 \text{ degrees} \\ &= 57^\circ 17' 44.8'' \text{ nearly.} \end{aligned}$$

$$\therefore 1 \text{ degree} = .0174533 \text{ radians nearly.}$$

In higher mathematics so far as theoretical investigations are concerned, as a matter of convenience, angles are usually measured in the circular unit, *i.e.* in radians. In this connection we may state the following theorem :

**Theorem II.** *The measure of any angle in radians is expressed by the ratio of the arc of any circle subtending that angle at its centre, to the radius.*

Let  $XOP$  be any angle.

With centre  $O$  and any radius  $OA$  draw a circle, and let  $AQ$  be the arc which subtends the angle  $XOP$  at the centre  $O$ . Let  $AB$  be the arc whose length is equal to the radius  $OA$ , so that, by definition,  $\angle AOB$  is one radian.

Now from Geometry, angles at the centre of a circle are proportional to the arcs which subtend them.




# INTERMEDIATE TRIGONOMETRY

$$\text{Hence, } \frac{\angle XOP}{\angle AOB} = \frac{\text{arc } AQ}{\text{arc } AB} = \frac{\text{arc } AQ}{\text{radius } OA},$$

$$\text{or, } \frac{\angle XOP}{1 \text{ radian}} = \frac{\text{arc } AQ}{\text{radius } OA}$$

$$\text{i.e. } \angle XOP = \frac{\text{arc } AQ}{\text{radius } OA} \text{ of a radian.}$$



Thus if  $\theta$  be the radian-measure of the  $\angle XOP$ ,  $s$  be the length of the arc  $AQ$ , and  $r$  the radius of the circle, then

$$\theta = \frac{s}{r} \text{ or, } s = r\theta.$$

**Note.** In higher mathematics, when an angle is expressed in radian measure, the unit is generally implied and not expressed, so that, when the measure of an angle is given without the unit being mentioned, we should always understand it to be in radians. For example, 'an angle is  $\frac{\pi}{2}$ ' means that the angle is  $\frac{\pi}{2}$  radians, which converted to degrees is  $90^\circ$  i.e. one right angle.

6. In working-out examples, the relations between the three systems of units should be carefully remembered, namely

$$1 \text{ rt. } \angle = 90^\circ = 100^\circ = \frac{\pi}{2} \text{ radians,}$$

whence,

$$\pi^\circ = 180^\circ.$$

**Ex. 1.** *Express*

- (i)  $63^{\circ} 22' 40''$  in centesimal measure  
and (ii)  $203^{\circ} 58' 73''$  in radians.

$$\begin{aligned}\text{Here (i) } 63^{\circ} 22' 40'' &= 63\frac{188}{800} \text{ deg.} = \frac{21888}{800} \times \frac{1}{90} \text{ rt. } \angle \\ &= \frac{21888}{800} \times \frac{1}{90} \times 100 \text{ grades} = \frac{2521}{80} \text{ grades} = 70^{\circ} 42'.\end{aligned}$$

$$\begin{aligned}\text{(ii) } 203^{\circ} 58' 73'' &= 203.5873 \text{ grades} \\ &= 2.035873 \text{ rt. } \angle = 2.035873 \times \frac{\pi}{2} \text{ radians} \\ &= 1.0179365 \pi \text{ radians.}\end{aligned}$$

**Ex. 2.** *Two angles of a triangle are  $72^{\circ} 53' 51''$ , and  $41^{\circ} 22' 50''$  respectively. Find the third angle in radians.*

$$\begin{aligned}41^{\circ} 22' 50'' &= 41.2250 \text{ grades} \\ &= \frac{41.225 \times 9}{10} \text{ degrees } [ 9^{\circ} = 10^{\circ} ] \\ &= 37.1025 \text{ degrees} \\ &= 37^{\circ} 6' 9''.\end{aligned}$$

The sum of the two given angles is therefore

$$72^{\circ} 53' 51'' + 37^{\circ} 6' 9'' = 110^{\circ}.$$

The sum of the three angles of a triangle being  $180^{\circ}$ , the third angle is

$$\begin{aligned}180^{\circ} - 110^{\circ} &= 70^{\circ} = 70 \times \frac{\pi}{180} \text{ radians } [ \pi^{\circ} = 180^{\circ} ] \\ &= \frac{7\pi}{18} \text{ radians.}\end{aligned}$$

**Ex. 3.** *Divide  $\frac{\pi}{4}$  radians into two parts such that the number of sexagesimal minutes in one may be to the number of centesimal seconds in the other part as 27 : 2500.*

$$\text{We have } \frac{\pi}{4} \text{ radians} = \frac{\pi}{4} \times \frac{2}{\pi} \text{ rt. } \angle = \frac{1}{2} \text{ rt. } \angle.$$

Let  $x$  be the number of centesimal seconds in the second part, so that  $\frac{27}{2500}x$  is the number of sexagesimal minutes in the first part.

$$\text{Now } x'' = \frac{x}{100 \times 100 \times 100} \text{ rt. } \angle$$

$$\text{and } \frac{27}{2500}x' = \frac{27x}{2500 \times 60 \times 90} \text{ rt. } \angle = \frac{x}{500000} \text{ rt. } \angle$$

$$\therefore \frac{x}{1000000} + \frac{x}{500000} = \frac{1}{2},$$

$$\text{whence } x = \frac{500000}{3}.$$

$$\text{Thus, second part is } \frac{500000''}{3} = \frac{500000}{3 \times 100 \times 100 \times 100} \text{ rt. } \angle$$

$$= \frac{1}{3} \text{ rt. } \angle = 15^\circ, \text{ and as the sum of the two parts is } \frac{1}{2} \text{ rt. } \angle \text{ i.e. } 45^\circ, \text{ the first part is } 30^\circ.$$

The two parts are therefore  $30^\circ$  and  $15^\circ$ .

**Ex. 4.** *The angles of a quadrilateral are in A.P., and the number of grades in the least angle is to the number of radians in the greatest as  $100 : \pi$ . Find the angles in degrees.*

Let the angles, expressed in degrees, be  $\alpha$ ,  $\alpha + \beta$ ,  $\alpha + 2\beta$  and  $\alpha + 3\beta$  respectively. Then

$$\alpha + \alpha + \beta + \alpha + 2\beta + \alpha + 3\beta = 360, \\ \text{i.e. } 2\alpha + 3\beta = 180. \quad \dots (i)$$

$$\text{Again the least angle, } \alpha^\circ = \frac{100}{\pi} \alpha^\circ$$

$$\text{and the greatest angle } (\alpha + 3\beta)^\circ = (\alpha + 3\beta) \frac{\pi^\circ}{180},$$

and so from the given condition,

$$\frac{10}{9} \alpha / (\alpha + 3\beta) \cdot \frac{\pi}{180} = 100/\pi,$$


$$\text{or, } \frac{2\alpha}{\alpha + 3\beta} = 1, \text{ whence } \alpha = 3\beta.$$

$\therefore$  using (i),  $3\alpha = 180$ , or,  $\alpha = 60$  and  $\beta = \frac{\alpha}{3} = 20$ .

Thus the angles are

$60^\circ, 80^\circ, 100^\circ$  and  $120^\circ$ .

**Ex. 5.** *At what distance does a man,  $5\frac{1}{2}$  ft. in height, subtend an angle of  $15''$ ?*



$AB$  being the man subtending an angle  $15''$  at  $O$ , let  $OA$  be  $r$  ft.

As the angle  $AOB$  is very small, so that  $AB$  is very small compared to  $AO$ , we may assume the small length  $AB$  to be practically a small arc of a circle whose centre is  $O$ . Now the measure of an angle in radians is the ratio of the arc which subtends it at the centre to the radius.

[ See Art. 5. ]

$$\therefore \frac{15}{60 \times 60} \times \frac{\pi}{180} = \frac{5\frac{1}{2}}{r},$$

$$\text{or, } r = \frac{11}{2} \times \frac{180 \times 60 \times 60}{15 \times \pi} \text{ ft.}$$

$$= \frac{11}{2} \times \frac{180 \times 60 \times 60 \times 7}{15 \times 22} \times \frac{1}{3 \times 1760} \text{ miles approx.}$$

$$= 14.32 \text{ miles nearly.}$$

### Examples I

1. Indicate the final position of a revolving line which has traced out the angle

(i)  $1122^\circ$ ;                      (ii)  $-810^\circ 29'$ ;

(iii)  $-617^\circ 51' 5''$ ; (iv)  $\frac{18\pi}{5}$  radians.

2. Express (i)  $55^{\circ} 12' 36''$  in centesimal measure ;  
(ii)  $195^{\circ} 35' 24''$  in degrees, minutes and secs.
3. How many radians are there in (i)  $50^{\circ} 75' 50''$  ;  
(ii)  $18^{\circ} 33' 45''$  ?
4. Express in each system of angular measurement, the angle between the minute hand and the hour hand of a clock at a quarter to twelve.
5. If  $x^{\circ}$  be taken as the unit angle, and the angles  $600^{\circ}$  and  $16^{\circ}$  expressed in that unit be  $\alpha$  and  $\beta$  respectively, find the relation between  $\alpha$  and  $\beta$ .
6. The difference of two angles is  $1^{\circ}$  ; the circular measure of their sum is 1 ; find the circular measure of the smaller angle.
7. Two angles are in the ratio 2 : 3, and the difference of their measure in grades and in degrees respectively is  $2\frac{1}{2}$  ; find the angles in degrees.
8. An angle is the excess of  $D^{\circ} M'$  over  $G^{\circ} m'$ . Find the ratio of this angle to a right angle.
9. The circular measure of a certain angle is equal to the ratio of the number of degrees in it to the number of centesimal minutes ; find the magnitude of the angle in degrees.
10. With two units of angular measurement differing by  $10^{\circ}$ , the measures of an angle are as 3 : 2 ; determine the units.
11. If an angle standing upon an arc of length ' $l$ ' at the centre of a circle of radius ' $r$ ' be taken as unit, and three angles  $D^{\circ}$ ,  $G^{\circ}$ , and  $C$  circular units expressed in that unit be  $x$ ,  $y$ ,  $z$  respectively, show that

$$x : y : z = \frac{D\pi}{18} : \frac{G\pi}{20} : 10C.$$

12. Three angles are in G. P. The number of grades in the greatest angle is to the number of circular units in the least as 800 to  $\pi$ , and the sum of the three angles is  $126^\circ$ . Find the angles in grades.

13. Divide  $54^\circ$  in three parts, such that the circular measure of the first exceeds that of the second by  $\frac{\pi}{10}$ , and the sum of the second and third is 30 grades.

14. Find at what times between 7 and 8 o'clock the angle between the two hands of a clock is (i)  $60^\circ$ , (ii)  $155^\circ$ .

15. The angles of a triangle are in A.P., and the number of radians in the greatest is to the number of grades in the least as  $\pi : 40$ . Find the angles in degrees.

16. In each of two triangles the angles are in G. P. ; the least angle of one of them is three times the least angle in the other, and the sum of the greatest angles is  $240^\circ$ . Find the circular measure of the angles.

17. One angle of a quadrilateral is  $\frac{2}{3}$  of another and the two other angles are  $66\frac{2}{3}$  grades and  $\frac{3\pi}{4}$  radians. Express the angles in degrees.

18. The angles of a polygon (which has no reflex angle) are in A. P. The least angle is  $\frac{2\pi}{3}$  radians and the common difference is  $5^\circ$ . Find the number of sides.

19. The number of sides of two regular polygons are as  $m : n$ , and the number of degrees in an angle of the first is to the number of grades in an angle of the second as  $p : q$ . Determine the number of sides in each polygon.

20. An arc of  $50^\circ$  in one circle equals one of  $60^\circ$  in another ; find the radian measure of an angle subtended at the centre of the first circle by an arc equal to the radius of the second.

21. Two regular figures are such that the number of degrees in an angle of one is to the number of degrees in an angle of the other as the number of sides in the first is to the number of sides in the second. The sum of the number of sides of the two figures being 9, determine the number of sides of each.

22. The wheel of a railway carriage is 4 ft. in diameter and makes 6 revolutions in a second ; how fast is the train going ?

23. The earth revolves round the sun in a circular orbit of radius 92700000 miles once a year. Find its velocity in miles per hour. If the apparent angular diameter of the sun observed from the earth be  $32'$ , find also the linear radius of the sun.

24. A tower subtends an angle of  $10'$  when the observer is at a distance of 6 miles ; find its height.

25. Find the radius of the earth, if an angle of  $1^\circ$  is subtended at its centre by an arc joining two places on it distant 69.1 miles.

26. A horse is tied to a post by a rope 27 feet long. If the horse moves along the circumference of a circle always keeping the rope tight, find how far the horse will have gone when the rope has traced out an angle of  $70^\circ$ . ( $\pi = \frac{22}{7}$ ).

27. A man running along a circular track at the rate of 10 miles per hour, traverses in 36 seconds, an arc which subtends  $56^\circ$  at the centre. Find the diameter of the circle. ( $\pi = \frac{22}{7}$ ).

28. An arc of  $30^\circ$  in one circle is double an arc in a second circle the radius of which is three times the radius of the first. Show that the arc of the second circle subtends  $5^\circ$  at its centre.

---

## CHAPTER II TRIGONOMETRICAL RATIOS

### 7. Trigonometrical ratios defined.\*




Fig. 1




Fig. 2

Let  $\theta$  be the measure of an angle  $XOP$  which may be supposed to be traced out by a revolving line starting from the initial position  $OX$ . From any point  $P$  on its other arm, draw a perpendicular  $PN$  on  $OX$  (produced if necessary, as in the second figure). A right-angled triangle is thereby formed. The trigonometrical ratios of the angle  $\theta$  are defined as follows :—

Sine of the angle  $\theta$ , written as  $\sin \theta = \frac{PN}{OP}$

*i.e.  $\frac{\text{opposite side}}{\text{hypotenuse}}$*

Cosine of  $\theta$ , written as  $\cos \theta = \frac{ON}{OP}$

*i.e.  $\frac{\text{adjacent side}}{\text{hypotenuse}}$*

---

\*For alternative definitions, see Appendix.



Tangent of  $\theta$ , written as  $\tan \theta = \frac{PN}{ON}$

i.e.  $\frac{\text{opposite side}}{\text{adjacent side}}$

Cosecant of  $\theta$ , written as  $\operatorname{cosec} \theta = \frac{OP}{PN}$

i.e.  $\frac{\text{hypotenuse}}{\text{opposite side}}$

Secant of  $\theta$ , written as  $\sec \theta = \frac{OP}{ON}$

i.e.  $\frac{\text{hypotenuse}}{\text{adjacent side}}$

Cotangent of  $\theta$ , written as  $\cot \theta = \frac{ON}{PN}$

i.e.  $\frac{\text{adjacent side}}{\text{opposite side}}$

In addition to these, we define two less important ratios of the angle  $\theta$  which are sometimes used, as follows :—

Versed sine of angle  $\theta$ , written as  $\operatorname{vers} \theta = 1 - \cos \theta$

Coversed sine of angle  $\theta$ , written as  $\operatorname{covers} \theta = 1 - \sin \theta$


## 8. Signs of Trigonometrical ratios.

$XOP$  being any angle, traced out by a revolving line which starts from  $OX$ , it has already been mentioned in the last Chapter that the plane may be divided into four quadrants by the two perpendicular lines  $XOX'$  and  $YOY'$ .

It is conventional, as in graphs, to consider distances measured along  $OX$  and  $OY$  as positive, and along  $OX'$  and  $OY'$  as negative. *The distance measured along  $OP$ , the final position of the revolving line corresponding to the angle  $XOP$ , in whichever quadrant it may lie, is however always considered positive.*

With this convention, if  $OP$  lies in the first quadrant as in Fig. (i) of the last article, the sides  $PN$ ,  $ON$  and  $OP$  of the right-angled triangle  $OPN$  are all positive. Hence all the Trigonometrical ratios are positive. If  $OP$  lies in the third quadrant as in Fig. (ii),  $ON$  and  $PN$  are both negative, but  $OP$  is positive. Hence from the definitions of the Trigonometrical ratios,  $\sin XOP \left( = \frac{PN}{OP} \right)$  is negative,  $\cos XOP \left( = \frac{ON}{OP} \right)$  is negative,  $\tan XOP \left( = \frac{PN}{ON} = \frac{\text{negative quantity}}{\text{negative quantity}} \right)$  is positive etc.


In this way, according to the final position of the revolving line (starting position being  $OX$ ), we can determine the signs of the Trigonometrical ratios of the angle  $XOP$  whether this angle traced out is positive or negative. If  $OP$  is in the *first quadrant*, the ratios are all positive. If  $OP$  falls in the *second quadrant*, sine and cosecant (which is evidently the reciprocal of sine), are positive; all the other ratios are negative. If  $OP$  be in the *third quadrant*, tangent and cotangent (which are reciprocals to each other) are positive; all the others are negative. In the *fourth quadrant*, cosine and secant are positive, others are negative. A symbolical figure will help the memory in this case, namely, that according to the position of  $OP$ ,



The positiveness of sine, cosine and tangent also implies the positiveness of their reciprocals, namely, cosecant, secant and cotangent respectively.

### 9. Constancy of Trigonometrical ratios.

So long as an angle remains the same, its Trigonometrical ratios are unique.



Let  $\angle XOP (= \theta)$  be any angle, and let  $PN$  and  $P'N'$  be drawn perpendiculars upon  $OX$  from any two points  $P$  and  $P'$  on  $OP$ . The two right-angled triangles  $OPN$  and  $OP'N'$  are similar. Hence  $\sin \theta$ , whether we take it as  $\frac{PN}{OP}$  or  $\frac{P'N'}{OP'}$ , is the same. If the angle be  $\angle XOP_1$ , when  $OP_1$  is not in the first quadrant, the right-angled triangles  $P_1N_1O$  and  $P'_1N'_1O$  are not only similar but also have their corresponding sides of the same sign. Hence the Trigonometrical ratios of the angle  $\angle XOP_1$ , whether defined from the triangle  $P_1N_1O$  or from  $P'_1N'_1O$  are the same in magnitude as well as in sign. Thus for any given angle, the Trigonometrical ratios are unique.

**Note.** In case of a *positive acute angle* like  $XOP$ , we might take any point  $Q$  on  $OX$  as well, and draw  $QM$  perpendicular upon  $OP$ , and define  $\sin XOP$  to be  $\frac{\text{opposite side}}{\text{hypotenuse}}$  i.e.  $\frac{QM}{OQ}$ ,  $\cos XOP$  to be  $\frac{OM}{OQ}$  etc. Now the two triangles  $QOM$  and  $PON$  are easily seen to be similar and both have their sides all positive; so that  $\frac{QM}{OQ} = \frac{PN}{OP}$ ,  $\frac{OM}{OQ} = \frac{ON}{OP}$  etc. Hence the Trigonometrical ratios of the angle  $XOP$ , even if defined from triangle  $QOM$ , will have the same values.

It may also be noted that for angles of any magnitude, positive or negative, any of the two arms may be supposed to be coincident with  $OX$ , and then the magnitude and sign of the angle will fix up the position of the other arm, and thereby will make the Trigonometrical ratios unique.

## 10. Fundamental relations between the Trigonometrical ratios of any angle.

From the very definitions given in Art. 7 of the Trigonometrical ratios of any angle  $XOP$  ( $=\theta$ ) of whatever magnitude and sign, we at once derive the following relations :

$$\operatorname{cosec} \theta = \frac{1}{\sin \theta}$$

$$\sec \theta = \frac{1}{\cos \theta}$$

$$\cot \theta = \frac{1}{\tan \theta}$$

$$\text{and since } \sin \theta = \frac{PN}{OP}, \cos \theta = \frac{ON}{OP}, \tan \theta = \frac{PN}{ON},$$

$$\cot \theta = \frac{ON}{PN}, \text{ we get}$$

$$\tan \theta = \frac{\sin \theta}{\cos \theta}$$

$$\cot \theta = \frac{\cos \theta}{\sin \theta}.$$

Again, since in the right-angled triangle  $OPN$ ,

$$OP^2 = PN^2 + ON^2,$$

dividing by  $OP^2$ ,  $ON^2$  and  $PN^2$  respectively, we get

$$\left(\frac{PN}{OP}\right)^2 + \left(\frac{ON}{OP}\right)^2 = 1 \quad \dots \quad \dots \quad (i)$$

$$\left(\frac{OP}{ON}\right)^2 = \left(\frac{PN}{ON}\right)^2 + 1 \quad \dots \quad \dots \quad (ii)$$

$$\left(\frac{OP}{PN}\right)^2 = 1 + \left(\frac{ON}{PN}\right)^2 \quad \dots \quad \dots \quad (iii)$$

From the definition of the Trigonometrical ratios, (i) gives

$$(\sin \theta)^2 + (\cos \theta)^2 = 1$$

Now it is usual to write  $(\sin \theta)^2$  in the form  $\sin^2 \theta$  and so for other ratios. The relation then reduces to the form

$$\sin^2 \theta + \cos^2 \theta = 1.$$

Similarly, (ii) and (iii) give respectively

$$\sec^2 \theta = 1 + \tan^2 \theta$$

$$\operatorname{cosec}^2 \theta = 1 + \cot^2 \theta.$$

These formulæ are also used in the forms

$$\sin^2 \theta = 1 - \cos^2 \theta, \cos^2 \theta = 1 - \sin^2 \theta,$$

$$\sec^2 \theta - \tan^2 \theta = 1, \tan^2 \theta = \sec^2 \theta - 1, \text{ etc.}$$

**Note.** The fundamental formulæ derived in this article are very important, and are true for all values of  $\theta$  whatever its magnitude and sign may be. For example, if we take  $\frac{\theta}{2}$  in place of  $\theta$ , we are simply taking a different angle for which the same relations are true, so that  $\sin^2 \frac{\theta}{2} + \cos^2 \frac{\theta}{2} = 1$ , etc.

## 11. Conversions of Trigonometrical ratios.

With the help of the formulæ of the previous article, we can express any Trigonometrical ratio of an angle in terms of any other ratio for the same angle; hence if the value of any Trigonometrical ratio of an angle be given, we can find the value of any other ratio.

**Ex. 1.** Express  $\sin \theta$  in terms of  $\cot \theta$ .

From the formulæ  $\operatorname{cosec} \theta = \frac{1}{\sin \theta}$

$$\text{and } \operatorname{cosec}^2 \theta = 1 + \cot^2 \theta,$$

$$\text{we get } \sin \theta = \frac{1}{\operatorname{cosec} \theta} = \pm \frac{1}{\sqrt{1 + \cot^2 \theta}}.$$

**Ex. 2.** Express  $\operatorname{cosec} \theta$  in terms of  $\sec \theta$ .

$$\begin{aligned} \operatorname{cosec} \theta &= \pm \sqrt{1 + \cot^2 \theta} = \pm \sqrt{1 + \frac{1}{\tan^2 \theta}} \\ &= \pm \sqrt{\frac{\tan^2 \theta + 1}{\tan^2 \theta}} = \pm \sqrt{\frac{\sec^2 \theta}{\sec^2 \theta - 1}} = \frac{\pm \sec \theta}{\sqrt{\sec^2 \theta - 1}}. \end{aligned}$$

**Ex. 3.** If  $\cos A = \frac{1}{3}$ , find  $\tan A$ .

$$\begin{aligned} \text{We have } \tan A &= \frac{\sin A}{\cos A} = \frac{\pm \sqrt{1 - \cos^2 A}}{\cos A} \\ &= \pm \frac{\sqrt{1 - \frac{1}{9}}}{\frac{1}{3}} = \pm \frac{\frac{2}{3}}{\frac{1}{3}} = \pm \frac{2}{1}. \end{aligned}$$

A more practical method in such cases is however to construct a right-angled triangle with the numerator and denominator as the two suitable sides, as shown below.

**Ex. 4.** If  $\sec A = \frac{41}{9}$ , find  $\cot A$ .


Let  $APN$  be a triangle right-angled at  $N$  in which the hypotenuse  $AP = 41$ ,  $AN = 9$ , so that  $\sec NAP = \frac{AP}{AN} = \frac{41}{9}$ .

Thus  $\angle NAP = A$ .

$$\begin{aligned} \text{Now } PN^2 &= AP^2 - AN^2 = 41^2 - 9^2 \\ &= 40^2, \end{aligned}$$

so that  $PN = \pm 40$ .

$$\cot A = \cot NAP = \frac{AN}{PN} = \pm \frac{9}{40}.$$



### 12. Restrictions on the magnitudes of Trigonometrical ratios.

From the relation  $\sin^2 \theta + \cos^2 \theta = 1$ , since  $\sin^2 \theta$  and  $\cos^2 \theta$  being square quantities are both positive, it is evident that neither  $\sin^2 \theta$  nor  $\cos^2 \theta$  can exceed 1, for if  $\sin^2 \theta$ , for example, be greater than 1,  $\cos^2 \theta$  (which is a square quantity) becomes negative, which is impossible. Thus  $\sin \theta$  as well as  $\cos \theta$  must have numerical values not exceeding 1; in other words, both  $\sin \theta$  and  $\cos \theta$  must lie between +1 and -1 whatever the magnitude of  $\theta$  may be. Any value numerically greater than 1, like -2 or +3.1 must be impossible for  $\sin \theta$  or  $\cos \theta$ .

$\sec \theta$  and  $\operatorname{cosec} \theta$  therefore, being reciprocals of  $\cos \theta$  and  $\sin \theta$  respectively, can never be numerically less than 1.

$\tan \theta$  and  $\cot \theta$  however, can have any numerical value greater than 1 or less than 1 according to the value of  $\theta$ .

13. A few examples on the applications of the fundamental formulæ are given below.

Ex. 1. Prove that  $\sqrt{\frac{1+\cos \theta}{1-\cos \theta}} = \operatorname{cosec} \theta + \cot \theta$ .

[C. U. 1937.]

$$\begin{aligned}\sqrt{\frac{1+\cos \theta}{1-\cos \theta}} &= \sqrt{\frac{(1+\cos \theta)^2}{1-\cos^2 \theta}} = \sqrt{\frac{(1+\cos \theta)^2}{\sin^2 \theta}} \\ &= \frac{1+\cos \theta}{\sin \theta} = \frac{1}{\sin \theta} + \frac{\cos \theta}{\sin \theta} = \operatorname{cosec} \theta + \cot \theta.\end{aligned}$$

Ex. 2. Prove that

$$\frac{1}{\sec A + \tan A} - \frac{1}{\cos A} = \frac{1}{\cos A} - \frac{1}{\sec A - \tan A}.$$

$$\begin{aligned}\text{We have } & \frac{1}{\sec A + \tan A} + \frac{1}{\sec A - \tan A} \\ &= \frac{\sec A - \tan A + \sec A + \tan A}{(\sec A + \tan A)(\sec A - \tan A)} = \frac{2 \sec A}{\sec^2 A - \tan^2 A} \\ &= 2 \sec A = \frac{2}{\cos A} = \frac{1}{\cos A} + \frac{1}{\cos A}.\end{aligned}$$

Hence by transposition,

$$\frac{1}{\sec A + \tan A} - \frac{1}{\cos A} = \frac{1}{\cos A} - \frac{1}{\sec A - \tan A}.$$

Ex. 3. Prove that 
$$\frac{1 + 2 \sin \theta \cos \theta}{(\sin \theta + \cos \theta)(\cot \theta + \tan \theta)} = \sin \theta \cos \theta (\sin \theta + \cos \theta).$$

We have 
$$\begin{aligned} & \frac{1 + 2 \sin \theta \cos \theta}{(\sin \theta + \cos \theta)(\cot \theta + \tan \theta)} \\ &= \frac{(\sin^2 \theta + \cos^2 \theta) + 2 \sin \theta \cos \theta}{(\sin \theta + \cos \theta) \left( \frac{\cos \theta}{\sin \theta} + \frac{\sin \theta}{\cos \theta} \right)} \\ &= \frac{(\sin \theta + \cos \theta)^2}{(\sin \theta + \cos \theta) \left( \frac{\cos^2 \theta + \sin^2 \theta}{\sin \theta \cos \theta} \right)} \\ &= \frac{(\sin \theta + \cos \theta) \sin \theta \cos \theta}{1} \\ &= \sin \theta \cos \theta (\sin \theta + \cos \theta). \end{aligned}$$

Ex. 4. If  $15 \sin^2 \theta + 2 \cos \theta = 7$ , find  $\tan \theta$ .

Here  $15(1 - \cos^2 \theta) + 2 \cos \theta = 7$ ,

whence  $15 \cos^2 \theta - 2 \cos \theta - 8 = 0$ ,

or,  $(5 \cos \theta - 4)(3 \cos \theta + 2) = 0$ ;  $\therefore \cos \theta = \frac{4}{5}$ , or,  $-\frac{2}{3}$ .

Case (i) when  $\cos \theta = \frac{4}{5}$ ,

$$\sin^2 \theta = 1 - \cos^2 \theta = 1 - \frac{16}{25} = \frac{9}{25}. \quad \therefore \sin \theta = \pm \frac{3}{5}.$$

and so  $\tan \theta = \frac{\sin \theta}{\cos \theta} = \pm \frac{3}{4}.$

Case (ii) when  $\cos \theta = -\frac{2}{3}$ ,

$$\sin^2 \theta = 1 - \cos^2 \theta = 1 - \frac{4}{9} = \frac{5}{9}. \quad \therefore \sin \theta = \pm \frac{\sqrt{5}}{3}.$$

$$\tan \theta = \frac{\sin \theta}{\cos \theta} = \pm \frac{\sqrt{5}}{2}.$$



## Examples II

Prove the following identities (*Ex. 1 to 24*) :—

- ✓1.  $\frac{\sin A + \cos A}{\sec A + \operatorname{cosec} A} = \sin A \cos A.$
- ✓2.  $\cot \theta + \tan \theta = \sec \theta \operatorname{cosec} \theta.$
- ✓3.  $\frac{1}{1 + \tan A} = \frac{\cot A}{1 + \cot A}.$
- ✓4.  $\operatorname{cosec}^6 A - \cot^6 A = 1 + 3 \operatorname{cosec}^2 A \cot^2 A.$
- ✓5.  $\cos^6 A + \sin^6 A = 1 - 3 \sin^2 A \cos^2 A.$
- ✓6.  $\frac{1}{\cos^2 A} - \operatorname{cosec}^2 A - 1 = 1.$
- ✓7.  $\cos A + \tan A \sin A = \sec A.$
- ✓8.  $\sec^4 A + \tan^4 A = 1 + 2 \sec^2 A \tan^2 A.$
- ✓9.  $\frac{1 + 3 \cos \theta - 4 \cos^3 \theta}{1 - \cos \theta} = (1 + 2 \cos \theta)^2.$
10.  $(\cot \theta + \operatorname{cosec} \theta)^2 = \frac{1 + \cos \theta}{1 - \cos \theta}.$
11.  $\frac{1 + \tan^2 \theta}{1 + \cot^2 \theta} = \left( \frac{1 - \tan \theta}{1 - \cot \theta} \right)^2.$
- ✓12.  $\frac{\tan^2 \alpha - \cot^2 \alpha}{1 + \cot^2 \alpha} = \frac{\sin^2 \alpha - \cos^2 \alpha}{\cos^2 \alpha}.$
- ✓13.  $1 + \tan \theta + \sec \theta = \frac{2}{1 + \cot \theta - \operatorname{cosec} \theta}.$
- ✓14.  $\frac{\tan \theta + \sec \theta - 1}{\tan \theta - \sec \theta + 1} = \frac{1 + \sin \theta}{\cos \theta}.$
- ✓15.  $\frac{\sin A - 2 \sin^3 A}{2 \cos^3 A - \cos A} = \tan A.$
16.  $\sqrt{\frac{1 + \sin \theta}{1 - \sin \theta}} - \sec \theta = \sec \theta - \sqrt{\frac{1 - \sin \theta}{1 + \sin \theta}}.$

$$17. \frac{\operatorname{cosec} A + \cot A}{\operatorname{cosec} A - \cot A} = \frac{\sin^2 A}{(1 - \cos A)^2}.$$

$$18. (1 + \sin A + \cos A)^2 = 2(1 + \sin A)(1 + \cos A).$$

$$19. \frac{\sec \theta + \tan \theta}{\operatorname{cosec} \theta + \cot \theta} - \frac{\sec \theta - \tan \theta}{\operatorname{cosec} \theta - \cot \theta} = 2(\sec \theta - \operatorname{cosec} \theta).$$

$$20. \frac{1}{1 + \sin^2 \theta} + \frac{1}{1 + \operatorname{cosec}^2 \theta} = 1.$$

$$21. \frac{\sin^3 \alpha + \cos^3 \alpha}{\sin \alpha + \cos \alpha} + \frac{\sin^3 \alpha - \cos^3 \alpha}{\sin \alpha - \cos \alpha} = 2.$$

$$22. \frac{\tan \theta}{\sec \theta - 1} - \frac{\sin \theta}{1 + \cos \theta} = 2 \cot \theta.$$

$$23. \frac{\cos \theta + \cos \phi}{\sin \theta - \sin \phi} = \frac{\sin \theta + \sin \phi}{\cos \phi - \cos \theta}.$$

$$24. 1 + 4 \operatorname{cosec}^2 \theta \cot^2 \theta = (\operatorname{cosec}^2 \theta + \cot^2 \theta)^2.$$

25. Express  $1 - 2 \sin \theta \cos \theta$  as a perfect square.  $\sim$

26. Express  $2 \sec^2 \theta - \sec^4 \theta - 2 \operatorname{cosec}^2 \theta + \operatorname{cosec}^4 \theta$  in terms of  $\tan \theta$ .

27. Prove that

$$(\sin \alpha \cos \beta + \cos \alpha \sin \beta)(\sin \alpha \cos \beta - \cos \alpha \sin \beta) = \sin^2 \alpha - \sin^2 \beta.$$

28. If  $\sin A + \sin^2 A = 1$ , then  $\cos^2 A + \cos^4 A = 1$ .

29. (i) If  $\sin \theta - \cos \theta = 0$ , prove that  $\sec \theta = \pm \sqrt{2}$ .

(ii) If  $7 \sin^2 \theta + 3 \cos^2 \theta = 4$ , show that  $\tan \theta = \pm \frac{1}{\sqrt{3}}$ .

(iii) If  $3 \sin \theta + 4 \cos \theta = 5$  show that  $\sin \theta = \frac{3}{5}$ .

30. If  $\tan \theta + \sec \theta = x$ , show that  $\sin \theta = \frac{x^2 - 1}{x^2 + 1}$ .

31. If  $\tan \theta = \frac{a}{b}$ , find the value of  $\frac{a \sin \theta - b \cos \theta}{a \sin \theta + b \cos \theta}$ .

32. If  $1 + 4x^2 = 4x \sec A$ , prove that  
 $\sec A + \tan A = 2x$  or  $1/2x$ .
33. Express  $\sin \alpha$  in terms of  $\sec \alpha$ , and  $\sec \theta$  in terms of  $\cot \theta$ .
34. Given  $\sin \theta = \frac{2}{3}$ ,  $\cos \phi = \frac{1}{3}$ , where  $\theta$  and  $\phi$  are acute angles, find the value of  $\frac{\tan \theta - \tan \phi}{1 + \tan \theta \tan \phi}$ .
35. If  $\cos \alpha + \sin \alpha = \sqrt{2} \cos \alpha$ , prove that  
 $\cos \alpha - \sin \alpha = \sqrt{2} \sin \alpha$ .
36. If  $\tan A = \frac{1}{\sqrt{3}}$ , find  $\frac{\operatorname{cosec}^2 A - \sec^2 A}{\operatorname{cosec}^2 A + \sec^2 A}$ .
37. If  $1 + \sin^2 A = 3 \sin A \cos A$ , find  $\tan A$ .
38. If  $\tan \theta + \sin \theta = m$ ,  $\tan \theta - \sin \theta = n$ , prove that  
 $m^2 - n^2 = 4 \sqrt{mn}$ .
39. If  $(a^2 - b^2) \sin \theta + 2ab \cos \theta = a^2 + b^2$ , find  $\tan \theta$  and  $\operatorname{cosec} \theta$ .
40. If  $\tan \theta = \frac{\sin \alpha - \cos \alpha}{\sin \alpha + \cos \alpha}$ , prove that  
 $\sqrt{2} \cos \theta = \sin \alpha + \cos \alpha$ .
41. Given  $\tan^2 \theta = 1 - e^2$ , show that  
 $\sec \theta + \tan^3 \theta \operatorname{cosec} \theta = (2 - e^2)^{\frac{5}{2}}$ .
42. If  $x$  and  $y$  are two unequal real quantities, show that the equations (i)  $\sin^2 \theta = \frac{(x+y)^2}{4xy}$  and (ii)  $\cos \theta = x + \frac{1}{x}$  are both impossible.
43. Eliminate  $\theta$  between  
 (i)  $x = a \cos \theta$ ,  $y = b \sin \theta$ .  
 (ii)  $x = c (\sec \theta + \tan \theta)$ ,  $y = c (\sec \theta - \tan \theta)$ .  
 (iii)  $a \cos \theta + b \sin \theta + c = 0$ ,  $a' \cos \theta + b' \sin \theta + c' = 0$ .  
 (vi)  $a \tan^2 \theta + b \tan \theta + c = a' \cot^2 \theta + b' \cot \theta + c' = 0$ .

## Examples II(A)

Prove the following identities (*Ex. 1 to 18*):—

1.  $\frac{\tan^3 a}{1 + \tan^2 a} + \frac{\cot^3 a}{1 + \cot^2 a} = \frac{1 - 2 \sin^2 a \cos^2 a}{\sin a \cos a}$ .
2.  $(\tan \theta + \cot \theta + \sec \theta)(\tan \theta + \cot \theta - \sec \theta) = \operatorname{cosec}^2 \theta$ .
3.  $\sin \theta(1 + \tan \theta) + \cos \theta(1 + \cot \theta) = \sec \theta + \operatorname{cosec} \theta$ .  
[C. U. 1935.]
4.  $(1 + \sin a - \cos a)^2 + (1 - \sin a + \cos a)^2 = 4(1 - \sin a \cos a)$ .
5.  $\sin^6 a + \sin^4 a \cos^2 a - \sin^2 a \cos^4 a - \cos^6 a = \sin^2 a - \cos^2 a$ .
6.  $3(\sin \theta + \cos \theta) - 2(\sin^3 \theta + \cos^3 \theta) = (\sin \theta + \cos \theta)^3$ .
7.  $\frac{1}{\operatorname{cosec} \theta - \cot \theta} - \frac{1}{\sin \theta} = \frac{1}{\sin \theta} - \frac{1}{\operatorname{cosec} \theta + \cot \theta}$ .
8.  $\frac{\cos x}{\sin x + \cos y} + \frac{\cos y}{\sin y - \cos x} = \frac{\cos x}{\sin x - \cos y} + \frac{\cos y}{\sin y + \cos x}$ .
9.  $(\sin \theta + \operatorname{cosec} \theta)^2 + (\cos \theta + \sec \theta)^2 = \tan^2 \theta + \cot^2 \theta + 7$ .
10.  $(\sec \theta - \cos \theta)(\operatorname{cosec} \theta - \sin \theta)(\tan \theta + \cot \theta) = 1$ .
11.  $\frac{1 + (\operatorname{cosec} x \tan y)^2}{1 + (\operatorname{cosec} z \tan y)^2} = \frac{1 + (\cot x \sin y)^2}{1 + (\cot z \sin y)^2}$ .
12.  $\sec^5 a \operatorname{cosec}^3 a - 3 \sec a \operatorname{cosec} a = \tan^3 a + \cot^3 a$ .
13.  $\sin^6 A - \cos^6 A = (\sin A + \cos A)(\sin A - \cos A) \times (1 + \sin A \cos A)(1 - \sin A \cos A)$ .
14.  $\frac{\tan a}{(1 + \tan^2 a)^2} + \frac{\cot a}{(1 + \cot^2 a)^2} = \sin a \cos a$ .
15.  $\sin^3 \theta \tan \theta - \cos^3 \theta \cot \theta + \sec \theta \operatorname{cosec} \theta = 2 \tan \theta$ .
16.  $\frac{\cos^3 A - \sin^3 A}{\sin A \cos^2 A - \cos A \sin^2 A} = \operatorname{cosec} A + \sec A$ .
17.  $\frac{\tan^3 A + \cot^3 A}{\tan^3 A - \cot^3 A} = \frac{\sin^4 A + \cos^4 A}{\sin^2 A - \cos^2 A}$ .

$$18. (\sin \alpha \cos \beta - \cos \alpha \sin \beta)^2 + (\cos \alpha \cos \beta + \sin \alpha \sin \beta)^2 = 1.$$

$$19. \text{ If } \cos^2 A - \sin^2 A = \tan^2 B, \\ \text{ then } \cos^2 B - \sin^2 B = \tan^2 A.$$

$$20. \text{ If } \sin^4 x + \sin^2 x = 1, \text{ then } \tan^4 x - \tan^2 x = 1.$$

21. Show that the difference between  $3 \sin^4 \theta - 2 \sin^6 \theta$  and  $2 \cos^6 \theta - 3 \cos^4 \theta$  is the same for all values of  $\theta$ .

$$(22.) \text{ If } x = \frac{1 + \sin \theta}{\cos \theta}, \text{ show that } \frac{1}{x} = \frac{1 - \sin \theta}{\cos \theta}.$$

$$23. \text{ If } \tan^2 A = 1 + 2 \tan^2 B, \text{ show that } \cos^2 B = 2 \cos^2 A. \checkmark$$

$$24. \text{ If } \sin \alpha + \cos \alpha = 1, \text{ then } \sin \alpha - \cos \alpha = \pm 1.$$

$$25. \text{ If } a \cos \theta - b \sin \theta = c, \text{ then show that} \\ a \sin \theta + b \cos \theta = \pm \sqrt{a^2 + b^2 - c^2}.$$

$$26. \text{ If } (1 + \sin x)(1 + \sin y)(1 + \sin z) \\ = (1 - \sin x)(1 - \sin y)(1 - \sin z),$$

• prove that each is equal to  $\pm \cos x \cos y \cos z$ .

$$27. \text{ If } x \sin^3 \alpha + y \cos^3 \alpha = \sin \alpha \cos \alpha, \text{ and} \\ x \sin \alpha - y \cos \alpha = 0, \text{ then } x^2 + y^2 = 1. \quad [C. U. 1937.]$$

$$28. \text{ If } \sin A = \frac{\sin x + \sin y}{1 + \sin x \sin y}, \text{ show that}$$

$$\cos A = \pm \frac{\cos x \cos y}{1 + \sin x \sin y}.$$


$$29. (i) \text{ If } \sin \alpha + \operatorname{cosec} \alpha = 2, \\ \text{ then } \sin^n \alpha + \operatorname{cosec}^n \alpha = 2.$$

$$(ii) \text{ If } \sec \alpha = \sec \beta \sec \gamma + \tan \beta \tan \gamma, \\ \text{ then } \sec \beta = \sec \gamma \sec \alpha \pm \tan \gamma \tan \alpha.$$

$$30. \text{ If } \frac{\cos^4 x}{\cos^2 y} + \frac{\sin^4 x}{\sin^2 y} = 1, \text{ then } \frac{\cos^4 y}{\cos^2 x} + \frac{\sin^4 y}{\sin^2 x} = 1.$$

CHAPTER III  
TRIGONOMETRICAL RATIOS OF SOME  
STANDARD ANGLES

**14. Ratios of  $30^\circ$ .**



Let the angle  $XOP$ , which may be supposed to be traced out by a revolving line starting from  $OX$ , be  $30^\circ$ . Let  $PN$  be drawn perpendicular upon  $OX$  from any point  $P$  on  $OP$ . The angle  $OPN$  is then  $60^\circ$ .

Produce  $PN$  to  $Q$ , making  $NQ = NP$ . Join  $OQ$ . The triangles  $PON$  and  $QON$  are easily seen to be equal in all respects, and so  $\angle OQN = \angle OPN = 60^\circ$ . Hence the triangle  $OPQ$  is equilateral, and so  $OP = PQ =$  double of  $PN$ .

Hence in the above figure if  $PN = a$ , then  $OP = 2a$  and so  $ON = \sqrt{OP^2 - PN^2} = \sqrt{4a^2 - a^2} = \sqrt{3}a$ . The sides  $ON$ ,  $PN$ , and  $OP$  are all positive in this case, since the angle is acute.

Hence

$$\sin 30^\circ = \sin PON = \frac{PN}{OP} = \frac{a}{2a} = \frac{1}{2}$$

$$\cos 30^\circ = \frac{ON}{OP} = \frac{\sqrt{3}a}{2a} = \frac{\sqrt{3}}{2}$$


$$\tan 30^\circ = \frac{PN}{ON} = \frac{1}{\sqrt{3}}$$

$$\cot 30^\circ = \frac{ON}{PN} = \sqrt{3}$$

$$\operatorname{cosec} 30^\circ = \frac{1}{\sin 30^\circ} = 2$$

$$\sec 30^\circ = \frac{1}{\cos 30^\circ} = \frac{2}{\sqrt{3}}$$

### 15. Ratios of $45^\circ$ .



Let  $\angle XOP = 45^\circ$ .  $PN$  is perpendicular on  $OX$ . In the right-angled triangle  $PON$ ,  $\angle PON = 45^\circ$ .

Therefore,  $\angle OPN$  is also  $45^\circ$  and so  $ON = PN = a$  suppose. Then  $OP = \sqrt{ON^2 + PN^2} = \sqrt{a^2 + a^2} = a\sqrt{2}$ .


Hence

$$\sin 45^\circ = \frac{PN}{OP} = \frac{1}{\sqrt{2}}$$

$$\cos 45^\circ = \frac{ON}{OP} = \frac{1}{\sqrt{2}}$$

$$\tan 45^\circ = \frac{PN}{ON} = 1$$

$$\sec 45^\circ = \operatorname{cosec} 45^\circ = \sqrt{2}, \cot 45^\circ = 1.$$

16. Ratios of  $60^\circ$ .

Let  $\angle XOP = 60^\circ$ . Now  $PN$  being perpendicular upon  $OX$ , along  $NX$  cut off  $NQ = ON$ . Join  $PQ$ . Then the two triangles  $OPN$  and  $QPN$  are easily seen to be congruent. Hence  $\angle PQN = \angle PON = 60^\circ$ . Thus the triangle  $POQ$  is equilateral, and so  $OP = OQ = \text{double of } ON$ .

If  $ON = a$ , then  $OP = 2a$  and hence  $PN = \sqrt{OP^2 - ON^2} = a\sqrt{3}$ .

$$\text{Then} \quad \sin 60^\circ = \frac{PN}{OP} = \frac{\sqrt{3}}{2}$$

$$\cos 60^\circ = \frac{ON}{OP} = \frac{1}{2}$$

$$\tan 60^\circ = \frac{PN}{ON} = \sqrt{3}$$

$$\cot 60^\circ = \frac{1}{\sqrt{3}}, \quad \sec 60^\circ = 2, \quad \operatorname{cosec} 60^\circ = \frac{2}{\sqrt{3}}.$$

**Note.** It may be noted from the values of the ratios, that  $\sin 60^\circ = \cos 30^\circ$ ,  $\cos 60^\circ = \sin 30^\circ$ ,  $\tan 60^\circ = \cot 30^\circ$ ,  $\cot 60^\circ = \tan 30^\circ$ ,  $\sec 60^\circ = \operatorname{cosec} 30^\circ$ ,  $\operatorname{cosec} 60^\circ = \sec 30^\circ$ . It will be proved more generally, in the next chapter, that for any two complementary angles sine of one is the cosine of the other and *vice-versa*, tangent of one is the cotangent of the other, and secant of one is the cosecant of the other. The angle  $45^\circ$  being its own complement, therefore, it should have its sine and cosine equal to one another, as is actually seen to be the case.



17. Ratios of  $90^\circ$ .

Let  $XOP$  be an acute angle very nearly  $90^\circ$ .  $PN$  being perpendicular upon  $OX$ ,  $ON$  is extremely small, and as  $\angle XOP$  approaches more and more to  $90^\circ$ ,  $ON$  becomes smaller and smaller. The length  $OP$  may however remain finite, and  $PN$  and  $OP$  will approach each other more and more closely. Ultimately when  $\angle XOP$  becomes  $90^\circ$ ,  $OP$  and  $PN$  coincide, and  $ON$  becomes zero ultimately. Hence the ratio  $PN/OP$  becomes 1 and  $ON/OP$  becomes zero.

Thus  $\sin 90^\circ = \frac{PN}{OP}$  in the limit = 1

$\cos 90^\circ = \frac{ON}{OP}$  in the limit = 0

$\tan 90^\circ = \frac{PN}{ON}$  in the limit =  $\infty^*$  (infinity)


(since  $ON \rightarrow 0$ , whereas  $PN$  remains finite)

$\cot 90^\circ = \frac{\cos 90^\circ}{\sin 90^\circ} = \frac{0}{1} = 0$

$\operatorname{cosec} 90^\circ = 1, \sec 90^\circ = OP/ON$  in the limit =  $\infty^*$ .

\* The symbol  $\infty$  is used to denote a quantity which exceeds any positive number, however large, and does not represent a definite number.

It should be noted that in determining  $\tan 90^\circ$ , we may start with an angle  $XOP$ , slightly greater than  $90^\circ$  (i.e. in the second quadrant), and make it approach  $90^\circ$ . Then  $ON$  will be negative and  $\rightarrow 0$ , whereas  $PN$  is positive. Accordingly we may also write  $\tan 90^\circ = -\infty$ . (Thus strictly speaking, we should write  $\tan 90^\circ = \pm \infty$ .) Similar remarks apply for  $\sec 90^\circ$ ,  $\cot 0^\circ$ ,  $\operatorname{cosec} 0^\circ$ .

18. Ratios of  $0^\circ$ .

Let  $\angle XOP$  be an infinitely small positive angle, and let  $PN$  be perpendicular on  $OX$ .

Then  $PN$  is infinitely small, whereas  $OP$  is finite. Now if  $\angle XOP$  be taken less and less and ultimately becomes less than any quantity we can assign, we denote it by zero, and in this case  $PN$  practically vanishes, whereas  $OP$  and  $ON$  remaining finite, coincide. Hence the ratio  $PN/OP$  becomes ultimately zero, and  $ON/OP$  becomes 1.

$$\text{Hence,} \quad \sin 0^\circ = \frac{PN}{OP} \text{ in the limit} = 0$$

$$\cos 0^\circ = \frac{ON}{OP} \text{ in the limit} = 1$$

$$\tan 0^\circ = \frac{\sin 0^\circ}{\cos 0^\circ} = \frac{0}{1} = 0$$

$$\cot 0^\circ = \frac{ON}{PN} \text{ in the limit} = \infty^*,$$

$$\operatorname{cosec} 0^\circ = \frac{OP}{PN} \text{ in the limit} = \infty^*,$$

$$\sec 0^\circ = \frac{1}{\cos 0^\circ} = \frac{1}{1} = 1.$$

**Note.** Note that  $0^\circ$  and  $90^\circ$  being complementary,  
 $\sin 0^\circ = \cos 90^\circ = 0$ ,  $\cos 0^\circ = \sin 90^\circ = 1$ , etc.

19. As the ratios of the standard angles  $0^\circ$ ,  $30^\circ$ ,  $45^\circ$ ,  $60^\circ$  and  $90^\circ$  are very often used, they should be remembered very

---

\* See foot note of Art. 17.

carefully. The first three ratios are given in the tabulated form below. The other three are reciprocals to these.

angle	sine	cosine	tangent
$0^\circ$ or $0^\circ$	0	1	0
$30^\circ$ or $\frac{\pi}{6}$	$\frac{1}{2}$	$\frac{\sqrt{3}}{2}$	$\frac{1}{\sqrt{3}}$
$45^\circ$ or $\frac{\pi}{4}$	$\frac{1}{\sqrt{2}}$	$\frac{1}{\sqrt{2}}$	1
$60^\circ$ or $\frac{\pi}{3}$	$\frac{\sqrt{3}}{2}$	$\frac{1}{2}$	$\sqrt{3}$
$90^\circ$ or $\frac{\pi}{2}$	1	0	$\pm \infty$

**Note.** The following device may be of use in remembering the *sines* and *cosines* of standard angles. The sines of the angles  $0^\circ$ ,  $30^\circ$ ,  $45^\circ$ ,  $60^\circ$ ,  $90^\circ$  are respectively the square roots of the fractions

$$\frac{0}{1}, \frac{1}{2}, \frac{1}{2}, \frac{3}{4}, \frac{4}{4}$$

and cosines of these angles are the square roots from right to left.

## 20. Examples worked out.

**Ex. 1.** If  $\theta = 30^\circ$ , verify that  $\cos 2\theta = \frac{1 - \tan^2 \theta}{1 + \tan^2 \theta}$ .

Hence  $\cos 2\theta = \cos 60^\circ = \frac{1}{2}$ . Also  $\tan \theta = \tan 30^\circ = \frac{1}{\sqrt{3}}$ .

$$\therefore \frac{1 - \tan^2 \theta}{1 + \tan^2 \theta} = \frac{1 - \frac{1}{3}}{1 + \frac{1}{3}} = \frac{1}{2}. \quad \text{Hence } \cos 2\theta = \frac{1 - \tan^2 \theta}{1 + \tan^2 \theta}$$

**Ex. 2.** Verify that

$$\sin 30^\circ = \sin 60^\circ \cos 30^\circ - \cos 60^\circ \sin 30^\circ.$$

The right-hand side, on substitution of the values,

$$= \frac{\sqrt{3}}{2} \cdot \frac{\sqrt{3}}{2} - \frac{1}{2} \cdot \frac{1}{2} = \frac{3}{4} - \frac{1}{4} = \frac{2}{4} = \sin 30^\circ.$$

Hence the result.

**Ex. 3.** Solve for  $\theta$ , where  $\theta$  is a positive acute angle, given  $\operatorname{cosec} \theta \cot \theta = 2\sqrt{3}$ .

From the given equation,  $\frac{1}{\sin \theta} \cdot \frac{\cos \theta}{\sin \theta} = 2\sqrt{3}$ ,

$$\text{or, } \cos \theta = 2\sqrt{3} \sin^2 \theta = 2\sqrt{3}(1 - \cos^2 \theta),$$

$$\text{whence, } 2\sqrt{3} \cos^2 \theta + \cos \theta - 2\sqrt{3} = 0$$

$$\text{giving } \cos \theta = \frac{-1 \pm \sqrt{1+48}}{4\sqrt{3}} = \frac{-1 \pm 7}{4\sqrt{3}}.$$

Since  $\theta$  is a positive acute angle,  $\cos \theta$  is positive, and so rejecting the negative value,

$$\cos \theta = \frac{6}{4\sqrt{3}} = \frac{\sqrt{3}}{2} = \cos 30^\circ. \quad \therefore \theta = 30^\circ \text{ i.e., } \frac{\pi}{6}.$$

### Examples III

Verify the results (Ex. 1 to 6) :—

$$1. \quad 1 - 2 \sin^2 30^\circ = 2 \cos^2 30^\circ - 1 = \cos 60^\circ.$$

$$2. \quad \frac{2 \tan 30^\circ}{1 - \tan^2 30^\circ} = \sqrt{3}.$$

$$\cos \frac{\pi}{3} \cos \frac{\pi}{4} + \sin \frac{\pi}{3} \sin \frac{\pi}{4} = \sin \frac{\pi}{4} \cos \frac{\pi}{6} + \cos \frac{\pi}{4} \sin \frac{\pi}{6}.$$

$$4. \text{ (i) } \tan(A+B) = \frac{\tan A + \tan B}{1 - \tan A \tan B};$$

$$\text{(ii) } \cos A = \cos^2 B - \sin^2 B, \\ \text{where } A = 60^\circ, B = 30^\circ.$$

5.  $\sin 3A = 3 \sin A - 4 \sin^3 A$ , where  $A = \frac{\pi}{6}$ .
6.  $\operatorname{cosec}^2 45^\circ \cdot \sec^2 30^\circ (\sin^2 30^\circ + 4 \cot^2 45^\circ - \sec^2 60^\circ) = \frac{1}{3}$ .
7. If  $\tan^2 \frac{\pi}{4} - \cos^2 \frac{\pi}{3} = x \sin \frac{\pi}{4} \cos \frac{\pi}{4} \tan \frac{\pi}{3}$ , find  $x$ .
8. If  $\theta$  be a positive acute angle, find  $\theta$ , when
- $2 \sin^2 \theta = 3 \cos \theta$ ;
  - $\tan \theta + \cot \theta = 2$ ;
  - $\operatorname{cosec}^2 \theta + 5 = 3 \sqrt{3} \cot \theta$ ;
  - $\sin \theta + \cos \theta = \sqrt{2}$ ;
  - $2(\cos^2 \theta - \sin^2 \theta) = 1$ ;
  - $6 \sin^3 \theta - 11 \sin \theta + 4 = 0$ ;
  - $\frac{\sin \theta - \cos \theta}{\sin \theta + \cos \theta} = \frac{1 - \sqrt{3}}{1 + \sqrt{3}}$ .
9. Given  $\theta$  and  $\phi$  to be positive acute angles, and  $\tan(\theta + \phi) = \sqrt{3}$ ,  $\tan(\theta - \phi) = 1$ , determine  $\theta$  and  $\phi$ .
10. Find  $\alpha$  and  $\beta$  ( $\alpha$  and  $\beta$  being positive acute angles), if
- $$\sin(2\alpha - \beta) = 1,$$
- and  $\cos(\alpha + \beta) = \frac{1}{2}$ .
11. Find  $A, B, C$  ( $A, B, C$  being positive acute angles), if
- $$\sin(B + C - A) = 1,$$
- $$\cos(C + A - B) = 1,$$
- and  $\tan(A + B - C) = 1$ .
12. Find the numerical values of :--
- $\cot^2 \frac{\pi}{6} - 2 \cos^2 \frac{\pi}{3} - \frac{3}{4} \sec^2 \frac{\pi}{4} - 4 \sec^2 \frac{\pi}{6}$ ;
  - $3 \tan^2 45^\circ - \sin^2 60^\circ - \frac{1}{3} \cot^2 30^\circ + \frac{1}{3} \sec^2 45^\circ$ .

CHAPTER IV  
TRIGONOMETRICAL RATIOS OF ANGLES ASSOCIATED  
WITH A GIVEN ANGLE  $\theta$

**21. Ratios of the angle  $(-\theta)$  in terms of those of  $\theta$ ,  $\theta$  having any magnitude.**




Fig. (i)




Fig. (ii)




Fig. (iii)




Fig. (iv)

Let the  $\angle XOP$  be  $\theta$  and the  $\angle XOQ$  described clockwise be  $-\theta$ . From any point  $P$  on  $OP$  draw  $PN$  perpendicular to  $OX$  [or  $OX'$  as in Figs. (ii) and (iii)], and produce it to meet  $OQ$  at  $Q$  say.

Now  $\angle XOP$  (measured anti-clockwisely) being equal to  $\angle XOQ$  (measured clockwisely),  $\angle PON = \angle QON$  in magnitude in all the figures, and therefore the two rt-angled triangles  $PON$  and  $QON$  are congruent. The corresponding sides are therefore equal in magnitude. Considering the signs of these sides according to the usual convention, we get in all the figures,

$$QN = -PN, \text{ and } OQ = OP$$

(both  $OP$  and  $OQ$  being always considered positive).

Hence, from definition,

$$\sin(-\theta) = \frac{QN}{OQ} = \frac{-PN}{OP} = -\sin \theta$$

$$\cos(-\theta) = \frac{ON}{OQ} = \frac{ON}{OP} = \cos \theta$$

$$\tan(-\theta) = \frac{QN}{ON} = \frac{-PN}{ON} = -\tan \theta$$

and the reciprocals of these give,

$$\operatorname{cosec}(-\theta) = -\operatorname{cosec} \theta,$$

$$\sec(-\theta) = \sec \theta,$$

$$\cot(-\theta) = -\cot \theta.$$

## 22. Ratios of $(90^\circ - \theta)$ .

Let the  $\angle XOP$  traced out by a revolving line be  $\theta$ , and let another revolving line, starting from  $OX$  trace out the angle  $XOY = 90^\circ$  and then revolve back, tracing out  $\angle YOQ = \theta$  in the clockwise direction, so that  $\angle XOQ = 90^\circ - \theta$ .

Take two equal lengths  $OP$  and  $OQ$  along  $OP$  and  $OQ$  respectively, and draw  $PN$  and  $QM$  perpendiculars on  $OX$ .

If  $OP$  be in the first or third quadrant as in Fig. (i) and Fig. (iii),  $OQ$  also lies in the same quadrant. If  $OP$  lies in the second quadrant as in Fig. (ii),  $OQ$  lies in the fourth quadrant; and if  $OP$  lies in the fourth,  $OQ$  lies in the

second, as in Fig. (iv). Now,  $\angle XOQ$  being equal to  $\angle YOQ$  in magnitude,  $\angle PON = \angle OQM$ , and since  $OP = OQ$ , the




Fig. (i)




Fig. (ii)




Fig. (iii)




Fig. (iv)

two rt.-angled triangles  $PON$ ,  $OQM$  are congruent. The corresponding sides are therefore equal in magnitude. Considering signs as well, we get in all the figures,

$$QM = ON, OM = PN, OQ = OP.$$

Hence, from definition,

$$\sin(90^\circ - \theta) = \sin \angle XOQ = \frac{QM}{OQ} = \frac{ON}{OP} = \cos \theta$$

$$\cos(90^\circ - \theta) = \frac{OM}{OQ} = \frac{PN}{OP} = \sin \theta$$

$$\tan(90^\circ - \theta) = \frac{QM}{OM} = \frac{ON}{PN} = \cot \theta$$



The reciprocals of these are

$$\operatorname{cosec} (90^\circ - \theta) = \sec \theta,$$

$$\sec (90^\circ - \theta) = \operatorname{cosec} \theta,$$

$$\cot (90^\circ - \theta) = \tan \theta.$$

**Obs.** The angle  $(90^\circ - \theta)$  is the complement of  $\theta$ , and we derive the result that for a pair of complementary angles, sine of one is the cosine of the other and *vice versa*, tangent of one is the cotangent of the other and secant of one is the cosecant of the other. This was verified in the last chapter in connection with the complementary pairs  $30^\circ$  and  $60^\circ$ , as also  $0^\circ$  and  $90^\circ$ .

### 23. Ratios of $(90^\circ + \theta)$ .




Fig. (i)




Fig. (ii)




Fig. (iii)




Fig. (iv)

Let a revolving line, starting from  $OX$ , trace out an  $\angle XOP = \theta$ , and further trace out an  $\angle POQ = 90^\circ$ , so that  $\angle XOQ = 90^\circ + \theta$ .

Cut off  $OP = OQ$  along  $OP$  and  $OQ$  respectively and let  $PN, QM$  be perpendiculars on  $OX$  (produced where necessary).

Now  $OQ$  being perpendicular to  $OP$ , the  $\angle PON =$  the complement of  $\angle QOM = \angle OQM$  in magnitude, and since  $OP = OQ$ , the two right-angled triangles  $OPN$  and  $OQM$  are congruent. The corresponding sides are therefore equal. Considering signs as well, we get, for all the figures,

$$QM = ON, OM = -PN, OQ = OP.$$

Hence from definition,

$$\sin(90^\circ + \theta) = \sin XOQ = \frac{QM}{OQ} = \frac{ON}{OP} = \cos \theta$$

$$\cos(90^\circ + \theta) = \frac{OM}{OQ} = \frac{-PN}{OP} = -\sin \theta$$

$$\tan(90^\circ + \theta) = \frac{QM}{OM} = \frac{ON}{-PN} = -\cot \theta$$

and considering their reciprocals,

$$\operatorname{cosec}(90^\circ + \theta) = \sec \theta,$$

$$\sec(90^\circ + \theta) = -\operatorname{cosec} \theta,$$

$$\cot(90^\circ + \theta) = -\tan \theta.$$

#### 24. Ratios of $(180^\circ - \theta)$ .




Fig. (i)




Fig. (ii)

Let  $\angle XOP = \theta$  be traced out by a revolving line, and let another revolving line, starting from  $OX$ , trace out an angle  $180^\circ$  coming up to  $OX'$  and then revolve back and describe an angle  $X'OQ = \theta$ , so that  $\angle XOQ = 180^\circ - \theta$ .

Two figures are given here, one with  $OP$  in the first quadrant and another with  $OP$  in the third quadrant. The two other figures may easily be drawn by the students.

Now cut off  $OP = OQ$ , and draw  $PN$  and  $QM$  perpendiculars on  $OX$  (or  $OX'$  as the case may be). Then  $\angle PON = \angle QOM$  in magnitude, and  $OP = OQ$ . Hence the right-angled triangles  $PON$  and  $QOM$  are congruent, and so have their corresponding sides equal in magnitude. Taking into consideration the signs, we get for all the figures,

$$QM = PN, OM = -ON, OQ = OP.$$

Hence for all values of  $\theta$ ,

$$\sin(180^\circ - \theta) = \sin XOQ = \frac{QM}{OQ} = \frac{PN}{OP} = \sin \theta$$

$$\cos(180^\circ - \theta) = \frac{OM}{OQ} = \frac{-ON}{OP} = -\cos \theta$$

$$\tan(180^\circ - \theta) = \frac{QM}{OM} = \frac{PN}{-ON} = -\tan \theta$$

and so taking reciprocals,

$$\operatorname{cosec}(180^\circ - \theta) = \operatorname{cosec} \theta,$$

$$\sec(180^\circ - \theta) = -\sec \theta,$$

$$\cot(180^\circ - \theta) = -\cot \theta.$$

**Note.** The first two formulæ may be expressed in the form "*sines of supplementary angles are equal, and cosines of supplementary angles are equal in magnitude but opposite in sign.*"

## 25. Ratios of $(180^\circ + \theta)$ .

Let a revolving line starting from  $OX$ , trace out an angle  $XOP = \theta$ , and further trace out an angle  $POQ = 180^\circ$ , so that  $\angle XOQ = 180^\circ + \theta$ .

$OP$  and  $OQ$  are then in one straight line.




Fig. (i)

Fig. (ii)

Cut off  $OP = OQ$ , and draw  $PN$  and  $QM$  perpendiculars on  $XOX'$ .

Two figures are given here with  $OP$  in the first and fourth quadrants, and the other two may be similarly drawn.

Now  $POQ$  being a straight line in this case,  $\angle PON = \angle QOM$  in magnitude. Also,  $OP = OQ$ . Hence the right-angled triangles  $PON$  and  $QOM$  are congruent and have their corresponding sides equal in magnitude. Considering signs, we get in all cases,

$$QM = -PN, OM = -ON, OQ = OP.$$

Thus for all values of  $\theta$ ,

$$\sin(180^\circ + \theta) = \sin XOQ = \frac{QM}{OQ} = \frac{-PN}{OP} = -\sin \theta$$

$$\cos(180^\circ + \theta) = \frac{OM}{OQ} = \frac{-ON}{OP} = -\cos \theta$$

$$\tan(180^\circ + \theta) = \frac{QM}{OM} = \frac{-PN}{-ON} = \frac{PN}{ON} = \tan \theta$$

and so,

$$\operatorname{cosec} (180^\circ + \theta) = -\operatorname{cosec} \theta,$$

$$\sec (180^\circ + \theta) = -\sec \theta,$$

$$\cot (180^\circ + \theta) = \cot \theta.$$

## 26. Ratios of $(270^\circ - \theta)$ .




Fig. (i)




Fig. (ii)

Let  $\angle XOP = \theta$  be traced out by a revolving line, and let another revolving line trace out an angle  $\angle XOY' = 270^\circ$ , thereby coming up to the position  $OY'$ , and then revolve back, tracing out an angle  $\angle Y'OQ = \theta$ , so that  $\angle XOQ = 270^\circ - \theta$ .

Two figures are given here with  $OP$  in the first and third quadrants. The other two may be drawn similarly.

Cut off  $OP = OQ$  and draw  $PN$ ,  $QM$  perpendiculars on  $XOX'$ .

Since  $\angle XOP = \angle Y'OQ$  in magnitude, we easily derive that  $\angle PON = \angle OQM$  in magnitude. Also  $OP = OQ$ . Hence the two right-angled triangles  $OPN$  and  $OQM$  are congruent. Considering signs, we get for all the figures,

$$QM = -ON, OM = -PN, OQ = OP.$$

Hence, for all values of  $\theta$ ,

$$\sin (270^\circ - \theta) = \sin \angle XOQ = \frac{QM}{OQ} = \frac{-ON}{OP} = -\cos \theta$$

$$\cos (270^\circ - \theta) = \frac{OM}{OQ} = \frac{-PN}{OP} = -\sin \theta$$

$$\tan (270^\circ - \theta) = \frac{QM}{OM} = \frac{-ON}{-PN} = \frac{ON}{PN} = \cot \theta ;$$

and thus,

$$\operatorname{cosec} (270^\circ - \theta) = -\sec \theta,$$

$$\sec (270^\circ - \theta) = -\operatorname{cosec} \theta,$$

$$\cot (270^\circ - \theta) = \tan \theta.$$

## 27. Ratios of $(270^\circ + \theta)$ .

We may proceed geometrically as in the previous cases. Otherwise we may proceed as follows :

$$\sin (270^\circ + \theta) = \sin (180^\circ + 90^\circ + \theta) = -\sin (90^\circ + \theta) \quad \left[ \begin{array}{l} \text{from } \S 25 \\ \dots \dots \dots \text{from } \S 28 \end{array} \right]$$

$$\cos (270^\circ + \theta) = \cos (180^\circ + 90^\circ + \theta) = -\cos (90^\circ + \theta) \\ = -(-\sin \theta) = \sin \theta$$

$$\tan (270^\circ + \theta) = \frac{\sin (270^\circ + \theta)}{\cos (270^\circ + \theta)} = \frac{-\cos \theta}{\sin \theta} = -\cot \theta ;$$

and hence,

$$\operatorname{cosec} (270^\circ + \theta) = -\sec \theta,$$

$$\sec (270^\circ + \theta) = \operatorname{cosec} \theta,$$

$$\cot (270^\circ + \theta) = -\tan \theta.$$

**Note.** The ratios of  $180^\circ - \theta$ ,  $180^\circ + \theta$ ,  $270^\circ - \theta$  can also be similarly deduced from the formulæ for ratios of  $90^\circ \pm \theta$ .

## 28. Ratios of $(360^\circ - \theta)$ , $(360^\circ + \theta)$ and $(n.360^\circ \pm \theta)$ .

It has already been remarked in Art. 2, Chapter I, that angles which differ by complete multiples of  $360^\circ$ , i.e. by an exact number of complete revolutions, have the final positions of the revolving lines coincident, if the initial lines are

the same. Hence all the trigonometrical ratios of two such angles must be identical in magnitude as well as in sign.

Thus trigonometrical ratios of  $360^\circ - \theta$  must be same as those of  $-\theta$ . Hence,

$$\sin(360^\circ - \theta) = \sin(-\theta) = -\sin \theta$$

$$\cos(360^\circ - \theta) = \cos(-\theta) = \cos \theta$$

$$\tan(360^\circ - \theta) = \tan(-\theta) = -\tan \theta, \text{ etc.}$$

Trigonometrical ratios of  $360^\circ + \theta$ , or of  $360^\circ \times n \pm \theta$ , where  $n$  is an integer, positive or negative, must similarly be same as those of  $\theta$ , or of  $\pm \theta$ .

Thus in *determining trigonometrical ratios of angles, complete multiples of  $360^\circ$  (i.e.,  $2\pi$ ) may be always added or subtracted.*

29. All the above results may, for easy remembrance, be summed up in a **simple rule**.

If  $\theta$  be associated with an even multiple of  $90^\circ$  by a + or - sign, (e.g.,  $180^\circ - \theta$ ,  $180^\circ + \theta$ ,  $360^\circ - \theta$ ,  $360^\circ + \theta$ , etc.), the ratio is not altered in form (i.e., sine remains sine, cosine remains cosine, etc.). To determine the sign, assuming  $\theta$  to be acute, find out the quadrant in which the associated angle lies, and determine the sign according to the rule "all, sin, tan, cos".

If  $\theta$  be associated with an odd multiple of  $90^\circ$  by a + or - sign, (e.g.,  $90^\circ - \theta$ ,  $90^\circ + \theta$ ,  $270^\circ - \theta$ ,  $270^\circ + \theta$ , etc.), the ratio is altered (sine becomes cosine, cosine becomes sine, tangent becomes cotangent, etc.). Moreover, the sign of the result is determined as in the previous paragraph.

**Example.** Consider formulae for  $\tan(270^\circ - \theta)$  and  $\sec(180^\circ + \theta)$ .

$$270^\circ - \theta = 3 \cdot 90^\circ - \theta \text{ (multiple of } 90^\circ \text{ is odd).}$$

Hence the ratio will be altered,  $\tan$  changing into  $\cot$ . Moreover,  $\theta$  being assumed acute (whether it actually is so or not, it does not matter),  $270^\circ - \theta$  falls in the third quadrant, where  $\tan$  is positive.

Hence,  $\tan (270^\circ - \theta) = +\cot \theta$ .

$180^\circ + \theta$  has got  $\theta$  associated with even multiple of  $90^\circ$ . Hence the ratio does not alter in form,  $\sec$  remaining  $\sec$ . Also,  $180^\circ + \theta$  falls in the third quadrant, if  $\theta$  be assumed acute, where  $\sec$  (by the rule "all, sin, tan, cos") is negative.

Hence,  $\sec (180^\circ + \theta) = -\sec \theta$ .

**N. B.** The angle " $-\theta$ " may be written as  $0.360^\circ - \theta$ , and 0 may be considered *even* in applying the above rule.

Thus,  $\theta$  being supposed acute,  $-\theta$  falls in the fourth quadrant, where  $\cos$  and  $\sec$  only are positive. The form of the ratio not changing in this case,  $\sin (-\theta) = -\sin \theta$ ,  $\cos (-\theta) = +\cos \theta$ , etc.

### 30. Special angles (*outside the first quadrant*).

In Art. 24, putting  $\theta = 60^\circ$ ,  $45^\circ$ ,  $30^\circ$  and  $0^\circ$  respectively we can deduce the following results :

$$\sin 120^\circ = \sin 60^\circ = \frac{\sqrt{3}}{2}; \quad \cos 120^\circ = -\cos 60^\circ = -\frac{1}{2}.$$

$$\sin 135^\circ = \sin 45^\circ = \frac{1}{\sqrt{2}}; \quad \cos 135^\circ = -\cos 45^\circ = -\frac{1}{\sqrt{2}}.$$

$$\sin 150^\circ = \sin 30^\circ = \frac{1}{2}; \quad \cos 150^\circ = -\cos 30^\circ = -\frac{\sqrt{3}}{2}.$$

$$\sin 180^\circ = \sin 0^\circ = 0; \quad \cos 180^\circ = -\cos 0^\circ = -1.$$

And similarly from Arts. 27 and 28, putting  $\theta = 0$ ,

$$\sin 270^\circ = -\cos 0^\circ = -1; \quad \cos 270^\circ = \sin 0^\circ = 0.$$

$$\sin 360^\circ = \sin 0^\circ = 0; \quad \cos 360^\circ = \cos 0^\circ = 1.$$

From the above, we get,

$$\tan 180^\circ = 0; \quad \tan 270^\circ = -\infty; \quad \tan 360^\circ = 0.$$



**Examples worked out.****Ex. 1.** Find the value of  $\cot (-1575^\circ)$ .

$$\begin{aligned}\cot (-1575^\circ) &= -\cot (1575^\circ) = -\cot (4 \times 360^\circ + 135^\circ) \\ &= -\cot (135^\circ) = -\cot (180^\circ - 45^\circ) \\ &= \cot 45^\circ = 1.\end{aligned}$$

**Ex. 2.** Find the value of  $\cot \theta - \tan \theta$ , where  $\theta = \frac{17\pi}{3}$ .

$\frac{17\pi}{3} = 6\pi - \frac{\pi}{3}$ , and omitting complete multiples of  $360^\circ$  i.e., of  $2\pi$ , whereby trigonometrical ratios are not altered, we get,

$$\begin{aligned}\cot \frac{17\pi}{3} &= \cot \left( -\frac{\pi}{3} \right) = -\cot \frac{\pi}{3} = -\cot 60^\circ = -\frac{1}{\sqrt{3}}. \\ \tan \frac{17\pi}{3} &= \tan \left( -\frac{\pi}{3} \right) = -\tan \frac{\pi}{3} = -\tan 60^\circ = -\sqrt{3}. \\ \therefore \cot \theta - \tan \theta &= -\frac{1}{\sqrt{3}} + \sqrt{3} = \frac{3-1}{\sqrt{3}} = \frac{2}{\sqrt{3}} = \frac{2\sqrt{3}}{3}.\end{aligned}$$

**Ex. 3.** Prove that

$$\sin (420^\circ) \cos (390^\circ) + \cos (-300^\circ) \sin (-330^\circ) = 1.$$

$$\begin{aligned}\text{L. H. side} &= \sin (360^\circ + 60^\circ) \cos (360^\circ + 30^\circ) \\ &\quad + \cos (-360^\circ + 60^\circ) \sin (-360^\circ + 30^\circ) \\ &= \sin 60^\circ \cos 30^\circ + \cos 60^\circ \sin 30^\circ \\ &= \frac{\sqrt{3}}{2} \cdot \frac{\sqrt{3}}{2} + \frac{1}{2} \cdot \frac{1}{2} = \frac{3}{4} + \frac{1}{4} = 1.\end{aligned}$$

**Ex. 4.** Express  $\cot (-1358^\circ)$  in terms of the ratio of a positive angle less than  $45^\circ$ .

$$\begin{aligned}\cot (-1358^\circ) &= \cot (-4 \times 360^\circ + 82^\circ) \\ &= \cot 82^\circ = \cot (90^\circ - 8^\circ) \\ &= \tan 8^\circ.\end{aligned}$$

**Note.** Ratios of angles of any magnitude and sign can always be expressed in terms of a ratio of a positive angle less than  $45^\circ$ .

Ex. 5. Express

$\frac{\cos (90^{\circ} + \theta) \sec (-\theta) \tan (180^{\circ} - \theta)}{\sec (360^{\circ} + \theta) \sin (180^{\circ} + \theta) \cot (90^{\circ} - \theta)}$  in its simplest form.

The given expression

$$\begin{aligned} &= \frac{-\sin \theta \sec \theta (-\tan \theta)}{\sec \theta \cdot (-\sin \theta) \cdot \tan \theta} \\ &= -1. \end{aligned}$$

### Examples IV

1. Write down the values of  $\sin 150^{\circ}$ ,  $\cot 840^{\circ}$ ,  $\operatorname{cosec} (-660^{\circ})$  and  $\tan (-1125^{\circ})$ .

2. Find the values of  $\sin \left(-\frac{11\pi}{4}\right)$ ,  $\operatorname{cosec} \left(\frac{16\pi}{3}\right)$ ,

$$\tan \left(\frac{3\pi}{2} + \frac{\pi}{3}\right) \text{ and } \cos \left(\frac{5\pi}{2} - \frac{19\pi}{3}\right).$$

3. Evaluate  $\sin \left(-1230^{\circ}\right) - \cos \left\{\left(2n+1\right)\pi + \frac{\pi}{3}\right\}$ , where  $n$  is a negative integer.

4. Find the value of  $\sin \left\{n\pi + (-1)^n \frac{\pi}{3}\right\}$ , where  $n$  is any integer.

5. Find all the values of

$$(i) \tan \left\{\frac{n\pi}{2} + (-1)^n \frac{\pi}{4}\right\};$$

$$(ii) \operatorname{cosec} \left\{\frac{n\pi}{2} + (-1)^n \frac{\pi}{6}\right\},$$

where  $n$  is any integer.

6. Show that  $\cos \left(2m\pi \pm \frac{\pi}{3}\right)$  and  $\tan \left(m\pi + \frac{\pi}{6}\right)$  have one value each for all integral values of  $m$ .

7. Prove that,  $n$  being any integer

$$(i) \cos (n\pi + a) = (-1)^n \cos a ;$$

$$(ii) \tan (n\pi + a) = \tan a .$$

8. Prove that

$$(i) \cos \theta = -\cos (\theta - 180^\circ) ;$$

$$(ii) \tan \theta = -\cot (\theta - \frac{3}{2}\pi) .$$

9. Prove that

$$(i) \sin (780^\circ) \cos (390^\circ) - \sin (330^\circ) \cos (-300^\circ) = 1 ;$$

$$(ii) \cos 306^\circ + \cos 234^\circ + \cos 162^\circ + \cos 18^\circ = 0 ;$$

$$(iii) \frac{\sin 250^\circ + \tan 290^\circ}{\cot 200^\circ + \cos 340^\circ} = -1 .$$

10. Simplify

$$\frac{\sin^3(\pi + \theta)}{\cos^2(\frac{1}{2}\pi + \theta)} \cdot \frac{\tan(2\pi - \theta)}{\operatorname{cosec}^2 \theta} \cdot \frac{\sec^2(\pi - \theta)}{\sin(\pi - \theta)}$$

and determine its value when  $\theta = 225^\circ$ .

11. Prove that

$$\begin{aligned} \sin(\frac{1}{2}\pi + \theta) \cos(\pi - \theta) \cot(\frac{3}{2}\pi + \theta) \\ = \sin(\frac{1}{2}\pi - \theta) \sin(\frac{3}{2}\pi - \theta) \cot(\frac{1}{2}\pi + \theta) . \end{aligned}$$

12. Evaluate

$$(i) \sin^2 \frac{\pi}{4} + \sin^2 \frac{3\pi}{4} + \sin^2 \frac{5\pi}{4} + \sin^2 \frac{7\pi}{4} .$$

$$(ii) \cot \frac{\pi}{20} \cot \frac{3\pi}{20} \cot \frac{5\pi}{20} \cot \frac{7\pi}{20} \cot \frac{9\pi}{20} .$$

$$(iii) \sin x + \sin(\pi + x) + \sin(2\pi + x) + \dots \text{ to } n \text{ terms.}$$

13. If  $\tan \theta = \frac{5}{12}$  and  $\cos \theta$  is negative, find the value of

$$\frac{\sin \theta + \cos(-\theta)}{\sec(-\theta) + \tan \theta} .$$

14. An angle  $\theta$  lies between  $180^\circ$  and  $270^\circ$ , and  $\operatorname{cosec} \theta = -\frac{7}{4}$ . Find  $\cot \theta$ .

15. Express in terms of ratios of positive angles less than  $45^\circ$  ;

$$(i) \cot (-1054^\circ) ; \quad (ii) \sin (1145^\circ) ;$$

$$(iii) \sec (-1491^\circ) ; \quad (iv) \cos \frac{35\pi}{9}.$$

16. Find the value of  $\theta$  when,

$$(i) \tan \theta = -\sqrt{3} \text{ and } \theta \text{ lies between } 270^\circ \text{ and } 360^\circ ;$$

$$(ii) \cos \theta = -\frac{1}{2}, \text{ and } 450^\circ < \theta < 540^\circ.$$

17. Solve for  $\theta$ , giving all the possible values, when  $0^\circ < \theta < 360^\circ$  ;

$$(i) \cos \theta + \sqrt{3} \sin \theta = 2 ; \quad \text{[ C. U. 1936. ]}$$

$$(ii) 2 \sin^2 \theta + 3 \cos \theta = 0 ;$$

$$(iii) 3(\sec^2 \theta + \tan^2 \theta) = 5 ;$$

$$(iv) \cot \theta + \tan \theta = 2 \sec \theta ;$$

$$(v) 1 - 2 \sin \theta - 2 \cos \theta + \cot \theta = 0.$$

18. If  $A, B, C$  be angles of a triangle, show that

$$\sin (A+B) - \cos C = \cos (A+B) + \sin C.$$

19. If  $A, B, C$  be angles of a triangle, show that

$$\tan (B+C) + \tan (C+A) + \tan (A+B) \\ \tan (\pi - A) + \tan (2\pi - B) + \tan (3\pi - C) = 1.$$

20. If  $A, B, C, D$  be the angles of a quadrilateral, show that

$$\cos \frac{1}{2}(A+C) + \cos \frac{1}{2}(B+D) = 0.$$


If the quadrilateral be cyclic, then

$$\cos A + \cos B + \cos C + \cos D = 0.$$

CHAPTER V  
SIMPLE PRACTICAL APPLICATIONS OF  
TRIGONOMETRY

**31.** One of the most important applications of Trigonometry is in the determination of *heights and distances* of distant objects which are not directly measurable, by observations of angles subtended by those objects at the eye of the observer. These angles may be measured by instruments known as Sextants, or Theodolites or by other angle-measuring instruments. Thus Trigonometry plays a very important part in *land survey*. It is also extensively used by Astronomers in determining the distances of the heavenly bodies like the sun, moon and stars.

Two angles are very often used in the practical applications of Trigonometry, and they are defined as follows :—




If a horizontal line  $OX$  be drawn through  $O$ , the eye of an observer, the angle which the line joining  $O$  to a point  $P$  above  $OX$  makes with  $OX$  is called the **Angle of Elevation** or *altitude* of  $P$  as seen from  $O$ .

If  $Q$  be below the horizontal line  $OX$ , the angle  $XOQ$  measured below  $OX$  is called the **Angle of Depression** of  $Q$  as seen from  $O$ .

### 32. Illustrative Examples.

**Ex. 1.** *From a distance of 40 feet from the foot of a palm tree in a horizontal field, the angle of elevation of the top of the tree is observed to be  $60^\circ$ . Find the height of the tree.*




Let  $h$  ft. be the height of the tree  $PN$ , and  $\angle NOP$ , the angle of elevation of  $P$  as seen from  $O$ , where  $ON = 40$  ft., is  $60^\circ$ .

$$\text{Then } \frac{h}{40} = \tan PON = \tan 60^\circ = \sqrt{3};$$

$$h = 40 \sqrt{3} \text{ ft.} = 69.28 \dots \text{ft.}$$

**Ex. 2.** *From one bank of a river, the top of a building just on the opposite bank is observed to have an elevation of  $45^\circ$ . On receding 50 ft. from the bank, perpendicular to its edge, the angle of elevation becomes  $30^\circ$ . Find the breadth of the river, and the height of the building.*

$AQ$  being the breadth of the river,  $PQ$  the height of the building,  $\angle PAQ = 45^\circ$ . Also  $AB$  being 50 ft.,  $\angle PBQ = 30^\circ$ .



$$\text{Now, } \frac{BQ}{PQ} = \cot 30^\circ, \quad \frac{AQ}{PQ} = \cot 45^\circ.$$

$$\text{Hence, subtracting, } \frac{AB}{PQ} = \cot 30^\circ - \cot 45^\circ,$$

$$\text{or, } \frac{50}{PQ} = \sqrt{3} - 1;$$

$$\therefore PQ = \frac{50}{\sqrt{3} - 1} = \frac{50(\sqrt{3} + 1)}{2} = 68.3 \text{ ft. nearly.}$$

$$\text{Also } \frac{AQ}{PQ} = \cot 45^\circ = 1; \therefore AQ = PQ = 68.3 \text{ ft.}$$


Thus, the breadth of the river and the height of the building are both 68.3 ft. nearly.

**Ex. 3.** *The angles of depression and elevation of the top of a tower 50 ft. high from the top and bottom of a second tower are  $60^\circ$  and  $20^\circ$  respectively. Find the height of the second tower to the nearest foot. [ Given  $\cot 20^\circ = 2.747$ . ]*

$PQ$  is the second tower, and  $\angle XPA = 60^\circ$ ,  $\angle BQA = 20^\circ$ ,  $AB = 50$  ft.,  $AC$  is parallel to  $BQ$  or  $PX$ , so that  $\angle PAC =$  the alternate angle  $\angle XPA = 60^\circ$ .

Now  $\frac{QB}{AB} = \cot 20^\circ$ ;  $\therefore QB = AB \cot 20^\circ$ .

Also  $\frac{PC}{CA} = \tan PAC = \tan 60^\circ$ ;  
 $PC = CA \tan 60^\circ = QB \tan 60^\circ$   
 $= AB \cot 20^\circ \tan 60^\circ$ .



$\therefore$  height  $PQ = PC + CQ = PC + AB$   
 $= AB (\cot 20^\circ \tan 60^\circ + 1)$   
 $= 50(2.747 \times \sqrt{3} + 1)$   
 $= 287.8 \dots \text{ft.} = 288 \text{ ft. nearly}$

**Ex. 4.** *The elevation of a hill from a place P due East of it is  $45^\circ$ , and at a place Q due South of P, the elevation is  $30^\circ$ . If the distance PQ be 400 yds., find the height of the hill.*


A is the top of the hill, B is the point vertically below it on the ground. BP is due East, PQ is due South, so that BPQ is a right angle. Also ABP and ABQ are both right angles.

Now  $\frac{BQ}{AB} = \cot AQB = \cot 30^\circ = \sqrt{3}$ ,

and  $\frac{BP}{AB} = \cot APB = \cot 45^\circ = 1$ .



Hence,  $BQ = AB \sqrt{3}$ ,  $BP = AB$ ,



$$\text{and } PQ^2 = BQ^2 - BP^2 = AB^2(3 - 1) = 2AB^2.$$

$$\therefore AB = \frac{PQ}{\sqrt{2}} = \frac{400}{\sqrt{2}} = 200\sqrt{2} = 283 \text{ yds. nearly.}$$

### Examples V

1. From the top of a tower by the sea side, 100 feet high, it was observed that the angle of depression of the bottom of a ship at anchor was  $30^\circ$ . Find the distance of the ship from the bottom of the tower.

2. Two straight roads, which cross one another, meet a river with a straight course at angles  $60^\circ$  and  $30^\circ$  respectively. If it be 5 miles by the longer of the two roads, from the crossing to the river, how far is it by the shorter? If there be a foot-path which goes the shortest way from the crossing to the river, what is the distance by it?

3. Two poles are of equal height; a person standing midway between the line joining their bases observes the

elevations of the poles to be  $30^\circ$ . After walking 40 feet towards one of them, he observes that the same pole now subtends an angle of  $60^\circ$ . Find their height and the distance between them.

4. A straight palm tree 60 feet high, is broken by the wind but not completely separated, and its upper part meets the ground at an angle of  $30^\circ$ . Find the distance of the point where the top of the tree meets the ground, from the root, and also the height at which the tree is broken.

5. Two posts are 120 ft. apart, and the height of one is double that of the other. From the middle point of the line joining their feet, an observer finds the angular elevations of their tops to be complementary. Find the height of the shorter post.

6. The Bally bridge subtends an angle of  $45^\circ$  at a given point at the edge of the river; 800 yds. higher up, it subtends an angle of  $30^\circ$ . The course of the river here is straight and perpendicular to the bridge. Find the length of the bridge.

7. The height of a house subtends a right angle at an opposite window, the top being  $60^\circ$  above a horizontal straight line through the window; find the height of the house, taking the breadth of the street to be 30 feet.

8. From an aeroplane vertically over a straight road, the angles of depression of two consecutive milestones are observed to be  $45^\circ$  and  $60^\circ$ ; find the height of the aeroplane.

9. From a ship sailing due South-East at the rate of 5 miles an hour, a light-house is observed to be  $30^\circ$  North of East, and after 4 hours, it is seen due North; find the distance of the light-house from the final position of the ship.

10. The shadow of a tower standing on a level plane is found to be 40 feet longer when the sun's altitude is  $45^\circ$  than when it is  $60^\circ$ . Find the height of the tower.

11. From the lower window of a house the angular elevation of a church-steeple is found to be  $45^\circ$  and from a window 20 feet above, the elevation is  $30^\circ$ . How far is the church from the house ?

12. A light-house facing East sends out a fan-shaped beam of light extending from *S. E.* to *N. E.* An observer sailing due North, after meeting the light continues to see it for  $10\sqrt{2}$  minutes. When leaving the fan of light, the ship is 10 miles from the light-house. Find the speed of the ship.

13. A pole 100 ft. high stands vertically at the centre of a horizontal equilateral triangle, each side of which subtends an angle of  $60^\circ$  at the top of the pole. Find the side of the triangle.

14. Two chimneys are of equal height. A person standing between them in the line joining their bases observes the elevation of the nearer one to be  $60^\circ$ . After walking 80 feet in a direction at right angles to the line joining their bases, he observes the elevations of the two to be  $45^\circ$  and  $30^\circ$  respectively. Find the height and the distance between them.

15. At the foot of a mountain the elevation of its summit is  $45^\circ$ ; after ascending 1 mile towards the mountain up an incline of  $30^\circ$ , the elevation changes to  $60^\circ$ . Find the height of the mountain.

16. From a station, two light-houses *A* and *B* are seen in directions North and  $30^\circ$  East of North respectively; if *A* were one-third as far off as it really is, it would appear due West of *B*. If the distance of *B* from the station be 10 miles, find the distance of *B* from *A*.

17. A person walking along a straight road observes a tall tree standing in front of a tower, both being on the road before him. The elevation of the top of the tower is  $45^\circ$ , and of the top of the tree  $30^\circ$ ; on advancing 100 feet he finds the tower and the tree to have the same elevation  $60^\circ$ ; supposing the height of the eye of the man to be 5 feet, find the height of the tower and of the tree.

18. A man on the top of a rock rising on a seashore, observes a boat coming towards it at an angle of depression  $30^\circ$ ; 10 minutes later the angle of depression is  $60^\circ$ . The height of the rock being 4000 feet, find the speed of the boat in miles per hour.

19. A person walking along a straight level road observes the elevation of the top of a hill to be  $60^\circ$  when he is nearest the hill, and after walking 200 yards in a direction perpendicular to the direction of the hill from this point, observes the elevation to be  $30^\circ$ . Find the approximate height of the hill.

20. A square tower stands on a horizontal plane. From a point in this plane, only three of its upper corners are visible, and their angles of elevation are  $45^\circ$ ,  $60^\circ$ ,  $45^\circ$ . Find the ratio of the height of the tower to its breadth.

21. Two wheels, the sum of whose radii is 10 feet, are placed flatly on a table with their centres at a distance of 20 ft. An endless string, quite stretched, is partly wrapped round the wheels and crosses itself between them. Show that the length of the string is nearly 76.5 feet.

22. On a still day, from a station *A* an airship is observed due north at an elevation of  $60^\circ$ , while from a station *B* it is observed due east at an elevation of  $45^\circ$ . At this instant of observation, a parachute message is dropped from the airship, and the observer at *A* has to walk a mile to reach the message. Find the distance between the two stations.

23. From the foot of a column the angle of elevation of the top of a tower is  $45^\circ$  and from the top of the column the angle of depression of the bottom of the tower is  $30^\circ$ . A man walks 10 ft. from the bottom of the column towards the tower and notices the angle of elevation of its top to be  $60^\circ$ . Find the height of the column.

---


CHAPTER VI  
COMPOUND ANGLES

33. To prove that

$$\sin(A+B) = \sin A \cos B + \cos A \sin B$$

$$\cos(A+B) = \cos A \cos B - \sin A \sin B,$$

when  $A$  and  $B$  are positive and acute and  $(A+B) < 90^\circ$ .



Let a revolving line starting from the position  $OX$  trace out an angle  $XOY = A$  and then revolving further, trace out an angle  $YOZ = B$ ; then  $\angle XOZ = A + B$ .

In  $OZ$ , the bounding line of the compound angle  $A + B$ , take any point  $P$  and draw  $PQ$  and  $PR$  perpendicular to  $OX$  and  $OY$  respectively; also draw  $RS$  and  $RT$  perpendicular to  $OX$  and  $PQ$  respectively.

From the right-angled  $\triangle POQ$ ,

$$\begin{aligned} \sin(A+B) &= \frac{PQ}{OP} = \frac{QT + TP}{OP} = \frac{RS + PT}{OP} = \frac{RS}{OP} + \frac{PT}{OP} \\ &= \frac{RS}{OR} \cdot \frac{OR}{OP} + \frac{PT}{PR} \cdot \frac{PR}{OP} \\ &= \sin A \cos B + \cos A \sin B. \end{aligned}$$

Now,  $\angle TPR = 90^\circ - \angle TRP = \angle TRQ = \angle ROS = A$ .

$$\therefore \sin (A+B)=\sin A \cos B+\cos A \sin B.$$

Again,


$$\begin{aligned}\cos (A+B) &= \frac{OQ}{OP} = \frac{OS - QS}{OP} = \frac{OS - TR}{OP} = \frac{OS}{OP} - \frac{TR}{OP} \\ &= \frac{OS}{OR} \cdot \frac{OR}{OP} - \frac{TR}{PR} \cdot \frac{PR}{OP} \\ &= \cos A \cos B - \sin A \sin B \\ &= \cos A \cos B - \sin A \sin B.\end{aligned}$$

34. To prove that

$$\sin (A-B)=\sin A \cos B-\cos A \sin B$$

$$\cos (A-B)=\cos A \cos B+\sin A \sin B,$$

when  $A$  and  $B$  are positive and acute, and  $A > B$ .



Let a revolving line start from the position  $OX$  and trace out an angle  $XOY = A$  and then, revolving back trace out an angle  $YOZ = B$ ; then  $\angle XOZ = A - B$ .

In  $OZ$ , the bounding line of the compound angle  $A - B$ , take any point  $P$ , and draw  $PQ$  and  $PR$  perpendicular to  $OX$  and  $OY$  respectively; and draw  $RS$  and  $RT$  perpendicular to  $OX$  and  $QP$  produced respectively.

From the right-angled  $\triangle POQ$ ,

$$\sin (A-B) = \frac{PQ}{OP} = \frac{TQ - PT}{OP} = \frac{RS - PT}{OP} = \frac{RS}{OP} - \frac{PT}{OP}$$

$$\begin{aligned}
 &= \frac{RS}{OR} \cdot \frac{OR}{OP} - \frac{PT}{PR} \cdot \frac{PR}{OP} \\
 &= \sin A \cos B - \cos A \sin B.
 \end{aligned}$$

But  $\angle TPR = 90^\circ - \angle TRP = \angle YRT = \angle YOX = A$ .

$$\therefore \sin(A - B) = \sin A \cos B - \cos A \sin B.$$

Again,

$$\begin{aligned}
 \cos(A - B) &= \frac{OQ}{OP} = \frac{OS + SQ}{OP} = \frac{OS + RT}{OP} = \frac{OS}{OP} + \frac{RT}{OP} \\
 &= \frac{OS}{OR} \cdot \frac{OR}{OP} + \frac{RT}{RP} \cdot \frac{RP}{OP} \\
 &= \cos A \cos B + \sin A \sin B.
 \end{aligned}$$

**Obs.** In the above Geometrical proofs, it is assumed that the angles  $A, B, A+B$  are all less than a right angle and that  $A-B$  is positive. If the angles are not so restricted, the same method of proof (there being some modifications in the figures) will apply, due attention being paid to the signs of the quantities involved.\*

Thus the above four formulæ are perfectly general.

**Note 1.** The sum or difference of two or more angles is called a *Compound angle*; such as,  $A+B, A-B, A+B+C$  etc.

The expansions  $\sin(A \pm B)$  and  $\cos(A \pm B)$  are generally called the "*Addition Formulæ* or *Addition and Subtraction Theorems*".

**Note 2.** Assuming the truth of the above formulæ for acute angles, they can be shown to be true for angles of any magnitude, as follows:

Let us consider  $\sin(A+B)$ .

Let  $A$  and  $B$  be acute and  $A+B < 90^\circ$ .

Let  $A_1 = 90^\circ + A$ ;  $B_1 = B$ .

$$\begin{aligned}
 \text{Now, } \sin(A_1 + B_1) &= \sin\{(90^\circ + A) + B\} = \sin\{90^\circ + (A+B)\} \\
 &= \cos(A+B) = \cos A \cos B - \sin A \sin B, \quad [\text{by Art. 33.}] \\
 &= \sin(90^\circ + A) \cos B + \cos(90^\circ + A) \sin B \\
 &= \sin A_1 \cos B_1 + \cos A_1 \sin B_1.
 \end{aligned}$$

\*See Appendix, Art. 2-4. Also for alternative proofs of Arts. 33 and 34, see Appendix, Art. 10.

Again, let  $A_1 = -A$ ,  $B_1 = B$ .

$$\begin{aligned}\text{Then } \sin (A_1 + B_1) &= \sin (-A + B) = -\sin (A - B) \\ &= -\sin A \cos B + \cos A \sin B, \quad [\text{by Art. 34.}] \\ &= \sin (-A) \cos B + \cos (-A) \sin B \\ &= \sin A_1 \cos B_1 + \cos A_1 \sin B_1.\end{aligned}$$

Thus, the above formula remains true if any of the two angles is either increased by  $90^\circ$ , or has its sign changed.

In the same way it may be shown that the other three formulae for  $\cos (A + B)$ ,  $\sin (A - B)$  and  $\cos (A - B)$  will continue to hold good unchanged in form, if any of the two angles be either increased by  $90^\circ$  or has its sign changed.

Now starting from positive acute-angled values of  $A$  and  $B$ , combining the two processes of increasing one of the angles by  $90^\circ$ , and reversing the sign of any one, we can arrive at values of  $A$  and  $B$  of any magnitude, positive or negative, and the four formulae will still hold good.

Thus the formulae for  $\sin (A \pm B)$  and  $\cos (A \pm B)$  are perfectly general.

**35. Ex. 1.** Find the values of

$\sin 75^\circ$ ,  $\cos 75^\circ$ ,  $\sin 15^\circ$  and  $\cos 15^\circ$ .

$$\sin 75^\circ = \sin (45^\circ + 30^\circ) = \sin 45^\circ \cos 30^\circ + \cos 45^\circ \sin 30^\circ$$

$$= \frac{1}{\sqrt{2}} \cdot \frac{\sqrt{3}}{2} + \frac{1}{\sqrt{2}} \cdot \frac{1}{2} = \frac{\sqrt{3} + 1}{2\sqrt{2}}.$$

$$\cos 75^\circ = \cos (45^\circ + 30^\circ) = \cos 45^\circ \cos 30^\circ - \sin 45^\circ \sin 30^\circ$$

$$= \frac{1}{\sqrt{2}} \cdot \frac{\sqrt{3}}{2} - \frac{1}{\sqrt{2}} \cdot \frac{1}{2} = \frac{\sqrt{3} - 1}{2\sqrt{2}}.$$

$\sin 15^\circ = \sin (45^\circ - 30^\circ) = \sin 45^\circ \cos 30^\circ - \cos 45^\circ \sin 30^\circ$   
and  $\cos 15^\circ = \cos (45^\circ - 30^\circ) = \cos 45^\circ \cos 30^\circ + \sin 45^\circ \sin 30^\circ$ ;  
therefore substituting the values of  $\sin 45^\circ$ ,  $\cos 45^\circ$  etc.  
as before, we get



$$\sin 15^\circ = \frac{\sqrt{3}-1}{2\sqrt{2}} \quad \text{and} \quad \cos 15^\circ = \frac{\sqrt{3}+1}{2\sqrt{2}}.$$

**Note.** The values of  $\sin 15^\circ$  and  $\cos 15^\circ$  can also be deduced from the fact that

$$\sin 15^\circ = \sin (90^\circ - 75^\circ) = \cos 75^\circ$$

$$\text{and} \quad \cos 15^\circ = \cos (90^\circ - 75^\circ) = \sin 75^\circ.$$

**Ex. 2.** Show that

$$\begin{aligned} \text{(i)} \quad \sin (A+B) \sin (A-B) &= \sin^2 A - \sin^2 B \\ &= \cos^2 B - \cos^2 A. \end{aligned}$$

$$\begin{aligned} \text{(ii)} \quad \cos(A+B) \cos (A-B) &= \cos^2 A - \sin^2 B \\ &= \cos^2 B - \sin^2 A. \end{aligned}$$

(i) Left side

$$\begin{aligned} &= (\sin A \cos B + \cos A \sin B)(\sin A \cos B - \cos A \sin B) \\ &= \sin^2 A \cos^2 B - \cos^2 A \sin^2 B \\ &= \sin^2 A (1 - \sin^2 B) - (1 - \sin^2 A) \sin^2 B \\ &= \sin^2 A - \sin^2 B \\ &= (1 - \cos^2 A) - (1 - \cos^2 B) = \cos^2 B - \cos^2 A. \end{aligned}$$

(ii) Left side

$$\begin{aligned} &= (\cos A \cos B - \sin A \sin B)(\cos A \cos B + \sin A \sin B) \\ &= \cos^2 A \cos^2 B - \sin^2 A \sin^2 B \\ &= \cos^2 A (1 - \sin^2 B) - (1 - \cos^2 A) \sin^2 B \\ &= \cos^2 A - \sin^2 B \\ &= (1 - \sin^2 A) - (1 - \cos^2 B) = \cos^2 B - \sin^2 A. \end{aligned}$$

**Note.** The results of Ex. 1 and Ex. 2 are very useful and should be carefully remembered.

**36.** To prove that

$$\text{(i)} \quad \tan (A+B) = \frac{\tan A + \tan B}{1 - \tan A \tan B}$$

$$\text{(ii)} \quad \tan (A-B) = \frac{\tan A - \tan B}{1 + \tan A \tan B}$$

we have

$$\tan (A+B) = \frac{\sin (A+B)}{\cos (A+B)} = \frac{\sin A \cos B + \cos A \sin B}{\cos A \cos B - \sin A \sin B}.$$

Now dividing the numerator and denominator by  $\cos A \cos B$ , we have

$$\begin{aligned} \tan (A+B) &= \frac{\frac{\sin A \cos B}{\cos A \cos B} + \frac{\cos A \sin B}{\cos A \cos B}}{\frac{\cos A \cos B}{\cos A \cos B} - \frac{\sin A \sin B}{\cos A \cos B}} \\ &= \frac{\tan A + \tan B}{1 - \tan A \tan B}. \end{aligned}$$

Again,

$$\tan (A-B) = \frac{\sin (A-B)}{\cos (A-B)} = \frac{\sin A \cos B - \cos A \sin B}{\cos A \cos B + \sin A \sin B}.$$

Now, dividing the numerator and denominator by  $\cos A \cos B$ , we have, as before,

$$\tan (A-B) = \frac{\tan A - \tan B}{1 + \tan A \tan B}.$$

37. To prove that

$$(i) \cot (A+B) = \frac{\cot A \cot B - 1}{\cot B + \cot A}.$$

$$(ii) \cot (A-B) = \frac{\cot A \cot B + 1}{\cot B - \cot A}.$$

$$\cot (A+B) = \frac{\cos (A+B)}{\sin (A+B)} = \frac{\cos A \cos B - \sin A \sin B}{\sin A \cos B + \cos A \sin B}.$$

Now, dividing the numerator and denominator by  $\sin A \sin B$ , we have,

$$\begin{aligned} \cot (A+B) &= \frac{\frac{\cos A \cos B}{\sin A \sin B} - \frac{\sin A \sin B}{\sin A \sin B}}{\frac{\sin A \cos B}{\sin A \sin B} + \frac{\cos A \sin B}{\sin A \sin B}} \\ &= \frac{\cot A \cot B - 1}{\cot B + \cot A}. \end{aligned}$$

$$\cot(A-B) = \frac{\cos(A-B)}{\sin(A-B)} = \frac{\cos A \cos B + \sin A \sin B}{\sin A \cos B - \cos A \sin B}.$$

Now, dividing the numerator and denominator by  $\sin A \sin B$ , we have, as before,

$$\cot(A-B) = \frac{\cot A \cot B + 1}{\cot B - \cot A}.$$

38. Ex. 1. Find the values of  $\tan 75^\circ$  and  $\tan 15^\circ$ .

$$\tan 75^\circ = \tan(45^\circ + 30^\circ) = \frac{\tan 45^\circ + \tan 30^\circ}{1 - \tan 45^\circ \tan 30^\circ}$$

$$\begin{aligned} &= \frac{1 + \frac{1}{\sqrt{3}}}{1 - \frac{1}{\sqrt{3}}} = \frac{\sqrt{3} + 1}{\sqrt{3} - 1} = \frac{(\sqrt{3} + 1)(\sqrt{3} + 1)}{3 - 1} \\ &= \frac{4 + 2\sqrt{3}}{2} = 2 + \sqrt{3}. \end{aligned}$$

$$\tan 15^\circ = \tan(45^\circ - 30^\circ) = \frac{\tan 45^\circ - \tan 30^\circ}{1 + \tan 45^\circ \tan 30^\circ}$$

$$\begin{aligned} &= \frac{1 - \frac{1}{\sqrt{3}}}{1 + \frac{1}{\sqrt{3}}} = \frac{\sqrt{3} - 1}{\sqrt{3} + 1} = \frac{(\sqrt{3} - 1)(\sqrt{3} - 1)}{(\sqrt{3} + 1)(\sqrt{3} - 1)} \\ &= \frac{4 - 2\sqrt{3}}{2} = 2 - \sqrt{3}. \end{aligned}$$

Ex. 2. Show that

$$(i) \tan(45^\circ + A) = \frac{1 + \tan A}{1 - \tan A}.$$

$$(ii) \tan(45^\circ - A) = \frac{1 - \tan A}{1 + \tan A}.$$

$$(i) \text{ Left side} = \frac{\tan 45^\circ + \tan A}{1 - \tan 45^\circ \tan A} = \frac{1 + \tan A}{1 - \tan A}.$$

(ii) This result follows similarly.

Ex. (3.) Show that

$$\cot 2A + \tan A = \operatorname{cosec} 2A. \quad [C. U. 1947.]$$

$$\begin{aligned} \text{Left side} &= \frac{\cos 2A}{\sin 2A} + \frac{\sin A}{\cos A} = \frac{\cos 2A \cos A + \sin 2A \sin A}{\sin 2A \cos A} \\ &= \frac{\cos (2A - A)}{\sin 2A \cos A} = \frac{\cos A}{\sin 2A \cos A} = \frac{1}{\sin 2A} \\ &= \operatorname{cosec} 2A. \end{aligned}$$

39. To find the expansions of

(i)  $\sin (A+B+C)$

(ii)  $\cos (A+B+C)$

(iii)  $\tan (A+B+C)$

(i)  $\sin (A+B+C)$

$$\begin{aligned} &= \sin \{(A+B) + C\} \\ &= \sin (A+B) \cos C + \cos (A+B) \sin C \\ &= (\sin A \cos B + \cos A \sin B) \cos C \\ &\quad + (\cos A \cos B - \sin A \sin B) \sin C \\ &= \sin A \cos B \cos C + \sin B \cos C \cos A \\ &\quad + \sin C \cos A \cos B - \sin A \sin B \sin C. \end{aligned}$$

**Note 1.** The expansion of  $\sin (A+B+C)$  can be easily put in the form

$$\cos A \cos B \cos C (\tan A + \tan B + \tan C - \tan A \tan B \tan C).$$

(ii)  $\cos (A+B+C)$

$$\begin{aligned} &= \cos \{(A+B) + C\} \\ &= \cos (A+B) \cos C - \sin (A+B) \sin C \\ &= (\cos A \cos B - \sin A \sin B) \cos C \\ &\quad - (\sin A \cos B + \cos A \sin B) \sin C \\ &= \cos A \cos B \cos C - \cos A \sin B \sin C \\ &\quad - \cos B \sin C \sin A - \cos C \sin A \sin B. \end{aligned}$$

**Note 2.** The expansion of  $\cos (A+B+C)$  can be easily put in the form

$$\cos A \cos B \cos C (1 - \tan B \tan C - \tan C \tan A - \tan A \tan B).$$

(iii)  $\tan (A+B+C)$ 

$$\begin{aligned}
 &= \tan \{(A+B)+C\} \\
 &= \frac{\tan (A+B) + \tan C}{1 - \tan (A+B) \tan C} \\
 &= \frac{\frac{\tan A + \tan B}{1 - \tan A \tan B} + \tan C}{1 - \frac{\tan A + \tan B}{1 - \tan A \tan B} \tan C} \\
 &= \frac{\tan A + \tan B + \tan C - \tan A \tan B \tan C}{1 - \tan B \tan C - \tan C \tan A - \tan A \tan B}
 \end{aligned}$$

**Note 3.** The expansion of  $\tan (A+B+C)$  can also be obtained thus.

$$\tan (A+B+C) = \frac{\sin (A+B+C)}{\cos (A+B+C)}.$$

Now, write down the expansions of  $\sin (A+B+C)$  and  $\cos (A+B+C)$  and divide the numerator and denominator by  $\cos A \cos B \cos C$  or simply write down the expansions of  $\sin (A+B+C)$  and  $\cos (A+B+C)$  as given in Notes 1 and 2.

**Obs.** Formulae for the Trigonometrical functions of the sum of four, five or more angles can be similarly obtained.

### Examples VI

Show that (*Ex. 1 to 20*):—

1. (i)  $\sin (A-B) = \frac{3}{5}$  and  $\cos (A+B) = \frac{4}{5}$ ,

if  $A$  and  $B$  are acute and if  $\sin A = \frac{3}{5}$ ,  $\cos B = \frac{4}{5}$ .

(ii)  $\cos 68^\circ 20' \cos 8^\circ 20' + \cos 81^\circ 40' \cos 21^\circ 40' = \frac{1}{2}$ .

(iii)  $\sec (x-y) = \frac{5}{4}$ , if  $\sec x = \frac{5}{3}$ ,  $\operatorname{cosec} y = \frac{5}{4}$ .

2. (i)  $\sin A \sin (B-C) + \sin B \sin (C-A)$   
 $+ \sin C \sin (A-B) = 0$ .

(ii)  $\cos A \sin (B-C) + \cos B \sin (C-A)$   
 $+ \cos C \sin (A-B) = 0$ .

(iii)  $\sin (B+C) \sin (B-C) + \sin (C+A) \sin (C-A)$   
 $+ \sin (A+B) \sin (A-B) = 0$ .

$$\begin{aligned} & \cdot (\text{iv}) \sin(\alpha - \theta) \sin(\beta - \gamma) + \sin(\beta - \theta) \sin(\gamma - \alpha) \\ & \quad + \sin(\gamma - \theta) \sin(\alpha - \beta) = 0. \end{aligned}$$

$$\begin{aligned} 3. \quad & \cos(60^\circ - A) \cos(30^\circ - B) - \sin(60^\circ - A) \sin(30^\circ - B) \\ & = \sin(A + B). \end{aligned}$$

$$\begin{aligned} 4. \quad & (\text{i}) \sin(n+1)x \cos(n-1)x - \cos(n+1)x \sin(n-1)x \\ & = \sin 2x. \end{aligned}$$

$$\begin{aligned} & (\text{ii}) \sin 2\theta \cos \theta + \cos 2\theta \sin \theta \\ & = \sin 4\theta \cos \theta - \cos 4\theta \sin \theta. \end{aligned}$$

$$5. \quad \frac{\sin B}{\sin A} = \frac{\sin(2A+B)}{\sin A} - 2 \cos(A+B).$$

$$6. \quad \frac{\sin(B-C)}{\cos B \cos C} + \frac{\sin(C-A)}{\cos C \cos A} + \frac{\sin(A-B)}{\cos A \cos B} = 0.$$

$$7. \quad \frac{\sin(B-C)}{\sin B \sin C} + \frac{\sin(C-A)}{\sin C \sin A} + \frac{\sin(A-B)}{\sin A \sin B} = 0.$$

$$8. \quad \tan(A+B) \tan(A-B) = \frac{\sin^2 A - \sin^2 B}{\cos^2 A - \sin^2 B}.$$

$$9. \quad \tan^2 A - \tan^2 B = \frac{\sin(A+B) \sin(A-B)}{\cos^2 A \cos^2 B}$$

$$10. \quad (\text{i}) \frac{\tan(\alpha + \beta) - \tan \alpha}{1 + \tan(\alpha + \beta) \tan \alpha} = \tan \beta.$$

(ii) If  $A + B + C = \pi$  and  $\cos A = \cos B \cos C$ , show that  $\tan A = \tan B + \tan C$ . [C. U. 1942.]

$$11. \quad 1 + \tan 2\theta \tan \theta = \sec 2\theta.$$

$$12. \quad \cot \theta - \cot 2\theta = \operatorname{cosec} 2\theta.$$

$$13. \quad \tan 20^\circ + \tan 25^\circ + \tan 25^\circ \tan 20^\circ = 1.$$

$$14. \quad (\text{i}) \tan(45^\circ + A) = \frac{\cos A + \sin A}{\cos A - \sin A}.$$

$$(\text{ii}) \sqrt{2} \sin(45^\circ + A) = \sin A + \cos A.$$

$$15. \quad \frac{\cos 8^\circ + \sin 8^\circ}{\cos 8^\circ - \sin 8^\circ} = \tan 53^\circ = \tan(45^\circ + 8^\circ)$$

$$16. \tan(45^\circ + A) \tan(45^\circ - A) = 1.$$

$$17. \tan(A+B) + \tan(A-B) = \frac{\sin 2A}{\cos^2 A - \sin^2 B}.$$

$$18. \frac{\sin(x+y)}{\sin(x-y)} = \frac{\tan x + \tan y}{\tan x - \tan y}.$$

$$19. \cot(45^\circ + x) = \frac{\cot x - 1}{\cot x + 1} = \frac{\cos x - \sin x}{\cos x + \sin x}.$$

$$20. \sec(x+y) = \frac{\sec x \sec y}{1 - \tan x \tan y}.$$

21. Find the expansions of  
 $\sin(A-B+C)$  and  $\tan(A-B-C).$

22. Express  $\cot(A+B+C)$  in terms of  
 $\cot A, \cot B, \cot C.$

23. (i) If  $a \cos(x+a) = b \cos(x-a)$ , prove that  
 $(a+b) \tan x = (a-b) \cot a.$

(ii) If  $\sin a \sin \beta - \cos a \cos \beta + 1 = 0$ , show that  
 $1 + \cot a \tan \beta = 0. \quad [C. U. 1939.]$

(iii) If  $A+B+C=\pi$  and  $\cos A = \cos B \cos C$ ,  
 then  $\cot B \cot C = \frac{1}{2}.$

24. If  $\tan \theta = \frac{a \sin x + b \sin y}{a \cos x + b \cos y}$ ,  
 then  $a \sin(\theta-x) + b \sin(\theta-y) = 0.$

25. An angle  $\theta$  is divided into two parts  $\alpha, \beta$  such that  
 $\tan \alpha : \tan \beta = x : y$ ; prove that

$$\sin(\alpha-\beta) = \frac{x-y}{x+y} \sin \theta.$$

26. If  $\cos(\beta-\gamma) + \cos(\gamma-\alpha) + \cos(\alpha-\beta) = -\frac{3}{2}$ ,  
 show that  $\Sigma \cos \alpha = 0$  and  $\Sigma \sin \alpha = 0.$

## CHAPTER VII

### TRANSFORMATION OF PRODUCTS AND SUMS

#### 40. Transformation of products into sums or differences.

We have from Arts. 33 and 34,

$$\sin A \cos B + \cos A \sin B = \sin (A + B) \quad \dots \quad (1)$$

$$\sin A \cos B - \cos A \sin B = \sin (A - B). \quad \dots \quad (2)$$

Adding (1) and (2), we get

$$2 \sin A \cos B = \sin (A + B) + \sin (A - B). \quad \dots \quad (3)$$

Subtracting (2) from (1), we get

$$2 \cos A \sin B = \sin (A + B) - \sin (A - B). \quad \dots \quad (4)$$

Again, from Arts. 33 and 34, we have,

$$\cos A \cos B - \sin A \sin B = \cos (A + B) \quad \dots \quad (5)$$

$$\cos A \cos B + \sin A \sin B = \cos (A - B). \quad \dots \quad (6)$$

Adding (5) and (6), we get

$$2 \cos A \cos B = \cos (A + B) + \cos (A - B). \quad \dots \quad (7)$$

Subtracting (5) from (6), we get

$$2 \sin A \sin B = \cos (A - B) - \cos (A + B). \quad \dots \quad (8)$$

Thus, we have the following formulæ for transforming a *product* of two sines and cosines into the *sum* or the *difference* of two sines or two cosines.

$$2 \sin A \cos B = \sin (A + B) + \sin (A - B). \quad \dots \quad (I)$$

$$2 \cos A \sin B = \sin (A + B) - \sin (A - B). \quad \dots \quad (II)$$

$$2 \cos A \cos B = \cos (A + B) + \cos (A - B). \quad \dots \quad (III)$$

$$2 \sin A \sin B = \cos (A - B) - \cos (A + B). \quad \dots \quad (IV)$$



#### 41. Transformation of sums or differences into products.

Let  $A + B = C$ , and  $A - B = D$

$$\text{then, } A = \frac{C+D}{2} \text{ and } B = \frac{C-D}{2}.$$

Making these substitutions for  $A$  and  $B$  in the results (3), (4), (7), (8) of Art. 40 and noting that the relation (8) can be written as

$$\begin{aligned} \cos(A+B) - \cos(A-B) &= -2 \sin A \sin B \\ &= -2 \sin A \sin(-B), \end{aligned}$$

we have the following four formulæ for transforming the *sum* or the *difference* of **two sines** only or **two cosines** only into a product of sines and cosines.

$$\sin C + \sin D = 2 \sin \frac{C+D}{2} \cos \frac{C-D}{2}. \quad \dots \quad \text{(I)}$$

$$\sin C - \sin D = 2 \cos \frac{C+D}{2} \sin \frac{C-D}{2}. \quad \dots \quad \text{(II)}$$

$$\cos C + \cos D = 2 \cos \frac{C+D}{2} \cos \frac{C-D}{2}. \quad \dots \quad \text{(III)}$$

$$\cos C - \cos D = 2 \sin \frac{C+D}{2} \sin \frac{D-C}{2}. \quad \dots \quad \text{(IV)}$$

**Note.** The following concise verbal statement of the above four formulæ is sometimes very convenient.

- (i)  $\sin + \sin = 2 \sin (\frac{1}{2} \text{ sum}). \cos (\frac{1}{2} \text{ diff.}).$
- (ii)  $\sin - \sin = 2 \cos (\frac{1}{2} \text{ sum}). \sin (\frac{1}{2} \text{ diff.}).$
- (iii)  $\cos + \cos = 2 \cos (\frac{1}{2} \text{ sum}). \cos (\frac{1}{2} \text{ diff.}).$
- (iv)  $\cos - \cos = 2 \sin (\frac{1}{2} \text{ sum}). \sin (\frac{1}{2} \text{ diff. reversed}).$

#### 42. Ex. 1. Prove that

- (i)  $\cos 20^\circ \cos 40^\circ \cos 80^\circ = \frac{1}{8}.$
- (ii)  $\cos 80^\circ + \cos 40^\circ - \cos 20^\circ = 0.$

$$\begin{aligned}
 \text{(i) Left side} &= \frac{1}{2} \cos 20^\circ (2 \cos 40^\circ \cos 80^\circ) \\
 &= \frac{1}{2} \cos 20^\circ (\cos 120^\circ + \cos 40^\circ) \\
 &= \frac{1}{2} \cos 20^\circ \left(-\frac{1}{2} + \cos 40^\circ\right) \\
 &= -\frac{1}{4} \cos 20^\circ + \frac{1}{2} \cos 20^\circ \cos 40^\circ \\
 &= -\frac{1}{4} \cos 20^\circ + \frac{1}{4} (\cos 60^\circ + \cos 20^\circ) \\
 &= -\frac{1}{4} \cos 20^\circ + \frac{1}{4} \left(\frac{1}{2} + \cos 20^\circ\right) \\
 &= \frac{1}{8}.
 \end{aligned}$$

$$\begin{aligned}
 \text{(ii) Left side} &= (\cos 80^\circ + \cos 40^\circ) - \cos 20^\circ \\
 &= 2 \cos 60^\circ \cos 20^\circ - \cos 20^\circ \\
 &= 2 \cdot \frac{1}{2} \cos 20^\circ - \cos 20^\circ = 0.
 \end{aligned}$$

**Ex. 2.** Show that

$$\frac{\sin \theta + \sin 2\theta + \sin 4\theta + \sin 5\theta}{\cos \theta + \cos 2\theta + \cos 4\theta + \cos 5\theta} = \tan 3\theta.$$

$$\begin{aligned}
 \text{Numerator} &= (\sin 5\theta + \sin \theta) + (\sin 4\theta + \sin 2\theta) \\
 &= 2 \sin 3\theta \cos 2\theta + 2 \sin 3\theta \cos \theta \\
 &= 2 \sin 3\theta (\cos 2\theta + \cos \theta);
 \end{aligned}$$

$$\begin{aligned}
 \text{Denominator} &= (\cos 5\theta + \cos \theta) + (\cos 4\theta + \cos 2\theta) \\
 &= 2 \cos 3\theta \cos 2\theta + 2 \cos 3\theta \cos \theta \\
 &= 2 \cos 3\theta (\cos 2\theta + \cos \theta).
 \end{aligned}$$

$$\therefore \text{Left side} = \frac{2 \sin 3\theta (\cos 2\theta + \cos \theta)}{2 \cos 3\theta (\cos 2\theta + \cos \theta)} = \frac{\sin 3\theta}{\cos 3\theta} = \tan 3\theta.$$

**Ex. 3.** Express  $4 \cos A \cos B \cos C$  as the sum of four cosines.

$$\begin{aligned}
 4 \cos A \cos B \cos C &= 2 \cos A \cdot (2 \cos B \cos C) \\
 &= 2 \cos A \{\cos (B+C) + \cos (B-C)\} \\
 &= 2 \cos A \cos (B+C) + 2 \cos A \cos (B-C) \\
 &= \cos (A+B+C) + \cos (A-B-C) \\
 &\quad + \cos (A+B-C) + \cos (A-B+C).
 \end{aligned}$$

**Ex. 4.** Express as the product of three sines

$$\begin{aligned}
 &\sin (B+C-A) + \sin (C+A-B) + \sin (A+B-C) \\
 &\quad - \sin (A+B+C).
 \end{aligned}$$

Grouping together the first two terms and grouping together the last two terms, the given expression

$$\begin{aligned}
 &= 2 \sin C \cos (B - A) + 2 \cos (A + B) \sin (-C) \\
 &= 2 \sin C \{\cos (B - A) - \cos (A + B)\} \\
 &= 2 \sin C (2 \sin B \sin A) \\
 &= 4 \sin A \sin B \sin C.
 \end{aligned}$$

### Examples VII

Prove that (Ex. 1 to 17) :

1.  $\frac{\sin A + \sin B}{\sin A - \sin B} = \tan \frac{A+B}{2} \cot \frac{A-B}{2}.$
2.  $\frac{\cos A + \cos B}{\cos A - \cos B} = \cot \frac{A+B}{2} \cot \frac{A-B}{2}.$
3.  $\cos 20^\circ + \cos 100^\circ + \cos 140^\circ = 0.$
4.  $\sin \theta \sin (60^\circ - \theta) \sin (60^\circ + \theta) = \frac{1}{4} \sin 3\theta.$
5.  $\cos \theta \cos (60^\circ - \theta) \cos (60^\circ + \theta) = \frac{1}{4} \cos 3\theta.$
6.  $(\sin 3\alpha + \sin \alpha) \sin \alpha + (\cos 3\alpha - \cos \alpha) \cos \alpha = 0.$
7.  $\cos (A - D) \sin (B - C) + \cos (B - D) \sin (C - A) + \cos (C - D) \sin (A - B) = 0.$
8.  $\cos 20^\circ \cos 40^\circ \cos 60^\circ \cos 80^\circ = \frac{1}{16}.$
9.  $\sin 20^\circ \sin 40^\circ \sin 60^\circ \sin 80^\circ = \frac{1}{16}.$
10.  $\frac{\sin A + \sin B}{\cos A + \cos B} = \tan \frac{A+B}{2}.$
11.  $\frac{\sin A - \sin B}{\cos A - \cos B} = \cot \frac{A+B}{2}.$
12.  $\frac{\sin \theta + \sin 3\theta + \sin 5\theta + \sin 7\theta}{\cos \theta + \cos 3\theta + \cos 5\theta + \cos 7\theta} = \tan 4\theta.$
13.  $\frac{\sin 2A + \sin 5A - \sin A}{\cos 2A + \cos 5A + \cos A} = \tan 2A.$

$$\textcircled{4} \quad \frac{\sin(a+\beta) - 2 \sin a + \sin(a-\beta)}{\cos(a+\beta) - 2 \cos a + \cos(a-\beta)} = \tan a.$$

$$15. \quad \frac{\cos 7a + \cos 3a - \cos 5a - \cos a}{\sin 7a - \sin 3a - \sin 5a + \sin a} = \cot 2a.$$

$$16. \quad \sin 2A + \sin 2B + \sin 2C - \sin 2(A+B+C) \\ = 4 \sin(B+C) \sin(C+A) \sin(A+B).$$

$$17. \quad \cos A + \cos B + \cos C + \cos(A+B+C) \\ = 4 \cos \frac{B+C}{2} \cos \frac{C+A}{2} \cos \frac{A+B}{2}.$$

$$\textcircled{18} \quad \text{If } \sin x = k \sin y, \text{ prove that}$$

$$\tan \frac{1}{2}(x-y) = \frac{k-1}{k+1} \tan \frac{1}{2}(x+y).$$

$$19. \quad \text{If } \cos x + \cos y = \frac{1}{2} \text{ and } \sin x + \sin y = \frac{1}{2}, \text{ prove that} \\ \tan \frac{1}{2}(x+y) = \frac{3}{4}.$$

$$20. \quad \text{If } x \cos \alpha + y \sin \alpha = k = x \cos \beta + y \sin \beta, \text{ prove that}$$

$$\cos \frac{x}{\frac{1}{2}(\alpha+\beta)} = \sin \frac{y}{\frac{1}{2}(\alpha+\beta)} = \frac{k}{\cos \frac{1}{2}(\alpha-\beta)}.$$

$$21. \quad \text{If } \sin \theta + \sin \phi = a, \cos \theta + \cos \phi = b, \text{ prove that}$$

$$\tan \frac{\theta-\phi}{2} = \pm \sqrt{\frac{4-a^2-b^2}{a^2+b^2}}.$$

$$22. \quad \text{Prove that } \frac{\cos 10^\circ - \sin 10^\circ}{\cos 10^\circ + \sin 10^\circ} = \tan 35^\circ.$$

[ Note that  $\sin \theta = \cos(90^\circ - \theta)$  and  $\cos \theta = \sin(90^\circ \pm \theta)$ . ]

$$23. \quad \text{If } \operatorname{cosec} A + \sec A = \operatorname{cosec} B + \sec B, \text{ then}$$

$$\tan A \tan B = \cot \frac{1}{2}(A+B). \quad [P. U. 1936.]$$

$$24. \quad \text{Prove that}$$

$$\left( \frac{\cos A + \cos B}{\sin A - \sin B} \right)^n + \left( \frac{\sin A + \sin B}{\cos A - \cos B} \right)^n = 2 \cot^n \frac{A-B}{2},$$

or zero, according as  $n$  is even or odd. [ P. U. 1933. ]

**Obs.** By a method similar to that of the previous article the Trigonometrical ratios of any higher multiple of  $A$  can be expressed in terms of those of  $A$ .

**45. Ex. 1.** Express  $\sin 2A$  and  $\cos 2A$  in terms of  $\tan A$ . [C. U. 1931.]

$$\begin{aligned}\sin 2A &= 2 \sin A \cos A = 2 \frac{\sin A}{\cos A} \cdot \cos^2 A \\ &= 2 \tan A \cdot \frac{1}{\sec^2 A} = \frac{2 \tan A}{1 + \tan^2 A} \\ \cos 2A &= \cos^2 A - \sin^2 A = \cos^2 A - \cos^2 A \cdot \frac{\sin^2 A}{\cos^2 A} \\ &= \cos^2 A \left( 1 - \frac{\sin^2 A}{\cos^2 A} \right) = \frac{1}{\sec^2 A} (1 - \tan^2 A) \\ &= \frac{1 - \tan^2 A}{1 + \tan^2 A}\end{aligned}$$

**Ex. 2.** Express  $\cos 4A$  in terms of  $\cos A$ .

$$\begin{aligned}\text{Putting } \theta = 2A, \cos 4A &= \cos 2\theta = 2 \cos^2 \theta - 1 \\ &= 2(\cos 2A)^2 - 1 \\ &= 2(2 \cos^2 A - 1)^2 - 1 \\ &= 8 \cos^4 A - 8 \cos^2 A + 1.\end{aligned}$$

**Ex. 3.** Show that  $\frac{1 - \tan^2(45^\circ - A)}{1 + \tan^2(45^\circ - A)} = \sin 2A$ .

Let  $\theta = 45^\circ - A$ ; then

$$\begin{aligned}\text{Left side} &= \frac{1 - \tan^2 \theta}{1 + \tan^2 \theta} = \frac{1 - \frac{\sin^2 \theta}{\cos^2 \theta}}{1 + \frac{\sin^2 \theta}{\cos^2 \theta}} = \frac{\cos^2 \theta - \sin^2 \theta}{\cos^2 \theta + \sin^2 \theta} \\ &= \cos^2 \theta - \sin^2 \theta = \cos 2\theta \\ &= \cos (90^\circ - 2A) = \sin 2A.\end{aligned}$$

### Examples VIII

Prove the following identities (*Ex. 1 to 25*)

$$1. \quad \frac{\sin 2A}{1 + \cos 2A} = \tan A.$$

$$2. \quad \frac{\sin 2A}{1 - \cos 2A} = \cot A.$$

$$3. \quad \cot A - \tan A = 2 \cot 2A.$$

$$4. (i) (2 \cos \theta + 1)(2 \cos \theta - 1) = 2 \cos 2\theta + 1.$$

$$(ii) \tan \theta(1 + \sec 2\theta) = \tan 2\theta.$$

$$5. \quad \frac{\cot A - \tan A}{\cot A + \tan A} = \cos 2A.$$

$$6. \quad \tan A + \cot A = 2 \operatorname{cosec} 2A.$$

$$7. \quad \cos^4 \theta - \sin^4 \theta = \cos 2\theta.$$

$$8. \quad \cos^6 \theta - \sin^6 \theta = \cos 2\theta(1 - \frac{1}{4} \sin^2 2\theta).$$

$$9. \quad \cos^6 \theta + \sin^6 \theta = \frac{1}{4}(1 + 3 \cos^2 2\theta).$$

$$10. \quad \frac{\sin^2 \alpha - \sin^2 \beta}{\sin \alpha \cos \alpha - \sin \beta \cos \beta} = \tan(\alpha + \beta).$$

$$11. (i) \frac{1 - \cos 2\theta + \sin 2\theta}{1 + \cos 2\theta + \sin 2\theta} = \tan \theta. \quad [C. U. 1938.]$$

$$(ii) \frac{\sin a - \sqrt{1 + \sin 2a}}{\cos a - \sqrt{1 + \sin 2a}} = \cot a, \quad [a \text{ being positive and}$$

acute, and the square root being taken with positive sign.]

$$12. \quad \frac{\cos \theta + \sin \theta}{\cos \theta - \sin \theta} - \frac{\cos \theta - \sin \theta}{\cos \theta + \sin \theta} = 2 \tan 2\theta.$$

$$13. (i) \frac{\sin A + \sin 2A}{1 + \cos A + \cos 2A} = \tan A.$$

$$(ii) \frac{\sin 4\theta}{\cos 2\theta} \frac{1 - \cos 2\theta}{1 - \cos 4\theta} = \tan \theta.$$

$$(i) \frac{\cos A - \sin A}{\cos A + \sin A} = \sec 2A - \tan 2A.$$

$$(ii) \frac{\cos^3 \theta + \sin^3 \theta}{\cos \theta + \sin \theta} = 1 - \frac{1}{4} \sin 2\theta.$$

$$15. \cos^3 A \cos 3A + \sin^3 A \sin 3A = \cos^3 2A.$$

$$16. (i) 4(\cos^3 10^\circ + \sin^3 20^\circ) = 3(\cos 10^\circ + \sin 20^\circ).$$

$$(ii) \frac{1}{\sin 10^\circ} - \frac{\sqrt{3}}{\cos 10^\circ} = 4.$$

$$17. \tan 3\theta - \tan 2\theta - \tan \theta = \tan 3\theta \tan 2\theta \tan \theta.$$

$$18. \frac{\cot A}{\cot A - \cot 3A} - \frac{\tan A}{\tan 3A - \tan A} = 1.$$

$$19. \frac{1}{\tan 3\theta - \tan \theta} - \frac{1}{\cot 3\theta - \cot \theta} = \cot 2\theta.$$

$$20. \sin 8\theta = 8 \sin \theta \cos \theta \cos 2\theta \cos 4\theta.$$

$$21. (i) \cos 5\theta = 16 \cos^5 \theta - 20 \cos^3 \theta + 5 \cos \theta.$$

$$(ii) \sin 5\theta = 16 \sin^5 \theta - 20 \sin^3 \theta + 5 \sin \theta.$$

$$22. (i) \cot 3\theta = \frac{\cot^3 \theta - 3 \cot \theta}{3 \cot^2 \theta - 1}.$$

$$(ii) \tan 4\theta = \frac{4 \tan \theta - 4 \tan^3 \theta}{1 - 6 \tan^2 \theta + \tan^4 \theta}.$$

$$23. (i) \cos(120^\circ - A) + \cos A + \cos(120^\circ + A) = 0.$$

$$(ii) \cos^2(A - 120^\circ) + \cos^2 A + \cos^2(A + 120^\circ) = \frac{3}{2}.$$

$$24. \frac{2 \cos 2^n \theta + 1}{2 \cos \theta + 1} = (2 \cos \theta - 1)(2 \cos 2\theta - 1)(2 \cos 2^2 \theta - 1) \dots (2 \cos 2^{n-1} \theta - 1).$$

$$[ \text{Use } (2 \cos \theta + 1)(2 \cos \theta - 1) = 2 \cos 2\theta + 1. ]$$

$$25. \frac{\tan 2^n \theta}{\tan \theta} = (1 + \sec 2\theta)(1 + \sec 2^2 \theta)(1 + \sec 2^3 \theta) \dots (1 + \sec 2^n \theta).$$

$$[ \text{Use } \tan \theta(1 + \sec 2\theta) = \tan 2\theta. ]$$

$$26. \text{ If } \theta = \frac{\pi}{2^n + 1}, \text{ prove that}$$

$$2^n \cos \theta \cos 2\theta \cos 2^2 \theta \dots \cos 2^{n-1} \theta = 1.$$

27. (i) If  $\tan x = b/a$ , find the value of  $a \cos 2x + b \sin 2x$ .

(ii) If  $\tan^2 x + 2 \tan x \tan 2y = \tan^2 y + 2 \tan y \tan 2x$ , prove that each side = 1, or else,  $\tan x = \pm \tan y$ .

28. If  $\tan^2 \theta = 1 + 2 \tan^2 \phi$ , show that  $\cos 2\phi = 1 + 2 \cos 2\theta$ .

29. (i) If  $2 \tan \alpha = 3 \tan \beta$ , prove that

$$\tan(\alpha - \beta) = \frac{\sin 2\beta}{5 - \cos 2\beta}. \quad [C. U. 1946.]$$

(ii) If  $\frac{\tan(\alpha + \beta - \gamma)}{\tan(\alpha - \beta + \gamma)} = \tan \gamma$ , show that

either,  $\sin(\beta - \gamma) = 0$ , or,  $\sin 2\alpha + \sin 2\beta + \sin 2\gamma = 0$ .

30. If  $\alpha$  and  $\beta$  are acute angles and  $\cos 2\alpha = \frac{3 \cos 2\beta - 1}{3 - \cos 2\beta}$ , show that  $\tan \alpha = \sqrt{2} \tan \beta$ . [C. U. 1941.]

31. If  $\cos \theta = \frac{1}{2}(a + a^{-1})$ , show that

$$(i) \cos 2\theta = \frac{1}{2}(a^2 + a^{-2});$$

$$(ii) \cos 3\theta = \frac{1}{2}(a^3 + a^{-3}).$$

Show that (Ex. 32 to 36) :—

$$32. \sin^4 \theta = \frac{3}{8} - \frac{1}{2} \cos 2\theta + \frac{1}{8} \cos 4\theta.$$

$$33. \cos^4 \theta + \sin^4 \theta = 1 - \sin^2 2\theta + \frac{1}{8} \sin^4 2\theta.$$

$$34. \tan\left(\frac{\pi}{4} + A\right) + \tan\left(\frac{\pi}{4} - A\right) = 2 \sec 2A.$$

$$35. \cos^3 \theta \frac{\sin 3\theta}{3} + \sin^3 \theta \frac{\cos 3\theta}{3} = \frac{\sin 4\theta}{4}.$$

$$36. \cos 4x - \cos 4y \\ = 8(\cos x - \cos y)(\cos x + \cos y)(\cos x - \sin y) \\ \times (\cos x + \sin y).$$



# CHAPTER IX

## SUBMULTIPLE ANGLES

46. From the usual formulæ for multiple angles, namely

$$\sin 2A = 2 \sin A \cos A$$

$$\cos 2A = \cos^2 A - \sin^2 A = 2 \cos^2 A - 1 = 1 - 2 \sin^2 A$$

$$1 + \cos 2A = 2 \cos^2 A ; 1 - \cos 2A = 2 \sin^2 A$$

$$\tan 2A = \frac{2 \tan A}{1 - \tan^2 A}$$

$$\sin 3A = 3 \sin A - 4 \sin^3 A$$

$$\cos 3A = 4 \cos^3 A - 3 \cos A$$

$$\tan 3A = \frac{3 \tan A - \tan^3 A}{1 - 3 \tan^2 A}$$

putting  $A = \frac{1}{2}\theta$  and  $\frac{1}{3}\theta$  respectively, we derive the following formulæ for submultiple angles.

$$\sin \theta = 2 \sin \frac{1}{2}\theta \cos \frac{1}{2}\theta$$

$$\cos \theta = \cos^2 \frac{1}{2}\theta - \sin^2 \frac{1}{2}\theta = 2 \cos^2 \frac{1}{2}\theta - 1 = 1 - 2 \sin^2 \frac{1}{2}\theta$$

$$1 + \cos \theta = 2 \cos^2 \frac{1}{2}\theta ; 1 - \cos \theta = 2 \sin^2 \frac{1}{2}\theta$$

$$\tan \theta = \frac{2 \tan \frac{1}{2}\theta}{1 - \tan^2 \frac{1}{2}\theta}$$

$$\sin \theta = 3 \sin \frac{1}{3}\theta - 4 \sin^3 \frac{1}{3}\theta$$

$$\cos \theta = 4 \cos^3 \frac{1}{3}\theta - 3 \cos \frac{1}{3}\theta$$

$$\tan \theta = \frac{3 \tan \frac{1}{3}\theta - \tan^3 \frac{1}{3}\theta}{1 - 3 \tan^2 \frac{1}{3}\theta}$$

47. Values of  $\sin \frac{1}{2}\theta$  and  $\cos \frac{1}{2}\theta$  in terms of  $\cos \theta$ .

From  $\cos \theta = 2 \cos^2 \frac{1}{2}\theta - 1 = 1 - 2 \sin^2 \frac{1}{2}\theta$ , we at once deduce

$$\sin \frac{1}{2}\theta = \pm \sqrt{\frac{1}{2}(1 - \cos \theta)}$$

$$\cos \frac{1}{2}\theta = \pm \sqrt{\frac{1}{2}(1 + \cos \theta)}$$

48. *Ambiguity of signs explained.*

When  $\cos \theta$  is given and not  $\theta$ ,  $\theta$  and consequently  $\frac{1}{2}\theta$  has a series of values as will be explained in Chapter XI. Thus  $\frac{1}{2}\theta$  may lie in any quadrant and  $\sin \frac{1}{2}\theta$  and  $\cos \frac{1}{2}\theta$  will then have corresponding signs.

If the quadrant in which  $\frac{1}{2}\theta$  lies be known, for example, when  $\theta$  is given along with  $\cos \theta$ , there is no ambiguity in choosing the proper signs of  $\cos \frac{1}{2}\theta$  and  $\sin \frac{1}{2}\theta$ , as shown in the following example.

Ex. 1. Find  $\sin 22\frac{1}{2}^\circ$  and  $\cos 22\frac{1}{2}^\circ$ .

$$\sin 22\frac{1}{2}^\circ = + \sqrt{\frac{1}{2}(1 - \cos 45^\circ)} = \sqrt{\frac{1}{2}\left(1 - \frac{1}{\sqrt{2}}\right)} = \frac{1}{2}\sqrt{2} - \sqrt{2}$$

$$\cos 22\frac{1}{2}^\circ = + \sqrt{\frac{1}{2}(1 + \cos 45^\circ)} = \sqrt{\frac{1}{2}\left(1 + \frac{1}{\sqrt{2}}\right)} = \frac{1}{2}\sqrt{2} + \sqrt{2}.$$

49. Values of  $\sin \frac{1}{2}\theta$  and  $\cos \frac{1}{2}\theta$  in terms of  $\sin \theta$ .

We know that  $\sin \theta = 2 \sin \frac{1}{2}\theta \cos \frac{1}{2}\theta$

$$\text{and} \quad 1 = \cos^2 \frac{1}{2}\theta + \sin^2 \frac{1}{2}\theta.$$

Therefore,  $1 + \sin \theta = (\cos \frac{1}{2}\theta + \sin \frac{1}{2}\theta)^2$

$$\text{and} \quad 1 - \sin \theta = (\cos \frac{1}{2}\theta - \sin \frac{1}{2}\theta)^2.$$

Hence,  $\cos \frac{1}{2}\theta + \sin \frac{1}{2}\theta = \pm \sqrt{1 + \sin \theta}$

$$\cos \frac{1}{2}\theta - \sin \frac{1}{2}\theta = \pm \sqrt{1 - \sin \theta}.$$

Thus,  $\cos \frac{1}{2}\theta = \pm \frac{1}{2}\sqrt{1 + \sin \theta} \pm \frac{1}{2}\sqrt{1 - \sin \theta}$

$$\text{and} \quad \sin \frac{1}{2}\theta = \pm \frac{1}{2}\sqrt{1 + \sin \theta} \mp \frac{1}{2}\sqrt{1 - \sin \theta}$$

50. *Ambiguity of signs explained.*

As before, when  $\sin \theta$  is given, and not  $\theta$ ,  $\theta$  has a series of values for the given value of  $\sin \theta$  as will be explained in Chapter XI;  $\frac{1}{2}\theta$  may therefore lie in any one of two possible quadrants.

$$\cos \frac{1}{2}\theta + \sin \frac{1}{2}\theta = \sqrt{2} \sin \left(\frac{1}{4}\pi + \frac{1}{2}\theta\right)$$

$$\text{and } \cos \frac{1}{2}\theta - \sin \frac{1}{2}\theta = \sqrt{2} \sin \left(\frac{1}{4}\pi - \frac{1}{2}\theta\right)$$

will have their signs determined accordingly.

Thus,  $\sin \frac{1}{2}\theta$  and  $\cos \frac{1}{2}\theta$  will be definitely known.

**Ex. 1.** Find  $\sin 15^\circ$  and  $\cos 15^\circ$ .

We have,  $\cos 15^\circ + \sin 15^\circ = + \sqrt{1 + \sin 30^\circ} = \sqrt{1 + \frac{1}{2}}$

$$\cos 15^\circ - \sin 15^\circ = + \sqrt{1 - \sin 30^\circ} = \sqrt{1 - \frac{1}{2}}.$$

[  $\cos 15^\circ - \sin 15^\circ = \sqrt{2} \sin \left(\frac{1}{4}\pi - 15^\circ\right)$  and is clearly positive. ]

$$\text{Thus, } \cos 15^\circ = \frac{1}{2} \left( \sqrt{\frac{3}{2}} + \sqrt{\frac{1}{2}} \right) = \frac{\sqrt{3} + 1}{2\sqrt{2}}$$

$$\sin 15^\circ = \frac{1}{2} \left( \sqrt{\frac{3}{2}} - \sqrt{\frac{1}{2}} \right) = \frac{\sqrt{3} - 1}{2\sqrt{2}}.$$

**51.**  $\tan \frac{1}{2}\theta$  in terms of  $\tan \theta$ .

$$\text{From the formula, } \tan \theta = \frac{2 \tan \frac{1}{2}\theta}{1 - \tan^2 \frac{1}{2}\theta},$$

$$\text{i.e. } \tan \theta \tan^2 \frac{1}{2}\theta + 2 \tan \frac{1}{2}\theta - \tan \theta = 0,$$

we easily deduce

$$\tan \frac{1}{2}\theta = \frac{-1 + \sqrt{1 + \tan^2 \theta}}{\tan \theta}.$$

The reason of the ambiguity is similar to those of the previous cases.

**52.** Ratios of  $\frac{1}{3}\theta$  from those of  $\theta$ .

By resolving the cubic equation

$$\sin \theta = 3 \sin \frac{1}{3}\theta - 4 \sin^3 \frac{1}{3}\theta \quad \dots (1)$$

we get  $\sin \frac{1}{3}\theta$ , if  $\sin \theta$  be known.

Similarly, by solving the cubic equations

$$\cos \theta = 4 \cos^3 \frac{1}{3}\theta - 3 \cos \frac{1}{3}\theta \quad \dots (2)$$

$$\text{and } \tan \theta = \frac{3 \tan \frac{1}{3}\theta - \tan^3 \frac{1}{3}\theta}{1 - 3 \tan^2 \frac{1}{3}\theta} \quad \dots (3)$$

we derive values of  $\cos \frac{1}{3}\theta$  from those of  $\cos \theta$ , and of  $\tan \frac{1}{3}\theta$  from those of  $\tan \theta$  respectively.

### 53. Ratios of $18^\circ$ and $36^\circ$ .

Let  $\theta = 18^\circ$ ; then  $5\theta = 90^\circ$ ;  $\therefore 2\theta = 90^\circ - 3\theta$ .

$$\therefore \sin 2\theta = \cos 3\theta \text{ or } 2 \sin \theta \cos \theta = \cos \theta (4 \cos^2 \theta - 3).$$

As  $\cos \theta$  (*i.e.*  $\cos 18^\circ$ ) is not zero, we have

$$2 \sin \theta = 4 \cos^2 \theta - 3 = 1 - 4 \sin^2 \theta,$$

$$\text{or, } 4 \sin^2 \theta + 2 \sin \theta - 1 = 0.$$

$$\therefore \sin \theta = \frac{-2 \pm \sqrt{4+16}}{8} = \frac{(+\sqrt{5}-1)}{4}.$$

Now, as  $\theta$  here is a positive acute angle, therefore, rejecting the negative value, we get

$$\sin 18^\circ = \frac{1}{4}(\sqrt{5}-1).$$

$$\cos 18^\circ = + \sqrt{1 - \sin^2 18^\circ} = \frac{1}{4}(\sqrt{10+2\sqrt{5}}).$$

$$\cos 36^\circ = 1 - 2 \sin^2 18^\circ = \frac{1}{2}(\sqrt{5}+1).$$

$$\sin 36^\circ = \sqrt{1 - \cos^2 36^\circ} = \frac{1}{4}(\sqrt{10-2\sqrt{5}}).$$

**Note.** Since  $54^\circ$  and  $36^\circ$  are complementary and  $72^\circ$  and  $18^\circ$  are complementary, from the above values we easily get the trigonometrical ratios of  $54^\circ$  and  $72^\circ$ .

### 54. Ratios of $3^\circ$ and multiples of $3^\circ$ .

$$\begin{aligned} \sin 3^\circ &= \sin (18^\circ - 15^\circ) = \sin 18^\circ \cos 15^\circ - \cos 18^\circ \sin 15^\circ \\ &= \frac{1}{4}(\sqrt{5}-1)(\sqrt{6}+\sqrt{2}) - \frac{1}{8}(\sqrt{3}-1)(\sqrt{5}+\sqrt{5}), \end{aligned}$$

on substituting the values of  $\sin 18^\circ$ ,  $\cos 15^\circ$  etc.

Similarly,

$$\cos 3^\circ = \frac{1}{8}(\sqrt{3}+1)(\sqrt{5}+\sqrt{5}) + \frac{1}{8}(\sqrt{6}-\sqrt{2})(\sqrt{5}-1).$$

From a knowledge of the ratios of  $3^\circ$ ,  $15^\circ$ ,  $18^\circ$ ,  $30^\circ$ ,  $36^\circ$  and  $45^\circ$ , we can deduce the ratios for all angles which

are multiples of  $3^\circ$ , (for,  $6^\circ = 36^\circ - 30^\circ$ ;  $9^\circ = 45^\circ - 36^\circ$ ;  $12^\circ = 30^\circ - 18^\circ$ ;  $21^\circ = 36^\circ - 15^\circ$ ; etc.). For angles greater than  $45^\circ$ , the ratios may be deduced from those of their complements which are less than  $45^\circ$ .

**Ex.** Show that

$$\sin x = 2^n \cos \frac{x}{2} \cos \frac{x}{2^2} \cos \frac{x}{2^3} \cdots \cos \frac{x}{2^n} \sin \frac{x}{2^n}.$$

We have,  $\sin x = 2 \sin \frac{x}{2} \cos \frac{x}{2}$

$$\sin \frac{x}{2} = 2 \sin \frac{x}{2^2} \cos \frac{x}{2^2}$$

$$\sin \frac{x}{2^2} = 2 \sin \frac{x}{2^3} \cos \frac{x}{2^3}.$$

Similarly,  $\sin \frac{x}{2^{n-1}} = 2 \sin \frac{x}{2^n} \cos \frac{x}{2^n}.$

Hence,  $\sin x = 2^n \cos \frac{x}{2} \cos \frac{x}{2^2} \cos \frac{x}{2^3} \cdots \cos \frac{x}{2^n} \sin \frac{x}{2^n}.$

### Examples IX

Prove that (*Ex. 1 to 14*):—

1.  $\frac{1 - \cos A}{\sin A} = \tan \frac{A}{2}.$       2.  $\frac{1 + \cos A}{\sin A} = \cot \frac{A}{2}.$

3.  $\left( \sin \frac{A}{2} \pm \cos \frac{A}{2} \right)^2 = 1 \pm \sin A.$

4.  $\sec \theta + \tan \theta = \tan \left( \frac{1}{2}\pi + \frac{1}{2}\theta \right).$  [ *C. U. 1939.* ]

5. (i)  $\frac{1 + \sin \theta - \cos \theta}{1 + \sin \theta + \cos \theta} = \tan \frac{\theta}{2}.$

(ii)  $\frac{\sin \frac{1}{2}a - \sqrt{1 + \sin a}}{\cos \frac{1}{2}a - \sqrt{1 + \sin a}} = \cot \frac{a}{2},$  where  $0 < a < \pi$ , and the square root is taken with positive sign.

6. (i)  $\frac{1 + \sin x}{1 - \sin x} = \tan^2 \left( \frac{\pi}{4} + \frac{x}{2} \right).$

(ii)  $\frac{2 \sin \theta - \sin 2\theta}{2 \sin \theta + \sin 2\theta} = \tan^2 \frac{1}{2}\theta.$

$$7. \textcircled{i} \frac{1 + \tan \frac{1}{2}A}{1 - \tan \frac{1}{2}A} = \frac{1 + \sin A}{\cos A}.$$

$$(ii) \cot \beta = \frac{1}{2}(\cot \frac{1}{2}\beta - \tan \frac{1}{2}\beta).$$

$$8. (i) \frac{\sin 2\theta}{1 + \cos 2\theta} \cdot \frac{\cos \theta}{1 + \cos \theta} = \tan \frac{\theta}{2}.$$

$$(ii) 8 \sin^4 \frac{1}{2}\theta - 8 \sin^2 \frac{1}{2}\theta + 1 = \cos 2\theta.$$

$$9. \sin \theta = \frac{2 \tan \frac{1}{2}\theta}{1 + \tan^2 \frac{1}{2}\theta}.$$

$$10. \cos \theta = \frac{1 - \tan^2 \frac{1}{2}\theta}{1 + \tan^2 \frac{1}{2}\theta}.$$

$$11. (\cos x + \cos y)^2 + (\sin x + \sin y)^2 = 4 \cos^2 \frac{1}{2}(x + y).$$

$$12. \tan 6^\circ \tan 42^\circ \tan 66^\circ \tan 78^\circ = 1.$$

$$\textcircled{13} \tan 7\frac{1}{2}^\circ = \sqrt{6} - \sqrt{3} + \sqrt{2} - 2.$$

$$14. 2 \cos \frac{1}{8}\pi = \sqrt{2} + \sqrt{2} + \sqrt{2}.$$

$$15. (i) \text{ If } \tan \frac{\theta}{2} = \sqrt{\frac{1-e}{1+e}} \cdot \tan \frac{\phi}{2}, \text{ show that}$$

$$\cos \phi = \frac{\cos \theta - e}{1 - e \cos \theta}.$$

$$(ii) \text{ If } \tan \theta = \frac{\sin a \sin \beta}{\cos a + \cos \beta}, \text{ show that one of the}$$

values of  $\tan \frac{1}{2}\theta$  is  $\tan \frac{1}{2}a \tan \frac{1}{2}\beta$ .

16. If  $\sin a + \sin \beta = a$  and  $\cos a + \cos \beta = b$ , find the value of  $\cos(a + \beta)$ .

17. (i) Prove that  $2 \sin \frac{1}{2}A = \pm \sqrt{1 + \sin A} \pm \sqrt{1 - \sin A}$ , and determine which are the correct signs when  $270^\circ > A > 180^\circ$ . [B. H. U. I. 1931.]

(ii) If  $\theta = 240^\circ$ , is the following statement correct?

$$2 \sin \frac{1}{2}\theta = \sqrt{1 + \sin \theta} - \sqrt{1 - \sin \theta}.$$

If not, how must it be modified?

18. If  $A = 320^\circ$ , prove that

$$\tan \frac{A}{2} = \frac{-1 + \sqrt{1 + \tan^2 A}}{\tan A}.$$

## CHAPTER X

### TRIGONOMETRICAL IDENTITIES

55. Many interesting identities involving functions of three or more angles can be established when there exists a relation among the angles. The most important of these identities are those in which the three angles are connected by the relation that their sum is equal to two right angles. In establishing this latter kind of identities, it will be necessary to make frequent use of the properties of supplementary and complementary angles.

Thus, since,  $A + B + C = \pi$ ,

$$\therefore B + C = \pi - A.$$

$$\therefore \sin(B + C) = \sin(\pi - A) = \sin A.$$

Similarly,  $\sin(C + A) = \sin B$  ;  $\sin(A + B) = \sin C$ .

Again,  $\cos(B + C) = \cos(\pi - A) = -\cos A$ .

Similarly,  $\cos(C + A) = -\cos B$  ;  $\cos(A + B) = -\cos C$ .

$$\tan(B + C) = \tan(\pi - A) = -\tan A.$$

Similarly,  $\tan(C + A) = -\tan B$  ;  $\tan(A + B) = -\tan C$ .

Again, since,  $\frac{A}{2} + \frac{B}{2} + \frac{C}{2} = \frac{\pi}{2}$ ,

$$\therefore \sin\left(\frac{B}{2} + \frac{C}{2}\right) = \sin\left(\frac{\pi}{2} - \frac{A}{2}\right) = \cos \frac{A}{2}$$

Similarly,  $\sin\left(\frac{C}{2} + \frac{A}{2}\right) = \cos \frac{B}{2}$  ;

$$\sin\left(\frac{A}{2} + \frac{B}{2}\right) = \cos \frac{C}{2}.$$

Again,  $\cos\left(\frac{B}{2} + \frac{C}{2}\right) = \cos\left(\frac{\pi}{2} - \frac{A}{2}\right) = \sin \frac{A}{2}$

$$\text{Similarly, } \cos \left( \frac{C}{2} + \frac{A}{2} \right) = \sin \frac{B}{2};$$

$$\cos \left( \frac{A}{2} + \frac{B}{2} \right) = \sin \frac{C}{2}$$

$$\tan \left( \frac{B}{2} + \frac{C}{2} \right) = \tan \left( \frac{\pi}{2} - \frac{A}{2} \right) = \cot \frac{A}{2}.$$

$$\text{Similarly, } \tan \left( \frac{C}{2} + \frac{A}{2} \right) = \cot \frac{B}{2};$$

$$\tan \left( \frac{A}{2} + \frac{B}{2} \right) = \cot \frac{C}{2}.$$

56. **Ex. 1.** *If  $A+B+C=\pi$ , prove that*

$$\sin 2A + \sin 2B + \sin 2C = 4 \sin A \sin B \sin C.$$

[ C. U. 1930, '33, '35. ]

$$\text{Left side} = (\sin 2A + \sin 2B) + \sin 2C$$

$$= 2 \sin (A+B) \cos (A-B) + 2 \sin C \cos C$$

$$= 2 \sin C \cos (A-B) + 2 \sin C \cos C$$

[  $\because A+B+C=\pi$ . ]

$$= 2 \sin C [\cos (A-B) + \cos C]$$

$$= 2 \sin C [\cos (A-B) - \cos (A+B)]$$

[  $\because A+B+C=\pi$ . ]

$$= 2 \sin C \cdot 2 \sin A \sin B$$

$$= 4 \sin A \sin B \sin C.$$

**Ex. 2.** *If  $A+B+C=\pi$ , prove that*

$$\cos 2A + \cos 2B + \cos 2C = -4 \cos A \cos B \cos C - 1.$$

$$\text{Left side} = (\cos 2A + \cos 2B) + \cos 2C$$

$$= 2 \cos (A+B) \cos (A-B) + 2 \cos^2 C - 1$$

$$= -2 \cos C \cos (A-B) + 2 \cos^2 C - 1$$

[  $\because A+B+C=\pi$ . ]

$$= -2 \cos C [\cos (A-B) - \cos C] - 1$$

$$= -2 \cos C [\cos (A-B) + \cos (A+B)] - 1$$

[  $\because A+B+C=\pi$ . ]

$$= -2 \cos C \cdot 2 \cos A \cos B - 1$$

$$= -4 \cos A \cos B \cos C - 1.$$



**Ex. 3.** If  $A+B+C=\pi$ , prove that

$$\sin A + \sin B + \sin C = 4 \cos \frac{A}{2} \cos \frac{B}{2} \cos \frac{C}{2}.$$

[C. U. 1910, '29.]

$$\text{Left side} = (\sin A + \sin B) + \sin C$$

$$= 2 \sin \frac{A+B}{2} \cos \frac{A-B}{2} + 2 \sin \frac{C}{2} \cos \frac{C}{2}$$

$$= 2 \cos \frac{C}{2} \cos \frac{A-B}{2} + 2 \sin \frac{C}{2} \cos \frac{C}{2}$$

$$\left[ \because \frac{A}{2} + \frac{B}{2} + \frac{C}{2} = \frac{\pi}{2} \right]$$

$$= 2 \cos \frac{C}{2} \left\{ \cos \frac{A-B}{2} + \sin \frac{C}{2} \right\}$$

$$= 2 \cos \frac{C}{2} \left[ \cos \frac{A-B}{2} + \cos \frac{A+B}{2} \right]$$

$$\left[ \because \frac{A}{2} + \frac{B}{2} + \frac{C}{2} = \frac{\pi}{2} \right]$$

$$= 2 \cos \frac{C}{2} \cdot 2 \cos \frac{A}{2} \cos \frac{B}{2}$$

$$= 4 \cos \frac{A}{2} \cos \frac{B}{2} \cos \frac{C}{2}$$

**Ex. 4.** If  $A+B+C=\pi$ , prove that

$$\cos A + \cos B + \cos C = 1 + 4 \sin \frac{A}{2} \sin \frac{B}{2} \sin \frac{C}{2}.$$

$$\text{Left side} = (\cos A + \cos B) + \cos C$$

$$= 2 \cos \frac{A+B}{2} \cos \frac{A-B}{2} + 1 - 2 \sin^2 \frac{C}{2}$$

$$= 2 \sin \frac{C}{2} \cos \frac{A-B}{2} - 2 \sin^2 \frac{C}{2} + 1$$

$$\left[ \because \frac{A}{2} + \frac{B}{2} + \frac{C}{2} = \frac{\pi}{2} \right]$$

$$\begin{aligned}
&= 2 \sin \frac{C}{2} \left[ \cos \frac{A-B}{2} - \sin \frac{C}{2} \right] + 1 \\
&= 2 \sin \frac{C}{2} \left[ \cos \frac{A-B}{2} - \cos \frac{A+B}{2} \right] + 1 \\
&\quad \left[ \because \frac{A}{2} + \frac{B}{2} + \frac{C}{2} = \frac{\pi}{2} \right] \\
&= 2 \sin \frac{C}{2} \cdot 2 \sin \frac{A}{2} \sin \frac{B}{2} + 1 \\
&= 1 + 4 \sin \frac{A}{2} \sin \frac{B}{2} \sin \frac{C}{2}
\end{aligned}$$

**Ex. 5.** *If  $A+B+C=\pi$ , prove that*

$$\tan A + \tan B + \tan C = \tan A \tan B \tan C.$$

Since,

$$B + C = \pi - A,$$

$$\therefore \tan(B + C) = \tan(\pi - A).$$

$$\therefore \frac{\tan B + \tan C}{1 - \tan B \tan C} = -\tan A,$$

$$\begin{aligned}
\text{i.e. } \tan B + \tan C &= -\tan A(1 - \tan B \tan C) \\
&= -\tan A + \tan A \tan B \tan C.
\end{aligned}$$

$$\therefore \tan A + \tan B + \tan C = \tan A \tan B \tan C.$$

*Otherwise :*

$$\tan(A + B + C) = \tan \pi = 0.$$

$$\therefore \frac{\tan A + \tan B + \tan C - \tan A \tan B \tan C}{1 - \tan B \tan C - \tan C \tan A - \tan A \tan B} = 0.$$

Since the fraction is zero, numerator must be zero.

$$\therefore \tan A + \tan B + \tan C - \tan A \tan B \tan C = 0,$$

$$\text{i.e. } \tan A + \tan B + \tan C = \tan A \tan B \tan C.$$

**Ex. 6.** *If  $A+B+C=\pi$ , prove that*

$$\tan \frac{B}{2} \tan \frac{C}{2} + \tan \frac{C}{2} \tan \frac{A}{2} + \tan \frac{A}{2} \tan \frac{B}{2} = 1.$$

[ C. U. 1936, '39. ]

Since,  $A + B + C = \pi$ ,  $\therefore \frac{A}{2} + \frac{B}{2} + \frac{C}{2} = \frac{\pi}{2}$ .

$$\therefore \tan \left( \frac{B}{2} + \frac{C}{2} \right) = \tan \left( \frac{\pi}{2} - \frac{A}{2} \right),$$

$$\therefore \frac{\tan \frac{B}{2} + \tan \frac{C}{2}}{1 - \tan \frac{B}{2} \tan \frac{C}{2}} = \cot \frac{A}{2} = \frac{1}{\tan \frac{A}{2}}$$

$$\text{or, } \tan \frac{A}{2} \left( \tan \frac{B}{2} + \tan \frac{C}{2} \right) = 1 - \tan \frac{B}{2} \tan \frac{C}{2}.$$

On simplification, the required result follows.

*Otherwise :*

$$\begin{aligned} \tan \left( \frac{A}{2} + \frac{B}{2} + \frac{C}{2} \right) &= \cot \left( \frac{A}{2} + \frac{B}{2} + \frac{C}{2} \right) = \cot \frac{\pi}{2} \\ \therefore \frac{1 - \tan \frac{B}{2} \tan \frac{C}{2} - \tan \frac{C}{2} \tan \frac{A}{2} - \tan \frac{A}{2} \tan \frac{B}{2}}{\tan \frac{A}{2} + \tan \frac{B}{2} + \tan \frac{C}{2} - \tan \frac{A}{2} \tan \frac{B}{2} \tan \frac{C}{2}} &= 0. \end{aligned}$$

Now the value of the fraction being zero, its numerator must be zero.

$$\therefore 1 - \tan \frac{B}{2} \tan \frac{C}{2} - \tan \frac{C}{2} \tan \frac{A}{2} - \tan \frac{A}{2} \tan \frac{B}{2} = 0,$$

whence the required result follows.

**Ex. 7.** If  $A+B+C=\pi$ , prove that

$$\cos \frac{A}{2} + \cos \frac{B}{2} + \cos \frac{C}{2} = 4 \cos \frac{\pi-A}{4} \cos \frac{\pi-B}{4} \cos \frac{\pi-C}{4}.$$

$$\begin{aligned} \text{Right side} &= 2 \cos \frac{\pi-A}{4} \left[ 2 \cos \frac{\pi-B}{4} \cos \frac{\pi-C}{4} \right] \\ &= 2 \cos \frac{\pi-A}{4} \left[ \cos \frac{2\pi-(B+C)}{4} + \cos \frac{B-C}{4} \right] \\ &= 2 \cos \frac{\pi-A}{4} \left[ \cos \frac{\pi+A}{4} + \cos \frac{B-C}{4} \right] \end{aligned}$$

$$[\because 2\pi-(B+C)=\pi+\pi-(B+C)=\pi+A, \text{ since, } A+B+C=\pi.]$$

$$\begin{aligned}
&= 2 \cos \frac{\pi-A}{4} \cos \frac{\pi+A}{4} + 2 \cos \frac{\pi-A}{4} \cos \frac{B-C}{4} \\
&= \left( \cos \frac{\pi}{2} + \cos \frac{A}{2} \right) + 2 \cos \frac{B+C}{4} \cos \frac{B-C}{4} \\
&\qquad\qquad\qquad A+B+C=\pi. ] \\
&= \cos \frac{A}{2} + \cos \frac{B}{2} + \cos \frac{C}{2}.
\end{aligned}$$

**Note.** Since  $\cos \frac{1}{2}(\pi-A) = \sin \{ \frac{1}{2}\pi - \frac{1}{2}(\pi-A) \} = \sin \frac{1}{2}(\pi+A)$   
and  $\cos \frac{1}{2}(\pi-A) = \cos \frac{1}{2}(A+B+C-A) = \cos \frac{1}{2}(B+C)$ ,

$$\begin{aligned}
\therefore \text{ we have also, } &\cos \frac{1}{2}A + \cos \frac{1}{2}B + \cos \frac{1}{2}C \\
&= 4 \sin \frac{1}{4}(\pi+A) \sin \frac{1}{4}(\pi+B) \sin \frac{1}{4}(\pi+C) \\
&= 4 \cos \frac{1}{4}(B+C) \cos \frac{1}{4}(C+A) \cos \frac{1}{4}(A+B).
\end{aligned}$$

**Ex. 8.** If  $A+B+C=\pi$ , prove that

$$\cos^2 A + \cos^2 B + \cos^2 C + 2 \cos A \cos B \cos C = 1.$$

[ C. U. 1932, '37, '47. ]

$$\begin{aligned}
&\cos^2 A + \cos^2 B + \cos^2 C \\
&= \frac{1}{2}(2 \cos^2 A + 2 \cos^2 B) + \cos^2 C \\
&= \frac{1}{2}(1 + \cos 2A + 1 + \cos 2B) + \cos^2 C \\
&= 1 + \frac{1}{2}(\cos 2A + \cos 2B) + \cos^2 C \\
&= 1 + \cos(A+B) \cos(A-B) + \cos C \cos C \\
&= 1 - \cos C \cos(A-B) - \cos C \cos(A+B) \\
&\qquad\qquad\qquad [ \because A+B=\pi-C. ] \\
&= 1 - \cos C [\cos(A-B) + \cos(A+B)] \\
&= 1 - \cos C [2 \cos A \cos B] \\
&= 1 - 2 \cos A \cos B \cos C
\end{aligned}$$

whence the required result follows.

✓ **Ex. 9.** Show that

$$\begin{aligned}
&\tan(\beta-\gamma) + \tan(\gamma-\alpha) + \tan(\alpha-\beta) \\
&\qquad\qquad\qquad = \tan(\beta-\gamma) \tan(\gamma-\alpha) \tan(\alpha-\beta).
\end{aligned}$$

Let  $A = \beta - \gamma$ ,  $B = \gamma - \alpha$ ,  $C = \alpha - \beta$  ;  
 then  $A + B + C = \beta - \gamma + \gamma - \alpha + \alpha - \beta = 0$   
 $\therefore \tan (A + B + C) = \tan 0 = 0$ .  
 $\therefore \tan A + \tan B + \tan C = \tan A \tan B \tan C$ .

Now, substituting the values for  $A, B, C$ , the required result follows.

**Ex. 10.** If  $x + y + z = xyz$ , prove that

$$x(1 - y^2)(1 - z^2) + y(1 - z^2)(1 - x^2) + z(1 - x^2)(1 - y^2) = 4xyz.$$

Putting  $x = \tan \alpha$ ,  $y = \tan \beta$ ,  $z = \tan \gamma$ , in the given relation, we have

$$\tan \alpha + \tan \beta + \tan \gamma = \tan \alpha \tan \beta \tan \gamma.$$

$\therefore$  by transposition,

$$\tan \alpha(1 - \tan \beta \tan \gamma) = -(\tan \beta + \tan \gamma),$$

$$\text{i.e. } \tan \alpha = -\frac{\tan \beta + \tan \gamma}{1 - \tan \beta \tan \gamma} = -\tan(\beta + \gamma).$$

$$\therefore \alpha = \pi - (\beta + \gamma). \quad \therefore \alpha + \beta + \gamma = \pi. \quad \therefore 2\alpha + 2\beta + 2\gamma = 2\pi.$$

$$\therefore \tan(2\alpha + 2\beta + 2\gamma) = \tan 2\pi = 0.$$

Therefore, as in Ex. 5, above,

$$\tan 2\alpha + \tan 2\beta + \tan 2\gamma = \tan 2\alpha \tan 2\beta \tan 2\gamma.$$

Now, expressing  $\tan 2\alpha$ ,  $\tan 2\beta$ ,  $\tan 2\gamma$  in terms of  $\tan \alpha$ ,  $\tan \beta$ ,  $\tan \gamma$ , and substituting  $x, y, z$ , for them, we get,

$$\frac{2x}{1-x^2} + \frac{2y}{1-y^2} + \frac{2z}{1-z^2} = \frac{8xyz}{(1-x^2)(1-y^2)(1-z^2)}.$$

On simplification, the required result follows.

### Examples X

If  $A + B + C = \pi$ , prove that (Ex. 1 to 16) :—

- ✓ 1.  $\sin A + \sin B - \sin C = 4 \sin \frac{A}{2} \sin \frac{B}{2} \cos \frac{C}{2}$ .
- ✓ 2.  $\cot B \cot C + \cot C \cot A + \cot A \cot B = 1$ .

3.  $\cot \frac{A}{2} + \cot \frac{B}{2} + \cot \frac{C}{2} = \cot \frac{A}{2} \cot \frac{B}{2} \cot \frac{C}{2}.$
4.  $\tan 2A + \tan 2B + \tan 2C = \tan 2A \tan 2B \tan 2C.$
5.  $(\cot B + \cot C)(\cot C + \cot A)(\cot A + \cot B)$   
 $= \operatorname{cosec} A \operatorname{cosec} B \operatorname{cosec} C.$
- ✓ 6.  $\frac{\cot B + \cot C}{\tan B + \tan C} + \frac{\cot C + \cot A}{\tan C + \tan A} + \frac{\cot A + \cot B}{\tan A + \tan B} = 1.$
7.  $\sin \frac{A}{2} + \sin \frac{B}{2} + \sin \frac{C}{2}$   
 $= 1 + 4 \sin \frac{\pi - A}{4} \sin \frac{\pi - B}{4} \sin \frac{\pi - C}{4}$   
 $= 1 + 4 \sin \frac{B+C}{4} \sin \frac{C+A}{4} \sin \frac{A+B}{4}.$
8.  $\cos^2 2A + \cos^2 2B + \cos^2 2C$   
 $= 1 + 2 \cos 2A \cos 2B \cos 2C.$
- ✓ 9.  $\sin^2 A + \sin^2 B + \sin^2 C = 2 + 2 \cos A \cos B \cos C.$
10.  $\sin^2 \frac{A}{2} + \sin^2 \frac{B}{2} + \sin^2 \frac{C}{2} = 1 - 2 \sin \frac{A}{2} \sin \frac{B}{2} \sin \frac{C}{2}.$
11.  $\frac{\cos A}{\sin B \sin C} + \frac{\cos B}{\sin C \sin A} + \frac{\cos C}{\sin A \sin B} = 2.$   
[ C. U. 1949. ]
12.  $\frac{\sin 2A + \sin 2B + \sin 2C}{\sin A + \sin B + \sin C} = 8 \sin \frac{A}{2} \sin \frac{B}{2} \sin \frac{C}{2}.$
- ✓ 13.  $\sin (B + C - A) + \sin (C + A - B) + \sin (A + B - C)$   
 $= 4 \sin A \sin B \sin C.$
14.  $\sin (B + 2C) + \sin (C + 2A) + \sin (A + 2B)$   
 $= 4 \sin \frac{B-C}{2} \sin \frac{C-A}{2} \sin \frac{A-B}{2}.$
- ✓ 15.  $\cos^2 A + \cos^2 B + 2 \cos A \cos B \cos C = \sin^2 C.$
16.  $\cos \frac{A}{2} \cos \frac{B-C}{2} + \cos \frac{B}{2} \cos \frac{C-A}{2} + \cos \frac{C}{2} \cos \frac{A-B}{2}$   
 $= \sin A + \sin B + \sin C.$

17. If  $\alpha + \beta + \gamma = \frac{1}{2}\pi$ , prove that

$$\checkmark (i) \sin^2 \alpha + \sin^2 \beta + \sin^2 \gamma + 2 \sin \alpha \sin \beta \sin \gamma = 1.$$

[ C. U. 1943. ]

$$\checkmark (ii) \tan \beta \tan \gamma + \tan \gamma \tan \alpha + \tan \alpha \tan \beta = 1.$$

18. If  $A, B, C, D$  are the angles of a quadrilateral, prove that

$$(i) \frac{\tan A + \tan B + \tan C + \tan D}{\cot A + \cot B + \cot C + \cot D} = \tan A \tan B \tan C \tan D.$$

$$(ii) \cos A + \cos B + \cos C + \cos D = 4 \cos \frac{1}{2}(A+B) \cos \frac{1}{2}(B+C) \cos \frac{1}{2}(C+A).$$

19. Show that

$$(i) \cos^2(\beta - \gamma) + \cos^2(\gamma - \alpha) + \cos^2(\alpha - \beta) = 1 + 2 \cos(\beta - \gamma) \cos(\gamma - \alpha) \cos(\alpha - \beta).$$

$$(ii) \sin^2 \alpha + \sin^2 \beta + 2 \sin \alpha \sin \beta \cos(\alpha + \beta) = \sin^2(\alpha + \beta).$$

$$(iii) \cos^2 \theta + \cos^2(\alpha + \theta) - 2 \cos \alpha \cos \theta \cos(\alpha + \theta) \text{ is independent of } \theta.$$

20. (i)  $\alpha + \beta = \gamma$ , show that

$$\cos^2 \alpha + \cos^2 \beta + \cos^2 \gamma = 1 + 2 \cos \alpha \cos \beta \cos \gamma.$$

[ C. U. 1940. ]

(ii) If  $\alpha + \beta + \gamma = 2\pi$ , show that

$$\cos^2 \alpha + \cos^2 \beta + \cos^2 \gamma - 2 \cos \alpha \cos \beta \cos \gamma = 1.$$

21. If  $\cos(A+B) \sin(C+D) = \cos(A-B) \sin(C-D)$ , show that

$$\cot A \cot B \cot C = \cot D.$$

22. If  $A+B+C=2S$ , prove that

$$(i) \sin(S-A) + \sin(S-B) + \sin(S-C) - \sin S = 4 \sin \frac{A}{2} \sin \frac{B}{2} \sin \frac{C}{2}.$$

$$(ii) \cos^2 A + \cos^2 B + \cos^2 C + 2 \cos A \cos B \cos C - 1 = 4 \cos S \cos(S-A) \cos(S-B) \cos(S-C).$$

23. If  $A + B + C = n\pi$  ( $n$  being zero or an integer),  
 $\tan A + \tan B + \tan C = \tan A \tan B \tan C$ .
24. Show that, if  $\alpha + \beta + \gamma = \pi$ ,  
 $\tan(\beta + \gamma - \alpha) + \tan(\gamma + \alpha - \beta) + \tan(\alpha + \beta - \gamma)$   
 $= \tan(\beta + \gamma - \alpha) \tan(\gamma + \alpha - \beta) \tan(\alpha + \beta - \gamma)$ .
25. If  $A + B + C = \pi$ , prove that  
 (i)  $\sin A \cos B \cos C + \sin B \cos C \cos A$   
 $+ \sin C \cos A \cos B = \sin A \sin B \sin C$ .  
 (ii)  $\cos A \sin B \sin C + \cos B \sin C \sin A$   
 $+ \cos C \sin A \sin B = 1 + \cos A \cos B \cos C$ .  
 (iii)  $\sin 5A + \sin 5B + \sin 5C$   
 $= 4 \cos \frac{5A}{2} \cos \frac{5B}{2} \cos \frac{5C}{2}$ .  
 (iv)  $(\tan A + \tan B + \tan C)(\cot A + \cot B + \cot C)$   
 $= 1 + \sec A \sec B \sec C$ .
- ✓ 26. If  $\cos A + \cos B + \cos C = 0$ , show that  
 $\cos 3A + \cos 3B + \cos 3C = 12 \cos A \cos B \cos C$ .  
 [Write  $\cos 3A = 4 \cos^3 A - 3 \cos A$ , etc.]
27. If  $x + y + z = \frac{1}{2}\pi$ , prove that  
 $\cos(x - y - z) + \cos(y - z - x) + \cos(z - x - y)$   
 $= 4 \cos x \cos y \cos z = 0$ .
28. Show that  
 $\sin(y - z) + \sin(z - x) + \sin(x - y)$   
 $+ 4 \sin \frac{y - z}{2} \sin \frac{z - x}{2} \sin \frac{x - y}{2} = 0$ .
29. If  $x + y + z = 0$ , show that  
 $\cot(z + x - y) \cot(x + y - z) + \cot(x + y - z) \cot(y + z - x)$   
 $+ \cot(y + z - x) \cot(z + x - y) = 1$ .
30. If  $x + y + z = xyz$ , prove that  
 $\frac{3x - x^3}{1 - 3x^3} + \frac{3y - y^3}{1 - 3y^3} + \frac{3z - z^3}{1 - 3z^3} = \frac{3x - x^3}{1 - 3x^3} \frac{3y - y^3}{1 - 3y^3} \frac{3z - z^3}{1 - 3z^3}$ .



## CHAPTER XI

### TRIGONOMETRICAL EQUATIONS AND GENERAL VALUES

57. It will be apparent from Chapter IV that there are infinitely many angles, the trigonometrical ratios of which have a given value. For example, if  $\sin \theta = \frac{1}{2}$ , one value of  $\theta$  (the smallest positive value) is known to be  $30^\circ$ . Now sines of supplementary angles are equal. Hence  $\sin 150^\circ$  being equal to  $\sin 30^\circ$  is also  $\frac{1}{2}$ . Again angles differing from  $30^\circ$  or  $150^\circ$  by complete multiples of  $360^\circ$  will have their sines (in fact all ratios) the same. Thus sine of each of the angles  $30^\circ$ ,  $150^\circ$ ,  $390^\circ$ ,  $510^\circ$ ,  $-330^\circ$ ,  $-210^\circ$ , etc. is equal to  $\frac{1}{2}$ . Similarly, if  $\cos \theta$  be given, equal to  $\frac{1}{\sqrt{2}}$  say,  $\theta$  may have any of the values  $+45^\circ$ ,  $+315^\circ$ ,  $+405^\circ$ ,  $-315^\circ$ ,  $-45^\circ$ , etc.; or else if  $\tan \theta = \sqrt{3}$ ,  $\theta$  may have any of the values  $60^\circ$ ,  $240^\circ$ ,  $420^\circ$ ,  $-300^\circ$ , etc.

It is very convenient for the solution of trigonometrical equations, as also for other purposes, to obtain a general expression in a compact form embracing all angles, the trigonometrical ratios of which have a given value.

58. General expression of all angles, one of whose trigonometrical ratios is zero.

If the sine of an angle be zero, from definition, the length of the perpendicular from any point of one of its arms upon another is zero, so that the two arms must be in the same straight line. Evidently, therefore, such angles must be some multiple of  $\pi$ , odd or even.

Thus, if  $\sin \theta = 0$ , then  $\theta = n\pi$ ,  
*n being zero, or any integer, positive or negative.*

When the cosine of an angle is zero, the projection of any length along one arm upon another is zero, and so the two arms must be at right angles to one another. The angles must therefore be evidently either  $\frac{\pi}{2}$  or  $\frac{3\pi}{2}$  or differ from these by complete revolutions; in other words, the angle may be any odd multiple of  $\frac{\pi}{2}$ .

Thus, if  $\cos \theta = 0$ , then  $\theta = (2n+1)\frac{\pi}{2}$ .

*n* being zero, or any integer, positive or negative.

Again if  $\tan \theta = 0$ , then its numerator  $\sin \theta$  is also zero;  
and so  $\theta = n\pi$ .

Similarly if  $\cot \theta = 0$ , then  $\cos \theta = 0$

and so  $\theta = (2n+1)\frac{\pi}{2}$ .

**Note.** The ratios  $\operatorname{cosec} \theta$  or  $\sec \theta$  can never be zero, for they can never be numerically less than unity.

**59. General expression of angles having the same sine (or cosecant).**

Let  $\alpha$  be any angle positive or negative such that its sine is equal to a given quantity  $k$  (numerically not greater than 1); for fixing up the idea, and for the sake of convenience in practice, the smallest positive angle having its sine for the given quantity  $k$  is taken as  $\alpha$ . Let  $\theta$  be any other angle whose sine is equal to  $k$ .

Then  $\sin \theta = \sin \alpha$ ,

or,  $\sin \theta - \sin \alpha = 0$ ,

or,  $2 \sin \frac{1}{2}(\theta - \alpha) \cos \frac{1}{2}(\theta + \alpha) = 0$ .

$\therefore$  either  $\sin \frac{1}{2}(\theta - \alpha) = 0$ ,

i.e.  $\frac{1}{2}(\theta - \alpha) = \text{any multiple of } \pi = m\pi, \dots (1)$

or, else  $\cos \frac{1}{2}(\theta + \alpha) = 0$ ,

i.e.  $\frac{1}{2}(\theta + \alpha) = \text{any odd multiple of } \frac{\pi}{2} = (2m+1)\frac{\pi}{2} \dots (2)$

From (1),  $\theta - \alpha = 2m\pi$ , i.e.  $\theta = \alpha + 2m\pi \dots (3)$

From (2),  $\theta + \alpha = (2m+1)\pi$ , i.e.  $\theta = -\alpha + (2m+1)\pi \dots (4)$

Combining (3) and (4),  $\theta = (-1)^n \alpha + n\pi \dots (5)$

where  $n$  is zero, or any integer, positive or negative, odd or even.

If  $\operatorname{cosec} \theta = \operatorname{cosec} \alpha$ , then  $\sin \theta = \sin \alpha$ ; hence all angles having the same cosecant as that of  $\alpha$  are also given by expression (5).

Thus all angles having the same sine or cosecant as that of  $\alpha$  are given by  $2n\pi + \alpha$  and  $(2n+1)\pi - \alpha$ ,

$$\text{or, } n\pi + (-1)^n \alpha,$$

$n$  being zero, or any integer, positive or negative.

**60. General expression of angles having the same cosine (or secant).**

Let  $\alpha$  be the smallest positive angle such that its cosine is equal to a given quantity  $k$  (numerically  $> 1$ ); and let  $\theta$  be any other angle whose cosine is equal to  $k$ .

Then,  $\cos \theta = \cos \alpha$ ,

or,  $\cos \alpha - \cos \theta = 0$ .

$\therefore 2 \sin \frac{1}{2}(\theta + \alpha) \sin \frac{1}{2}(\theta - \alpha) = 0$ .

$\therefore$  either  $\sin \frac{1}{2}(\theta + \alpha) = 0$ ,

i.e.  $\frac{1}{2}(\theta + \alpha) = \text{any multiple of } \pi = n\pi \dots (1)$

or else,  $\sin \frac{1}{2}(\theta - \alpha) = 0$ ,

i.e.  $\frac{1}{2}(\theta - \alpha) = \text{any multiple of } \pi = n\pi \dots (2)$

From (1),  $\theta + \alpha = 2n\pi$ , or  $\theta = 2n\pi - \alpha \dots (3)$

From (2),  $\theta - \alpha = 2n\pi$ , or  $\theta = 2n\pi + \alpha \dots (4)$

From (3) and (4), we have  $\theta = 2n\pi \pm \alpha$ , ... (5)  
 where  $n$  is zero, or any integer, positive or negative.

It is also evident as in the previous case that all angles having the same secant as that of  $\alpha$  are also included in the expression (5).

Hence, *all angles having the same cosine or secant as that of  $\alpha$  are given by*

$$2n\pi \pm \alpha$$

*$n$  being zero, or any integer, positive or negative.*

**Note.** As in Art. 59, instead of taking the smallest positive angle, we might take  $\alpha$  to be any one angle having for its cosine the given quantity  $k$ . The general value of  $\theta$  satisfying  $\cos \theta = \cos \alpha$  as obtained above, would not be affected at all.

### 61. General expression of all angles having the same tangent (or cotangent).

Let  $\alpha$  be the smallest positive angle such that its tangent is equal to a given quantity  $k$ ; and let  $\theta$  be any other angle whose tangent is equal to  $k$ .

$$\text{Then} \quad \tan \theta = \tan \alpha,$$

$$\text{or,} \quad \frac{\sin \theta}{\cos \theta} - \frac{\sin \alpha}{\cos \alpha} = 0,$$

$$\text{or,} \quad \frac{\sin \theta \cos \alpha - \cos \theta \sin \alpha}{\cos \theta \cos \alpha} = 0,$$

$$\text{or,} \quad \frac{\sin (\theta - \alpha)}{\cos \theta \cos \alpha} = 0.$$

$$\therefore \sin (\theta - \alpha) = 0,$$

$$\text{i.e.} \quad \theta - \alpha = \text{any multiple of } \pi = n\pi.$$

$$\therefore \theta = \alpha + n\pi. \quad \dots \quad (1)$$

The factor  $\frac{1}{\cos \theta \cos \alpha}$  cannot be zero, for cosine of an angle cannot have an infinitely large value.

It is also evident as in the previous case that all angles having the same cotangent as that of  $\alpha$  are given by the expression (1).

Hence, *all angles having the same tangent or cotangent as that of  $\alpha$  are given by*

$$n\pi + \alpha$$

*n being zero, or any integer, positive or negative.*

**Note.** The remark below Art. 60 is applicable here also.

## 62. Special cases.

From Art. 59, considering both cases when  $n$  is odd or even, it may be easily seen that

$$\text{if } \sin \theta = 1 = \sin \frac{\pi}{2}, \quad \theta = 2n\pi + \frac{\pi}{2} = (4n+1) \frac{\pi}{2}$$

$$\text{and if } \sin \theta = -1 = \sin \left( -\frac{\pi}{2} \right), \quad \theta = 2n\pi - \frac{\pi}{2} = (4n-1) \frac{\pi}{2}$$

$$\text{or,} \quad = (4k+3) \frac{\pi}{2}$$

where  $n$  (or  $k=n-1$ ) is zero, or any integer, positive or negative.

Similarly from Art. 60, it may be seen that

$$\text{if } \cos \theta = 1, \quad \theta = 2n\pi$$


$$\text{and if } \cos \theta = -1, \quad \theta = (2n+1)\pi$$

*n being zero, or any integer, positive or negative.*

These are the usual forms in which the above special cases are used in practice.

**63. Geometrical Treatment.**

(i) *Geometrical construction of an angle whose sine (or cosecant) is given, and to obtain a general expression of all such angles.*



Let the sine of an angle be given equal to ' $a$ '.

Taking the perpendicular lines  $XOX'$  and  $YOY'$  for reference, draw a circle of unit radius with centre  $O$ .

Measure off  $ON = a$  along  $OY$  (or along  $OY'$  if  $a$  be negative). Through  $N$  draw a straight line  $PNQ$  parallel to  $XOX'$  meeting the circle at  $P$  and  $Q$ .

Then  $\angle POX = a$  say, is one of the required angles, for  
 $\sin a = \sin OPN = \frac{ON}{OP} = \frac{a}{1} = a$ .

Another angle with the same sine, as is apparent from the figure, is  $\angle QOX = \pi - a$  (or  $3\pi - a$  if  $a = ON$  be negative, which is trigonometrically the same as  $\pi - a$ ).


' $a$ ' being given in magnitude and sign, the position of  $N$  on  $YOY'$  is fixed and thus in one revolution, i.e., from 0 to

$2\pi$  there are, as is clear from the figure, only two angles  $\alpha$  and  $\pi - \alpha$  having the given sine.\*

Now the addition or subtraction of any multiple of  $2\pi$  makes no difference in the values of the trigonometrical ratios of an angle (*See Art. 28*).

Hence all the angles having the same sine as that of  $\alpha$  are contained in the formulæ  $2m\pi + \alpha$  and  $2m\pi + \pi - \alpha$  i.e.,  $(2m + 1)\pi - \alpha$ , where  $m$  is zero, or any integer, positive or negative. Both the sets of angles are evidently included in the formula  $n\pi + (-1)^n\alpha$ ,  $n$  being zero, or any integer, positive or negative.

(ii) *Angles with given cosine (or secant).*




Let the given cosine be ' $a$ '. As before, measure off  $ON = a$  along  $OX$  (or along  $OX'$  if ' $a$ ' be negative), and through  $N$  draw  $PNQ$  parallel to  $YOY'$  to meet the circle with centre  $O$ , and radius unity, at  $P$  and  $Q$ .

\*In the same quadrant there cannot be two distinct angles (without being coterminals), having the same sine, for the corresponding triangles will then be congruent.

Let  $\angle POX = \alpha$ . Then  $\alpha$  is a required angle. Also from the figure, the only angles in the first four quadrants which have the given cosine are  $\alpha$  and  $2\pi - \alpha$ .

Adding or subtracting multiples of  $2\pi$  to these, all the angles having the same cosine as that of  $\alpha$  are given by  $2m\pi + \alpha$  or  $2m\pi + 2\pi - \alpha$ , both of which are included in the formula  $2n\pi \pm \alpha$ ,  $n$  being zero, or any integer, positive or negative.

(iii) *Angles with given tangent (or cotangent).*



Let ' $a$ ' be the given tangent. Along  $OX$  or  $OX'$  measure off  $ON$  of unit length, and then measure off  $NP$  perpendicular to it of length whose numerical value is ' $a$ '. If ' $a$ ' be positive, both  $ON$  and  $NP$  will be positive, or both will be negative, and so the  $\angle XOP$  will be either in the first or in the third quadrant. If ' $a$ ' be negative, the angle will be either in the second or in the fourth quadrant. In any case, there are only two angles, within one revolution, i.e., from 0 to  $2\pi$  as is apparent from the figure, with the given tangent.\*

\*The ratio  $PN : ON$  being given, and the included angle  $PNO$  being right, the triangle  $PNO$  constructed remains always similar to itself and so in the same quadrant the  $\angle PON$  of the triangle is unique.



One of the angles being  $\alpha$ , the other is evidently (from the figure)  $\pi + \alpha$ . Adding or subtracting multiples of  $2\pi$ , all the angles having the same tangent as that of  $\alpha$  are given by  $2m\pi + \alpha$  or  $2m\pi + \pi + \alpha$  both of which are included in the formula  $n\pi + \alpha$  where  $n$  is zero, or any integer, positive or negative, odd or even.

**Ex. 1.** Solve  $2(\cos^2 \theta - \sin^2 \theta) = 1$ .

The given equation can be written as

$$2 \cos 2\theta = 1. \quad \therefore \cos 2\theta = \frac{1}{2} = \cos \frac{1}{3}\pi.$$

$$\therefore 2\theta = 2n\pi \pm \frac{1}{3}\pi. \quad \therefore \theta = n\pi \pm \frac{1}{6}\pi.$$

**Note.** It may be observed that a trigonometrical equation can be solved in several ways; and the results though different in forms will give the same series of angles. To illustrate this we work out the above example in another way.

The equation can also be written in the form

$$2(\cos^2 \theta - 1 + \cos^2 \theta) = 1, \quad \text{or, } 4 \cos^2 \theta = 3.$$

$$\therefore \cos \theta = \pm \frac{\sqrt{3}}{2} = \cos \frac{\pi}{6}, \quad \text{or, } \cos \frac{5\pi}{6}.$$

$$\therefore \theta = 2m\pi \pm \frac{\pi}{6}, \quad \text{or, } 2m\pi \pm \frac{5\pi}{6}.$$

$$\text{Now } 2m\pi \pm \frac{5\pi}{6} = (2m+1)\pi - \frac{\pi}{6}, \quad \text{or, } (2m-1)\pi + \frac{\pi}{6}.$$

All the four sets of solutions,  $m$  being any integer, can be included in the expression  $n\pi \pm \frac{1}{6}\pi$ , in which form the result has already been obtained by the previous process.

**Ex. 2.** Solve  $4 \cos^2 x + 6 \sin^2 x = 5$ .

The equation can be written as

$$4 \cos^2 x + 6 \sin^2 x = 5(\sin^2 x + \cos^2 x).$$

$$\therefore \sin^2 x = \cos^2 x, \quad \text{or, } \tan^2 x = 1.$$

$$\therefore \tan x = \pm 1. \quad \therefore x = n\pi \pm \frac{1}{4}\pi.$$

**Note.** Equations of the form  $a \cos^2 x + b \sin^2 x = c$  can be easily solved by the above method, or by expressing sine in terms of cosine or cosine in terms of sine.

**Ex. 3.** Solve  $2 \sin^2 x + \sin^2 2x = 2$ . [C. U. 1940.]

The given equation can be written as

$$2(1 - \sin^2 x) - \sin^2 2x = 0, \quad \text{or, } 2 \cos^2 x - 4 \sin^2 x \cos^2 x = 0.$$

$$\text{or, } 2 \cos^2 x (1 - 2 \sin^2 x) = 0, \quad \text{or, } \cos^2 x \cos 2x = 0.$$

$$\therefore \text{either } \cos x = 0, \text{ i.e., } x = n\pi + \frac{1}{2}\pi, \text{ or } (2n+1)\frac{\pi}{2}$$

$$\text{or, } \cos 2x = 0, \text{ i.e., } 2x = 2n\pi \pm \frac{1}{2}\pi. \quad \therefore x = n\pi \pm \frac{1}{4}\pi.$$

$$\text{Ex. 4. Solve } \cos \theta - \sin \theta = \frac{1}{\sqrt{2}}.$$

Dividing both sides of the equation by  $\sqrt{1^2 + 1^2}$  i.e.,  $\sqrt{2}$ , we have

$$\cos \theta \cdot \frac{1}{\sqrt{2}} - \sin \theta \cdot \frac{1}{\sqrt{2}} = \frac{1}{2}$$

$$\text{i.e., } \cos \theta \cos \frac{1}{4}\pi - \sin \theta \sin \frac{1}{4}\pi = \frac{1}{2}.$$

$$\therefore \cos (\theta + \frac{1}{4}\pi) = \cos \frac{1}{2}\pi. \quad \therefore \theta + \frac{1}{4}\pi = 2n\pi \pm \frac{1}{2}\pi.$$

$$\therefore \theta = 2n\pi + \frac{1}{4}\pi, \text{ or, } 2n\pi - \frac{7}{4}\pi.$$

**Note. Extraneous solutions.**

In general, as pointed out in Ex. 1 above, the same trigonometrical equation may be solved by different methods, and the forms of the result we arrive at, though apparently different in some cases, are ultimately equivalent. In some cases, however, we may be tempted to solve a trigonometrical equation by methods which have flaws in them, leading to solutions which include in addition to the correct solutions, some extraneous solutions which do not satisfy the given equation. The given equation which is of the type  $a \cos \theta + b \sin \theta = c$  is an example. We proceed to demonstrate it as follows:

$$\text{Here } \cos \theta - \frac{1}{\sqrt{2}} = \sin \theta.$$

$$\therefore \cos^2 \theta - \sqrt{2} \cos \theta + \frac{1}{2} = \sin^2 \theta = 1 - \cos^2 \theta,$$

$$\text{whence } 2 \cos^2 \theta - \sqrt{2} \cos \theta - \frac{1}{2} = 0.$$

$$\therefore \cos \theta = \frac{\sqrt{2} \pm \sqrt{2 + 4}}{4} = \frac{1 \pm \sqrt{3}}{2\sqrt{2}} = \cos \frac{\pi}{12} \quad \text{or} \quad \cos \frac{7\pi}{12}.$$

$$\therefore \theta = 2n\pi \pm \frac{1}{12}\pi, \text{ or } 2n\pi \pm \frac{7}{12}\pi.$$

But it can be easily seen on substitution that

$2n\pi - \frac{1}{12}\pi$  and  $2n\pi + \frac{7}{12}\pi$  do not satisfy the given equation. The error in the method lies in squaring the equation as we have done; for the squared equation includes the equation  $\cos \theta - \frac{1}{\sqrt{2}} = -\sin \theta$  i.e.,  $\cos \theta + \sin \theta = \frac{1}{\sqrt{2}}$  of which the solutions are  $2n\pi - \frac{1}{12}\pi$  and  $2n\pi + \frac{7}{12}\pi$ .

Equations of this type are therefore best solved as in the next example, and not by squaring.

Thus while solving any trigonometrical equation it is always advisable to verify the roots obtained ; for thereby extraneous roots, if any, can be easily detected.

**Ex. 5.** Solve  $a \cos \theta + b \sin \theta = c$ . ( $c > \sqrt{a^2 + b^2}$ .)

Put  $a = r \cos \alpha$ ,  $b = r \sin \alpha$ , choosing the smallest positive value of  $\alpha$ , keeping  $r$  positive.

$$\text{Then } r = \sqrt{a^2 + b^2} \text{ and } \sin \alpha = \frac{b}{\sqrt{a^2 + b^2}}$$

$$\text{and } \cos \alpha = \frac{a}{\sqrt{a^2 + b^2}}.$$

The signs of  $a$  and  $b$  will determine the quadrant in which  $\alpha$  lies, and  $a$  and  $b$  being given,  $r$  and  $\alpha$  are definitely known.

The equation now becomes,

$$r \cos (\theta - \alpha) = c,$$

$$\text{or, } \cos (\theta - \alpha) = \frac{c}{\sqrt{a^2 + b^2}} = \cos \beta,$$

where  $\beta$  is the smallest positive angle whose cosine is  $\frac{c}{\sqrt{a^2 + b^2}}$ , and  $a, b, c$  being known,  $\beta$  is also known.

Hence  $\theta - \alpha = 2n\pi \pm \beta$ , or,  $\theta = 2n\pi + \alpha \pm \beta$ .

**Note.** An angle which is introduced in a trigonometrical work to facilitate calculations is called a *subsidiary* angle. Thus  $\alpha$  and  $\beta$  are here subsidiary angles.

**Ex. 6.** Solve  $4 \cos x + 5 \sin x = 5$ , given  $\tan 51^\circ 21' = \frac{5}{4}$ .

Dividing both sides of the given equation by  $\sqrt{4^2 + 5^2}$  i.e. by  $\sqrt{41}$ , we get

$$\frac{4}{\sqrt{41}} \cos x + \frac{5}{\sqrt{41}} \sin x = \frac{5}{\sqrt{41}}. \quad \dots (1)$$

Since,  $\tan 51^\circ 21' = \frac{5}{4}$ ,

$$\therefore \sin 51^\circ 21' = \frac{5}{\sqrt{41}}, \cos 51^\circ 21' = \frac{4}{\sqrt{41}}.$$

$\therefore$  (1) reduces to

$$\cos x \cos 51^\circ 21' + \sin x \sin 51^\circ 21' = \sin 51^\circ 21',$$

$$\text{or, } \cos (x - 51^\circ 21') = \sin 51^\circ 21' = \cos 38^\circ 39'.$$

$$\therefore x - 51^\circ 21' = 2n\pi \pm 38^\circ 39'.$$

$$\therefore x = 2n\pi + 90^\circ, \text{ or, } 2n\pi + 12^\circ 42'.$$

**Ex. 7.** (i) Solve  $2 \sin^2 x + \sin^2 2x = 2$  for  $-\pi < x < \pi$ .

From Ex. 3 above, we see that  $x = n\pi + \frac{1}{2}\pi \dots$  (1)

$$\text{or, } x = n\pi \pm \frac{1}{4}\pi. \dots \dots (2)$$

Putting  $n = 0, -1$  in (1), we get  $x = \frac{1}{2}\pi, -\frac{1}{2}\pi$ , which lie in the given interval. Putting  $n = 0, 1, -1$  in (2), we get  $x = \pm \frac{1}{4}\pi, \frac{3}{4}\pi, -\frac{3}{4}\pi$  which also lie in the given interval.

Hence the required values of  $x$  are  $\pm \frac{1}{4}\pi, \pm \frac{1}{2}\pi, \pm \frac{3}{4}\pi$ .

(ii) Solve  $\cos \theta + \sqrt{3} \sin \theta = 2$

for  $-2\pi < \theta < 2\pi$  and  $3\pi < \theta < 5\pi$ .

Dividing both sides of the equation by  $\sqrt{1+3}$  i.e., 2, we have

$$\cos \theta \cdot \frac{1}{2} + \sin \theta \cdot \frac{\sqrt{3}}{2} = 1,$$

$$\text{i.e., } \cos \theta \cos \frac{1}{2}\pi + \sin \theta \sin \frac{1}{2}\pi = 1,$$

$$\text{i.e., } \cos (\theta - \frac{1}{2}\pi) = 1.$$

$$\therefore \theta - \frac{1}{2}\pi = 2n\pi, \text{ i.e., } \theta = 2n\pi + \frac{1}{2}\pi.$$

Putting  $n = 0, -1$ , we get  $\theta = \frac{1}{2}\pi, -\frac{1}{2}\pi$  which lie in the 1st interval.

Again putting  $n = 1, 2$ , we get  $\theta = \frac{5}{2}\pi, \frac{9}{2}\pi$ , which lie in the 2nd interval.

**Ex. 8.** Solve  $\tan ax = \cot bx$ .

Here,  $\tan ax = \cot bx = \tan (\frac{1}{2}\pi - bx)$ .

$$\therefore ax = n\pi + \frac{1}{2}\pi - bx,$$

$$\therefore x = \frac{2n+1}{a+b} \cdot \frac{\pi}{2}.$$

## Examples XI

Solve the following equations (*Ex. 1 to 23*) :—

1.  $\cot^2 x + \operatorname{cosec}^2 x = 3$ .
2. (i)  $2 \cos^2 \theta + 4 \sin^2 \theta = 3$ .  
(ii)  $\tan^2 \theta = 3 \operatorname{cosec}^2 \theta - 1$ . [ *C. U. 1939.* ]
3.  $\tan x - \cot x = \operatorname{cosec} x$ .
4.  $\cot x - \cot 2x = 2$ .
5.  $2 \sin \theta \tan \theta + 1 = \tan \theta + 2 \sin \theta$ .
6.  $\sin 5\theta + \sin \theta = \sin 3\theta$ .
7.  $\sin m\theta + \sin n\theta = 0$ .
8.  $\cos x + \cos 3x + \cos 5x + \cos 7x = 0$ .
9.  $\cot 2x = \cos x + \sin x$ .
10.  $\sin x + \cos x = \sqrt{2}$ , for  $-\pi < x < \pi$ .
11.  $\sin 2x \tan x + 1 = \sin 2x + \tan x$ .
12.  $\cot x - \tan x = 2$ . [ *C. U. 1934, '37.* ]
13.  $\sin x + \sqrt{3} \cos x = \sqrt{2}$ . [ *C. U. 1938, '47.* ]
14.  $2 \sin x \sin 3x = 1$ .
15.  $\sin \theta + 2 \cos \theta = 1$ . [ *C. U. 1933.* ]
16.  $\tan x + \tan 2x + \tan 3x = \tan x \tan 2x \tan 3x$ .
17.  $\tan (\frac{1}{2}\pi + \theta) + \tan (\frac{1}{2}\pi - \theta) = 4$ . [ *C. U. 1949.* ]
18.  $\tan x + \tan 2x + \tan x \tan 2x = 1$ . [ *C. U. 1941, '45.* ]
19.  $\cos \theta + \sqrt{3} \sin \theta = \sqrt{2}$ . [ *C. U. 1944.* ]
20.  $\sqrt{3} \cos x + \sin x = 1$ , for  $-2\pi < x < 2\pi$ .
21.  $\cos 2x = \cos x \sin x$ .
22.  $2 \cot x + \sin x = 2 \operatorname{cosec} x$ .
23.  $\cos x + \sin x = \cos 2x + \sin 2x$ . [ *C. U. 1943.* ]
24. Solve  $2 \sin^2 x + \sin x = 3$ ; and find all the angles between  $0^\circ$  and  $1000^\circ$  which satisfy it.
25. Find the solution of the equations (general solution not required)  

$$\tan x + \tan y = 2$$

$$2 \cos x \cos y = 1.$$

(26) If  $\tan ax - \tan bx = 0$ , show that the values of  $x$  form a series in A. P.

27. Solve

(i)  $\cos 3x + \cos 2x + \cos x = 0$ . [C. U. 1942, '46.]

(ii)  $\cos 9x \cos 7x = \cos 5x \cos 3x$ ,  $-\frac{1}{2}\pi < x < \frac{1}{2}\pi$ .

(iii)  $\tan x + \tan 2x + \tan 3x = 0$ . [A. I. 1941.]

(iv)  $\cos x - \sin x = \cos a + \sin a$ . [B. H. U. 1938.]

(v)  $\cos^3 x - \cos x \sin x - \sin^3 x = 1$ .

(vi)  $\cos 6x + \cos 4x = \sin 3x + \sin x$ .

(vii)  $\frac{\sin a}{\sin 2x} + \frac{\cos a}{\cos 2x} = 2$ .

28. Solve  $5 \cos \theta + 2 \sin \theta = 2$ , given  $\tan 68^\circ 12' = 2\frac{1}{2}$ .

29. Find those pairs of solutions of the following equations which correspond to positive solutions less than  $2\pi$  of each individual equation :—

(i)  $\sin (a - \beta) = 0$  ;  $\sin (a + \beta) = 1$ .

(ii)  $\sin (a - \beta) = \cos (a + \beta) = \frac{1}{2}$ .

30. If  $\sin A = \sin B$ ,  $\cos A = \cos B$ , prove that either  $A$  and  $B$  are equal or they differ by some multiple of four right angles. [C. U. 1935]

31. Show that the three equations

$$\sin^2 \theta = \sin^2 a, \cos^2 \theta = \cos^2 a, \tan^2 \theta = \tan^2 a$$

are all identical and the solution is always  $n\pi \pm a$ .

32. Show that the same two series of angles are given by the equations

$$x + \frac{\pi}{4} = n\pi + (-1)^n \frac{\pi}{6} \quad \text{and} \quad x - \frac{\pi}{4} = 2n\pi \pm \frac{\pi}{3}.$$

## CHAPTER XII

### INVERSE CIRCULAR FUNCTIONS

64. The equation  $\sin \theta = x$  means that  $\theta$  is an angle whose sine is  $x$ . It is often convenient to express this statement *inversely* by writing  $\theta = \sin^{-1}x$ . Thus the symbol  $\sin^{-1}x$  denotes an angle whose sine is  $x$ . Hence  $\sin^{-1}x$  is an angle, whereas  $\sin \theta$  is a number. The two relations  $\sin \theta = x$  and  $\theta = \sin^{-1}x$  are identical; if one is given the other follows. The symbol  $\sin^{-1}x$  is usually read as "*sine inverse x*". Sometimes it is also denoted by *arc sin x*.

**Note.**  $\sin^{-1}x$  must not be confused with  $(\sin x)^{-1}$  i.e.  $\frac{1}{\sin x}$ .

65. We know that if  $\theta$  be any one angle whose sine is equal to  $x$ , then sines of all the angles given by  $n\pi + (-1)^n\theta$  are equal to  $x$ . Hence,  $\sin^{-1}x$  has got an infinite number of values, and as such,  $\sin^{-1}x$  is a *multiple-valued function*.

Hence the *general value* of  $\sin^{-1}x = n\pi + (-1)^n \sin^{-1}x$  where on the right-hand side  $\sin^{-1}x$  stands for any particular angle whose sine is  $x$ .

Similarly, the *general value* of

$$\cos^{-1}x = 2n\pi \pm \cos^{-1}x$$

$$\text{and of } \tan^{-1}x = n\pi + \tan^{-1}x.$$

The smallest numerical value, either positive or negative, of  $\theta$  is called the *principal value* of  $\sin^{-1}x$ . Thus, the principal value of  $\sin^{-1}\frac{1}{2}$  is  $30^\circ$ . If corresponding to the same ratio, there are two numerically equal angles, one positive and the other negative, it is customary to take the positive angle as the principal value; thus the principal value of  $\cos^{-1}\frac{1}{2}$  is  $60^\circ$ , and not  $(-60^\circ)$  although  $\cos(-60^\circ) = \frac{1}{2}$ .

In all numerical examples, the principal value is generally taken.

$\cos^{-1}x$ ,  $\tan^{-1}x$ ,  $\operatorname{cosec}^{-1}x$ ,  $\sec^{-1}x$ ,  $\cot^{-1}x$  have similar significance and all properties as those of  $\sin^{-1}x$ . These expressions are called **Inverse Circular Functions**.

66. If  $\sin \theta = x$ , then  $\theta = \sin^{-1}x$ , i.e.  $\theta = \sin^{-1} \sin \theta$ .

Similarly,  $\theta = \cos^{-1} \cos \theta = \tan^{-1} \tan \theta$ ; etc.

Again, if  $\theta = \sin^{-1}x$ ,  $\sin \theta = x$ , i.e.  $\sin \sin^{-1}x = x$ .

Similarly,  $\cos \cos^{-1}x = x$ ;  $\tan \tan^{-1}x = x$ ; etc.

Also, we have

$$\operatorname{cosec}^{-1}x = \sin^{-1} \frac{1}{x}; \cot^{-1}x = \tan^{-1} \frac{1}{x}; \sec^{-1}x = \cos^{-1} \frac{1}{x}$$

Let  $\operatorname{cosec}^{-1}x = \theta$ ; then  $\operatorname{cosec} \theta = x$ .

$$\therefore \sin \theta = \frac{1}{\operatorname{cosec} \theta} = \frac{1}{x}.$$

Hence  $\theta = \sin^{-1} \frac{1}{x}$ , and therefore  $\operatorname{cosec}^{-1}x = \sin^{-1} \frac{1}{x}$ .

In the same way we have,  $\sec^{-1} \frac{1}{x} = \cos^{-1}x$ .

The other relations follow similarly.

67. As all the trigonometrical ratios can be expressed in terms of any one, similarly, all the inverse trigonometrical ratios can be expressed in terms of any one inverse ratio.

Thus, let  $\sin^{-1}x = \theta$ ; then  $\sin \theta = x$ .

$$\therefore \cos \theta = \sqrt{1-x^2}; \tan \theta = \frac{x}{\sqrt{1-x^2}}; \cot \theta = \frac{\sqrt{1-x^2}}{x};$$

$$\sec \theta = \frac{1}{\sqrt{1-x^2}} \text{ and } \operatorname{cosec} \theta = \frac{1}{x}.$$

$$\begin{aligned} \therefore \theta = \sin^{-1}x &= \cos^{-1} \sqrt{1-x^2} = \tan^{-1} \frac{x}{\sqrt{1-x^2}} \\ &= \cot^{-1} \frac{\sqrt{1-x^2}}{x} = \sec^{-1} \frac{1}{\sqrt{1-x^2}} = \operatorname{cosec}^{-1} \frac{1}{x}. \end{aligned}$$



To prove that

$$(i) \sin^{-1}x + \cos^{-1}x = \frac{\pi}{2}.$$

$$(ii) \tan^{-1}x + \cot^{-1}x = \frac{\pi}{2}.$$

$$(iii) \operatorname{cosec}^{-1}x + \sec^{-1}x = \frac{\pi}{2}.$$

(i) Let  $\sin^{-1}x = \theta$  ; then  $\sin \theta = x$ .

Now,  $\sin \theta = \cos (\frac{1}{2}\pi - \theta)$ .

$\therefore \cos (\frac{1}{2}\pi - \theta) = x$  and hence  $\cos^{-1}x = \frac{1}{2}\pi - \theta$ .

Therefore,  $\sin^{-1}x + \cos^{-1}x = \theta + \frac{1}{2}\pi - \theta = \frac{1}{2}\pi$ .

(ii) Let  $\tan^{-1}x = \theta$  ; then  $\tan \theta = x$ .

Now,  $\tan \theta = \cot (\frac{1}{2}\pi - \theta)$ .

$\therefore \cot (\frac{1}{2}\pi - \theta) = x$ .  $\therefore \cot^{-1}x = \frac{1}{2}\pi - \theta$ .

$\therefore \tan^{-1}x + \cot^{-1}x = \theta + \frac{1}{2}\pi - \theta = \frac{1}{2}\pi$ .

(iii) Let  $\operatorname{cosec}^{-1}x = \theta$  ; then  $\operatorname{cosec} \theta = x$ .

Now,  $\operatorname{cosec} \theta = \sec (\frac{1}{2}\pi - \theta)$ .

$\therefore \sec (\frac{1}{2}\pi - \theta) = x$ .  $\therefore \sec^{-1}x = \frac{1}{2}\pi - \theta$ .

$\therefore \operatorname{cosec}^{-1}x + \sec^{-1}x = \theta + \frac{1}{2}\pi - \theta = \frac{1}{2}\pi$ .

69. To prove that

(i)  $\tan^{-1}x + \tan^{-1}y = \tan^{-1} \frac{x+y}{1-xy}$ .

(ii)  $\tan^{-1}x - \tan^{-1}y = \tan^{-1} \frac{x-y}{1+xy}$ .

Let  $\tan^{-1}x = \alpha$  ; and  $\tan^{-1}y = \beta$  ;

then  $\tan \alpha = x$  ; and  $\tan \beta = y$ .

Now,  $\tan (\alpha + \beta) = \frac{\tan \alpha + \tan \beta}{1 - \tan \alpha \tan \beta} = \frac{x+y}{1-xy}$ .

$$\therefore a + \beta = \tan^{-1} \frac{x+y}{1-xy}$$

$$\text{i.e. } \tan^{-1}x + \tan^{-1}y = \tan^{-1} \frac{x+y}{1-xy}$$

$$\text{Again, } \tan(a - \beta) = \frac{\tan a - \tan \beta}{1 + \tan a \tan \beta} = \frac{x-y}{1+xy}$$

$$\therefore a - \beta = \tan^{-1} \frac{x-y}{1+xy}$$

$$\text{i.e. } \tan^{-1}x - \tan^{-1}y = \tan^{-1} \frac{x-y}{1+xy}$$

Note. It can be easily proved as above that

$$\cot^{-1}x \pm \cot^{-1}y = \cot^{-1} \frac{xy \mp 1}{y \pm x}$$

70. To prove that

$$\tan^{-1}x + \tan^{-1}y + \tan^{-1}z = \tan^{-1} \frac{x+y+z-xyz}{1-yz-zx-xy}$$

$$\text{Let } \tan^{-1}x = a; \tan^{-1}y = \beta; \tan^{-1}z = \gamma.$$

$$\therefore \tan a = x, \quad \tan \beta = y, \quad \tan \gamma = z.$$

Now,  $\tan(a + \beta + \gamma)$

$$\begin{aligned} &= \frac{\tan a + \tan \beta + \tan \gamma - \tan a \tan \beta \tan \gamma}{1 - \tan \beta \tan \gamma - \tan \gamma \tan a - \tan a \tan \beta} \\ &= \frac{x + y + z - xyz}{1 - yz - zx - xy} \end{aligned}$$

$$\text{Hence, } a + \beta + \gamma = \tan^{-1} \frac{x+y+z-xyz}{1-yz-zx-xy}$$

Since,  $a + \beta + \gamma = \tan^{-1}x + \tan^{-1}y + \tan^{-1}z$ , the required result follows.

Note. This relation can also be deduced by applying twice the formula of Art. 69. Thus,

$$\text{Left side} = (\tan^{-1}x + \tan^{-1}y) + \tan^{-1}z$$

$$= \tan^{-1} \frac{x+y}{1-xy} + \tan^{-1}z; \text{ now again apply Art. 69.}$$

71. In fact for most of the formulæ involving ordinary circular functions, corresponding relations connecting the inverse circular functions can be easily deduced. In addition to those given above, some are illustrated in the following examples.

**Ex. 1.** *Show that*

$$(i) \sin^{-1}x \pm \sin^{-1}y = \sin^{-1}\{x\sqrt{1-y^2} \pm y\sqrt{1-x^2}\}.$$

$$(ii) \cos^{-1}x \pm \cos^{-1}y = \cos^{-1}\{xy \mp \sqrt{(1-x^2)(1-y^2)}\}.$$

(i) Let  $\sin^{-1}x = a$ .  $\therefore \sin a = x$  and  $\cos a = \sqrt{1-x^2}$ ;  
also let  $\sin^{-1}y = \beta$ .  $\therefore \sin \beta = y$  and  $\cos \beta = \sqrt{1-y^2}$ .

$$\begin{aligned} \text{Now, } \sin(a \pm \beta) &= \sin a \cos \beta \pm \cos a \sin \beta \\ &= x\sqrt{1-y^2} \pm y\sqrt{1-x^2}. \end{aligned}$$

$$\therefore a \pm \beta = \sin^{-1}\{x\sqrt{1-y^2} \pm y\sqrt{1-x^2}\}.$$

Since,  $a \pm \beta = \sin^{-1}x \pm \sin^{-1}y$ , the required result follows.

(ii) These relations follow similarly from the value of  $\cos(a \pm \beta)$ .

**Ex. 2.** *Show that*

$$(i) 2 \sin^{-1}x = \sin^{-1}(2x\sqrt{1-x^2}).$$

$$(ii) 2 \cos^{-1}x = \cos^{-1}(2x^2 - 1).$$

$$(iii) 2 \tan^{-1}x = \tan^{-1} \frac{2x}{1-x^2}.$$

(i) Let  $\sin^{-1}x = a$ .  $\therefore \sin a = x$ ,  $\cos a = \sqrt{1-x^2}$ .

$$\text{Now, } \sin 2a = 2 \sin a \cos a = 2x\sqrt{1-x^2}.$$

$$\therefore 2a = \sin^{-1}(2x\sqrt{1-x^2}).$$

Since,  $a = \sin^{-1}x$ , the required result follows.

(ii) & (iii). These relations follow similarly from the corresponding values of  $\cos 2a$  in terms of  $\cos a$  and  $\tan 2a$  in terms of  $\tan a$ . [See Art. 43]

**Note.** The above three relations can also be deduced by putting  $x$  for  $y$  in the values of  $\sin^{-1}x + \sin^{-1}y$ ,  $\cos^{-1}x + \cos^{-1}y$  and  $\tan^{-1}x + \tan^{-1}y$ .

**Ex. 3.** Show that

$$(i) \quad 3 \sin^{-1}x = \sin^{-1}(3x - 4x^3).$$

$$(ii) \quad 3 \cos^{-1}x = \cos^{-1}(4x^3 - 3x).$$

$$(iii) \quad 3 \tan^{-1}x = \tan^{-1} \frac{3x - x^3}{1 - 3x^2}. \quad [C. U. 1938.]$$

(i) Let  $\sin^{-1}x = \theta$ ; then  $\sin \theta = x$ .

$$\text{Now, } \sin 3\theta = 3 \sin \theta - 4 \sin^3 \theta = 3x - 4x^3.$$

$$\therefore \quad 3\theta, \text{ i.e. } 3 \sin^{-1}x = \sin^{-1}(3x - 4x^3).$$

(ii) & (iii). These relations follow similarly from the corresponding values of  $\cos 3\theta$  in terms of  $\cos \theta$  and of  $\tan 3\theta$  in terms of  $\tan \theta$ . [See Art. 44.]

**Note.** The result of (iii) may also be deduced by putting  $y = x = x$  in the formula of Art. 70.

**Ex. 4.** Show that

$$2 \tan^{-1}x = \sin^{-1} \frac{2x}{1+x^2} = \cos^{-1} \frac{1-x^2}{1+x^2} = \tan^{-1} \frac{2x}{1-x^2}.$$

$$\text{Let } \tan^{-1}x = \theta, \quad \therefore \tan \theta = x.$$

$$\text{Since, } \sin 2\theta = \frac{2 \tan \theta}{1 + \tan^2 \theta} = \frac{2x}{1+x^2}, \quad [\text{Art. 45, Ex. I.}]$$

$$\therefore \quad 2\theta \text{ i.e. } 2 \tan^{-1}x = \sin^{-1} \frac{2x}{1+x^2}.$$

$$\text{Since, } \cos 2\theta = \frac{1 - \tan^2 \theta}{1 + \tan^2 \theta} = \frac{1 - x^2}{1 + x^2},$$

$$\text{and } \tan 2\theta = \frac{2 \tan \theta}{1 - \tan^2 \theta} = \frac{2x}{1 - x^2},$$

the remaining relations follow similarly.

**Ex. 5.** Show that

$$\tan^{-1} \frac{a-b}{1+ab} + \tan^{-1} \frac{b-c}{1+bc} + \tan^{-1} \frac{c-a}{1+ca} = 0.$$

1st term of left side =  $\tan^{-1} a - \tan^{-1} b$  [By Art. 69 (ii)]

2nd ... .. =  $\tan^{-1} b - \tan^{-1} c$ .

3rd ... .. =  $\tan^{-1} c - \tan^{-1} a$ .

Hence, adding up the three terms, the required result follows.

**Ex. 6.** Show that

$$2 \tan^{-1} \frac{1}{5} + \tan^{-1} \frac{1}{4} = \tan^{-1} \frac{4}{3}.$$

Since,  $2 \tan^{-1} x = \tan^{-1} \frac{2x}{1-x^2}$ , [See Ex. 1.]

$$\therefore 2 \tan^{-1} \frac{1}{5} = \tan^{-1} \frac{\frac{2}{5}}{1 - \frac{1}{25}} = \tan^{-1} \frac{2}{12}.$$

$$\therefore \text{Left side} = \tan^{-1} \frac{2}{12} + \tan^{-1} \frac{1}{4} = \tan^{-1} \frac{\frac{2}{12} + \frac{1}{4}}{1 - \frac{2}{12} \cdot \frac{1}{4}} = \tan^{-1} \frac{4}{3}.$$

**Ex. 7.** Solve

$$\sin^{-1} \frac{2a}{1+a^2} + \sin^{-1} \frac{2b}{1+b^2} = 2 \tan^{-1} x.$$

[C. U. 1917.]

Since,  $\sin^{-1} \frac{2x}{1+x^2} = 2 \tan^{-1} x$ . [See Ex. 4.]

$$\therefore \text{Left side} = 2 \tan^{-1} a + 2 \tan^{-1} b.$$

$\therefore$  the equation reduces to

$$2 \tan^{-1} x = 2 \tan^{-1} a + 2 \tan^{-1} b.$$

$$\therefore \tan^{-1} x = \tan^{-1} a + \tan^{-1} b = \tan^{-1} \frac{a+b}{1-ab}.$$

$$\therefore x = \frac{a+b}{1-ab}.$$

Ex. 8. Solve

$$\tan^{-1} \frac{x-1}{x-2} + \tan^{-1} \frac{x+1}{x+2} = \frac{\pi}{4}.$$

$$\text{Left side} = \tan^{-1} \frac{\frac{x-1}{x-2} + \frac{x+1}{x+2}}{1 - \frac{x^2-1}{x^2-4}} = \tan^{-1} \frac{x^2-4}{-3}.$$

$\therefore$  the equation reduces to

$$\tan^{-1} \frac{2x^2-4}{-3} = \frac{\pi}{4} = \tan^{-1} 1.$$

$$\therefore \frac{2x^2-4}{-3} = 1 \text{ or, } 2x^2 = 1 \text{ or, } x = \pm \frac{1}{\sqrt{2}}.$$

### Examples XII

Prove (Ex. 1 to 17) that :—

1. (i)  $\tan^{-1} \frac{1}{2} + \tan^{-1} \frac{1}{3} = \frac{1}{4}\pi.$

(ii)  $\tan^{-1} x + \tan^{-1} \frac{2x}{1-x^2} = \tan^{-1} \frac{3x-x^3}{1-3x^2}.$

(iii)  $\tan^{-1} \frac{1}{7} + \tan^{-1} \frac{1}{8} + \tan^{-1} \frac{1}{15} = \cot^{-1} 3.$

2.  $\tan^{-1} \frac{4}{1} + \cot^{-1} \frac{2}{7} = \tan^{-1} \frac{1}{2}.$

3.  $\tan^{-1} 1 + \tan^{-1} 2 + \tan^{-1} 3 = \pi$   
 $= 2(\tan^{-1} 1 + \tan^{-1} \frac{1}{2} + \tan^{-1} \frac{1}{3}).$

4. (i)  $\tan^{-1} x + \cot^{-1} (x+1) = \tan^{-1} (x^2+x+1).$

(ii)  $\tan^{-1} \frac{1}{p+q} + \tan^{-1} \frac{q}{p^2+pq+1} = \tan^{-1} \frac{1}{p}.$

5.  $\tan^{-1} a - \tan^{-1} c = \tan^{-1} \frac{a-b}{1+ab} + \tan^{-1} \frac{b-c}{1+bc}.$

6.  $\tan^{-1} \frac{2}{3} + \sin^{-1} \frac{3}{5} = \tan^{-1} \frac{27}{11}.$

7.  $\tan^{-1} \frac{1}{3} + \tan^{-1} \frac{1}{2} + \tan^{-1} \frac{1}{7} + \tan^{-1} \frac{1}{8} = \frac{1}{2}\pi.$

[ C. U. 1942. ]

8.  $2 \tan^{-1} \frac{1}{3} + \tan^{-1} \frac{1}{7} = \frac{1}{2}\pi.$

[ C. U. 1937. ]

$$9. (i) \sin(2 \sin^{-1} x) = 2x \sqrt{1-x^2}.$$

$$(ii) \{\cos(\sin^{-1} x)\}^2 = \{\sin(\cos^{-1} x)\}^2.$$

$$10. \cos^{-1} x = 2 \sin^{-1} \sqrt{\frac{1-x}{2}} = 2 \cos^{-1} \sqrt{\frac{1+x}{2}}.$$

$$11. \tan^{-1} \sqrt{x} = \frac{1}{2} \cos^{-1} \frac{1-x}{1+x}. \quad [C. U. 1943.]$$

$$12. \sin^{-1} \sqrt{\frac{x-b}{a-b}} = \cos^{-1} \sqrt{\frac{a-x}{a-b}} = \tan^{-1} \sqrt{\frac{x-b}{a-x}}.$$

$$13. \tan^{-1} \frac{a-b}{1+ab} + \tan^{-1} \frac{b-c}{1+bc} + \tan^{-1} \frac{c-a}{1+ca} \\ = \tan^{-1} \frac{a^2-b^2}{1+a^2b^2} + \tan^{-1} \frac{b^2-c^2}{1+b^2c^2} + \tan^{-1} \frac{c^2-a^2}{1+c^2a^2}.$$

$$14. \sec^2(\tan^{-1} 2) + \operatorname{cosec}^2(\cot^{-1} 3) = 15.$$

$$15. \cot^{-1}(\tan 2x) + \cot^{-1}(-\tan 3x) = x.$$

$$16. \sin^{-1} \frac{4}{5} + \sin^{-1} \frac{5}{13} + \sin^{-1} \frac{16}{65} = \frac{1}{2}\pi. \quad [C. U. 1941.]$$

$$17. 4(\cot^{-1} 3 + \operatorname{cosec}^{-1} \sqrt{5}) = \pi. \quad [C. U. 1939.]$$

$$18. \text{If } \tan^{-1} x + \tan^{-1} y + \tan^{-1} z = \pi, \text{ show that} \\ x + y + z = xyz.$$

$$19. \text{If } \tan^{-1} x + \tan^{-1} y + \tan^{-1} z = \frac{1}{2}\pi, \text{ show that} \\ yz + zx + xy = 1.$$

$$20. \text{If } \cos^{-1} x + \cos^{-1} y + \cos^{-1} z = \pi, \text{ show that} \\ x^2 + y^2 + z^2 + 2xyz = 1.$$

$$21. \text{If } \sin^{-1} x + \sin^{-1} y + \sin^{-1} z = \pi, \text{ show that} \\ x \sqrt{1-x^2} + y \sqrt{1-y^2} + z \sqrt{1-z^2} = 2xyz.$$

$$22. \text{Find the values of}$$

$$(i) \sin(\sin^{-1} \frac{1}{2} + \cos^{-1} \frac{1}{2}). \quad [C. U. 1935.]$$

$$(ii) \tan(\tan^{-1} a + \cot^{-1} a).$$

$$(iii) \tan\left(\frac{1}{2} \sin^{-1} \frac{2x}{1+x^2} + \frac{1}{2} \cos^{-1} \frac{1-y^2}{1+y^2}\right).$$

23. If  $\tan^{-1}y = 4 \tan^{-1}x$ , find  $y$  as an algebraic function of  $x$ .

24. If  $\tan^{-1}x, \tan^{-1}y, \tan^{-1}z$  are in A.P., find out the algebraic relation between  $x, y, z$ . If in addition,  $x, y, z$  are also in A.P., prove that  $x=y=z$ . [ $y \neq 0, 1$ , or  $-1$ ]

25. Solve the following equations :

(i)  $\tan^{-1}(x+1) + \tan^{-1}(x-1) = \tan^{-1} \frac{8}{3}$ .

(ii)  $\tan^{-1} \frac{2x}{1-x^2} = \sin^{-1} \frac{2a}{1+a^2} - \cos^{-1} \frac{1-b^2}{1+b^2}$ .

(iii)  $\tan(\cos^{-1}x) = \sin(\tan^{-1}2)$ .

(iv)  $\tan^{-1} \frac{1-x}{1+x} = \frac{1}{2} \tan^{-1}x$ .

(v)  $\tan^{-1} \frac{x-1}{x+1} + \tan^{-1} \frac{2x-1}{2x+1} = \tan^{-1} \frac{23}{36}$ .

(vi)  $\sin^{-1}x + \sin^{-1}2x = \frac{\pi}{3}$ .

(vii)  $\sin^{-1}x + \sin^{-1}(1-x) = \cos^{-1}x$

(viii)  $\tan^{-1}(x-1) + \tan^{-1}x + \tan^{-1}(x+1) = \tan^{-1}3x$ .

(ix)  $\tan^{-1} \frac{2x}{1-x^2} + \cot^{-1} \frac{1-x^2}{2x} = \frac{\pi}{3}$ .

(x)  $\cot^{-1}(x-1) + \cot^{-1}(x-2) + \cot^{-1}(x-3) = 0$ .

26. Show that

(i)  $\cot^{-1} \frac{xy+1}{x-y} + \cot^{-1} \frac{yz+1}{y-z} + \cot^{-1} \frac{zx+1}{z-x} = 0$ .

(ii)  $\tan(\tan^{-1}x + \tan^{-1}y + \tan^{-1}z)$   
 $= \cot(\cot^{-1}x + \cot^{-1}y + \cot^{-1}z)$ .

(iii)  $\tan^{-1}(\cot x) + \cot^{-1}(\tan x) = \pi - 2x$ .



### Miscellaneous Examples I

1. If  $3 \sin \theta + 4 \cos \theta = 5$ , show that  $\tan \theta = \frac{3}{4}$ .
2. If  $a^2 \sec^2 x - b^2 \tan^2 x = c^2$ , find  $\operatorname{cosec} x$ .
3. If  $x = r \cos \theta \cos \phi$ ,  $y = r \cos \theta \sin \phi$ ,  $z = r \sin \theta$ , show that  $x^2 + y^2 + z^2 = r^2$ .

4. If  $\sin \theta = \frac{x-y}{x+y}$ , show that  $\tan \left( \frac{\pi}{4} - \frac{\theta}{2} \right) = \pm \sqrt{\frac{y}{x}}$ .

5. If  $x = r \sin (\theta + 45^\circ)$  and  $y = r \sin (\theta - 45^\circ)$ , then  $x^2 + y^2 = r^2$ .

6. If  $\cos (a + \beta) \sin (\gamma + \beta) = \cos (a - \beta) \sin (\gamma - \theta)$ , then  $\tan \theta = \tan a \tan \beta \tan \gamma$ .

Show that (Ex. 7 to 9) :—

7.  $(\cos x - \cos y)^2 + (\sin x - \sin y)^2 = 4 \sin^2 \frac{x-y}{2}$ .

8.  $\sin A + \sin B + \sin C - \sin (A + B + C)$   
 $= 4 \sin \frac{A+B}{2} \sin \frac{B+C}{2} \sin \frac{C+A}{2}$ .

9.  $4 \sin \frac{A+B+C}{2} \sin \frac{B+C-A}{2} \sin \frac{C+A-B}{2} \sin \frac{A+B-C}{2}$   
 $= 1 - \cos^2 A - \cos^2 B - \cos^2 C + 2 \cos A \cos B \cos C$ .

10. If  $\tan \beta = \frac{2 \sin \alpha \sin \gamma}{\sin (a + \gamma)}$ , then  $\tan a$ ,  $\tan \beta$ ,  $\tan \gamma$  are in harmonical progression.

11. If  $a + \beta + \gamma = (2n + 1) \frac{\pi}{2}$ , then

(i)  $\tan \beta \tan \gamma + \tan \gamma \tan a + \tan a \tan \beta = 1$ .

(ii)  $\sin 2a + \sin 2\beta + \sin 2\gamma = \pm 4 \cos a \cos \beta \cos \gamma$ .

12. If the angles  $A, B, C$  be in A. P., then

$$\frac{\sin A - \sin C}{\cos C - \cos A} = \frac{\cos B}{\sin B}$$

13. If  $\operatorname{cosec} 2A + \operatorname{cosec} 2B + \operatorname{cosec} 2C = 0$ , show that  
 $\tan A + \tan B + \tan C + \cot A + \cot B + \cot C = 0$ .
14. If  $\tan \alpha = \frac{a \sin \beta}{1 - a \cos \beta}$  and  $\tan \beta = \frac{b \sin \alpha}{1 - b \cos \alpha}$ ,  
 then  $\frac{\sin \alpha}{\sin \beta} = \frac{a}{b}$ .
15. Show that  
 $\tan \theta + 2 \tan 3\theta + 4 \tan 4\theta + 8 \cot 8\theta = \cot \theta$ .
16. If  $\cos(\theta - \psi) \cos \phi = \cos(\theta - \phi + \psi)$ , then  $\tan \theta$ ,  $\tan \phi$ ,  
 $\tan \psi$  are in harmonical progression.
17. If  $1 + \cos(y - z) \cos(z - x) \cos(x - y) = 0$ , show that  
 either  $(y - z)$ , or  $(z - x)$ , or  $(x - y)$  is an odd multiple of  $\pi$ .
18. If  $\sin \theta + \sin \phi = \sqrt{3}(\cos \phi - \cos \theta)$ , show that  
 $\sin 3\theta + \sin 3\phi = 0$ .
19. Eliminate  $\alpha$  and  $\beta$  from  
 $\sin \alpha + \sin \beta = a$ ,  $\cos \alpha + \cos \beta = b$ ,  $\cos(\alpha - \beta) = c$ .
20. If  $A + C + C = \pi$ , prove that  
 (i)  $\tan B \tan C + \tan C \tan A + \tan A \tan B$   
 $= 1 + \sec A \sec B \sec C$ .  
 (ii)  $\cot A + \cot B + \cot C = \cot A \cot B \cot C$   
 $+ \operatorname{cosec} A \operatorname{cosec} B \operatorname{cosec} C$ .
21. If  $A + B + C = \pi$ , and if  
 $\sin^2 A + \sin^2 B + \sin^2 C = \sin B \sin C + \sin C \sin A$   
 $+ \sin A \sin B$ , then  $A = B = C$ .
22. If  $A, B, C$  be the angles of a triangle, and if  
 $\cot A + \cot B + \cot C = \sqrt{3}$ , show that the triangle  
 is equilateral.
23. If  $\sec ax + \sec bx = 0$ , show that the values of  $x$   
 form two series in A. P.

## CHAPTER XIII

### LOGARITHMS

#### 72. Definition of Logarithm.

Logarithm of a number with respect to a given base is the index of the power to which the base is to be raised in order to give the number.

Mathematically if  $a^x = N$ , then ' $x$ ' is the power to which ' $a$ ' (which is called the base) is raised to give ' $N$ '. Hence by definition, ' $x$ ' is the logarithm of ' $N$ ' with respect to the base ' $a$ ', and it is usually written as  $x = \log_a N$ .

As a numerical example,  $\log_2 8 = 3$ , for  $2^3 = 8$  i.e. 3 is the power to which 2 is to be raised to give 8. Again, since  $3^4 = 81$ ,  $4 = \log_3 81$ .

Any result involving indices can be expressed as a result in logarithm, and *vice versa*.

For example,

$$\text{if } p^q = r, \text{ then, } q = \log_p r.$$

$$\text{If } m^n = z^k, \text{ then } n = \log_m (z^k)$$

$$\text{or } k = \log_z (m^n).$$

Similarly, if  $\log_y x = z$ ,

$$\text{then } y^z = x.$$

It should be noted that the logarithm of the same number with respect to different bases will be different ; for example, to get the same number 64, we must raise 2 to the power 6, whereas we are to raise 4 to the power 3 and 8 to the power 2 only ; hence  $\log_2 64 = 6$ ,  $\log_4 64 = 3$ ,  $\log_8 64 = 2$ .

Thus so long as the base is not stated, logarithm of a number has no meaning.

### 73. Special Cases.

We know from Algebra that if  $a$  be any real finite quantity, other than zero, then  $a^0 = 1$ .

Hence,  $\log_a 1 = 0$ ; in other words,

(i) *logarithm of 1 with respect to any finite quantity (other than zero) as base, is zero.*

Again,  $a$  being any quantity,  $a^1 = a$ .

Hence,  $1 = \log_a a$ . In other words,

(ii) *logarithm of any number with respect to itself as base is unity.*

**Note 1.** If  $a^x = 0$ , then  $x = -\infty$  if  $a > 1$ , and  $x = +\infty$  if  $a < 1$ .

Thus, we have  $\log_a 0 = \mp \infty$  according as  $a >$  or  $< 1$ . Hence, *logarithm of zero to a base greater than unity is minus infinity, and to a base less than unity is plus infinity.*

**Note 2.** Since the equation  $a^x = -n$  ( $a$  and  $n$  being real positive quantities), cannot be satisfied by any real value of  $x$ , whether positive or negative, provided we consider the principal value\* only of  $a^x$ ,

therefore, *logarithm of a negative quantity (in a system of logarithms whose base is a real positive quantity) must be imaginary.*

### 74. Fundamental formulæ in logarithms.

From the definition it is clear that logarithms are but indices in another form. Hence corresponding to the three fundamental results in the theory of indices in Algebra, namely that if  $a, x, y$  be any real quantities,

$$(i) a^x \times a^y = a^{x+y},$$

$$(ii) a^x \div a^y = a^{x-y} \text{ and}$$

$$(iii) (a^x)^y = a^{xy},$$

we get three fundamental laws of logarithms which are given below.

---

\*See a treatise on Higher Trigonometry.

$$(i) \log_a (m \times n) = \log_a m + \log_a n$$

in other words, *logarithm of the product of two quantities is equal to the sum of their logarithms taken separately, base remaining the same always.*

$$\text{Proof. Put } \log_a m = x, \log_a n = y$$

$$\text{and } \log_a (m \times n) = z$$

then from definition,

$$a^x = m, a^y = n \text{ and } a^z = m \times n = a^x \times a^y = a^{x+y},$$

so that,  $z = x + y$ .

Replacing values,

$$\log_a (mn) = \log_a m + \log_a n.$$

$$\text{Cor. } \log_a (m.n.p.\dots) = \log_a m + \log_a n + \log_a p + \dots$$

$$(ii) \log_a \left( \frac{m}{n} \right) = \log_a m - \log_a n$$

in other words, *logarithm of the quotient of two numbers is equal to the difference of their logarithms (logarithm of the numerator minus logarithm of the denominator).*

$$\text{Proof. Put } \log_a m = x, \log_a n = y$$

$$\text{and } \log_a \left( \frac{m}{n} \right) = z.$$

Then from definition,

$$a^x = m, a^y = n$$

$$\text{and } a^z = \frac{m}{n} = \frac{a^x}{a^y} = a^{x-y}$$

so that

$$z = x - y,$$

or replacing values,

$$\log_a \left( \frac{m}{n} \right) = \log_a m - \log_a n.$$

(iii)  $\log_a (m)^n = n \log_a m$ .

Or, *logarithm of a power of a number is the product of the power and the logarithm of the number.*

**Proof.** Put  $\log_a m = x$ , and  $\log_a (m)^n = z$ .

Then by definition,

$$a^x = m \text{ and}$$

$$a^z = (m)^n = (a^x)^n = a^{nx}.$$

$$\therefore z = nx,$$

or replacing values,

$$\log_a (m)^n = n \log_a m.$$

**Ex. 1.** Reduce to a simple form  $\log_a \frac{x^p y^q}{z^s}$ .

$$\begin{aligned} \log_a \frac{x^p y^q}{z^s} &= \log_a (x^p y^q) - \log_a (z^s) \\ &= \log_a x^p + \log_a y^q - \log_a z^s \\ &= p \log_a x + q \log_a y - s \log_a z. \end{aligned}$$

**Ex. 2.** Simplify  $\log_{10} \sqrt[3]{\frac{25}{88}}$ .

$$\begin{aligned} \log_{10} \sqrt[3]{\frac{25}{88}} &= \log_{10} \left( \frac{5^2}{8 \cdot 11} \right)^{\frac{1}{3}} = \frac{1}{3} \log_{10} \frac{5^2}{2^3 \cdot 11} \\ &= \frac{1}{3} \log_{10} \frac{10^2}{2^3 \cdot 11} \\ &= \frac{1}{3} [\log_{10} 10^2 - \log_{10} (2^3 \cdot 11)] \\ &= \frac{1}{3} [2 \log_{10} 10 - (\log_{10} 2^3 + \log_{10} 11)] \\ &= \frac{1}{3} [2 - 5 \log_{10} 2 - \log_{10} 11]. \end{aligned}$$

## 75. Change of base.

There is a fourth standard formula whereby logarithms of numbers with respect to one base being given, those with respect to a different base may be obtained. The formula is

$$\log_a m = \log_b m \times \log_a b$$

**Proof.** Put  $\log_a m = x$ ,  $\log_b m = y$  and  $\log_a b = z$ .

Then from definition,

$$a^x = m, \quad b^y = m, \quad a^z = b.$$

Hence, 
$$a^x = m = b^y = (a^z)^y = a^{yz},$$

or, 
$$x = yz.$$

Replacing values,

$$\log_a m = \log_b m \times \log_a b.$$

**Cor. 1.** In the above result, put  $m = a$ . Then remembering that  $\log_a a = 1$ , we get

$$\log_b a \times \log_a b = 1.$$

Since the above relation is very important, we add here an *independent proof* of it.

Let  $\log_b a = x$ , and  $\log_a b = y$ .

Then  $b^x = a$  and  $a^y = b$ .

$$\therefore a = b^x = (a^y)^x = a^{xy}. \quad \therefore xy = 1,$$

i.e.  $\log_b a \times \log_a b = 1$ ,

or, 
$$\log_b a = \frac{1}{\log_a b}.$$

**Cor. 2.** The result of the above article may be written with the help of Cor. 1, in the form

$$\log_a m = \log_b m / \log_b a.$$

Thus if logarithms of both  $m$  and  $a$  with respect to  $b$  be known, logarithm of  $m$  with respect to  $a$  is obtained.

## 76. Common system of logarithms.

For all practical purposes, wherever logarithms are used for numerical calculations, the base is invariably taken as 10. Logarithms of numbers with respect to the base 10 are

referred to as the *Common system* of logarithms. The advantage of the common system of logarithms for practical applications will be clear presently, from the article 77, Theorems I & II.

**Note.** In higher mathematics, for *theoretical* investigations, another quantity '*e*' (defined in books of Algebra), whose value is nearly 2.718..., is used as the base of logarithms, and logarithms to this base *e* are called *Napierian* logarithms.

With the help of the logarithmic series established in books on Algebra, Napierian logarithms of numbers are tabulated. The factor  $\frac{1}{\log_{e}10}$  which is known as the *modulus of the common system*, applied to the Napierian logarithms will convert them to common logarithms (*See Art. 75*). Thus a table of common logarithms is prepared.

Henceforth we shall proceed with the consideration of the common system of logarithms, and the base being understood to be 10, will not be written.

### 77. Characteristic and Mantissa of common logarithms.

It is only in very few cases that the logarithm of a number is integral. In most cases, however, the logarithm of a number is partly integral and partly fractional (or decimal).

**Def.** The integral portion of the logarithm of a number is called the *characteristic*, and the decimal portion is called the *mantissa*.

In case the logarithm of a number is negative, and partly integral and partly decimal, the decimal portion, *i.e.*, the mantissa is always kept positive by altering the integral part *i.e.*, the characteristic suitably. Thus the *mantissa part of the logarithm of a number is always positive*. For instance, if the logarithm of a number is  $-2.3$ , we write it as  $-3 + .7$  and call  $-3$  as the characteristic and  $.7$  (and not  $-.3$ ) as the mantissa.  $-3 + .7$  is often abbreviated in the form  $\bar{3}.7$ .



**Theorem I.** *The characteristic of the common logarithm of (i) any number greater than 1 is positive, and numerically one less than the number of digits in the integral part of the quantity whose logarithm is sought; and (ii) of any positive\* number less than 1, is negative, and numerically one greater than the number of zeroes immediately after the decimal point in the quantity whose logarithm is wanted.*

(i) Let the number be greater than unity.

Any number, say 7'209, which consists of 1 digit only in its integral part, lies between 1 and 10.

Now  $10^0 = 1$  and  $10^1 = 10$ .

Hence if  $10^x = 7'209$ , clearly  $x$  must be greater than 0 and less than 1.

Thus  $\log 7'209$ , must lie between 0 and 1, i.e., of the form 0'..., having its characteristic 0.

Similarly, numbers of the type 53'0528, which consists of 2 digits in their integral parts, must lie between 10 and 100, i.e., between  $10^1$  and  $10^2$ .

Hence the index to which 10 should be raised to give 53'0528 must be greater than 1 and less than 2, i.e.,  $\log 53'0528$  must be of the form 1'... having the characteristic 1.

$\log 10$  is 1, and 10 also falls in this category of two digits.

In the same way, a number which has  $n$  digits in its integral part lies between  $10^{n-1}$  (which also has  $n$  digits) and  $10^n$  (which has  $n+1$  digits). Thus the logarithm of such numbers must lie between  $n-1$  and  $n$ , i.e.  $(n-1)$  + some positive proper fraction. Hence the characteristic in such cases is  $n-1$ .

Hence the result.

\*Logarithms of negative numbers are easily seen to be imaginary, for there is no real power, positive, or negative, to which 10 may be raised to give a negative result. [ See Note 2, Art. 73. ]

(ii) Let the number be positive, and less than 1 (*i.e.* between 0 and 1).

We notice that

$$\begin{array}{ll} 10^0 & = 1 \\ 10^{-1} = \frac{1}{10} & = \cdot 1 \\ 10^{-2} = \frac{1}{100} & = \cdot 01 \\ 10^{-3} = \frac{1}{1000} & = \cdot 001 \\ 10^{-4} = \frac{1}{10000} & = \cdot 0001 \\ \text{etc.} & \text{etc.} \quad \text{etc.} \end{array}$$

Now a number less than 1, with no zero immediately after the decimal point, like '3015, must be greater than '1 and less than 1; hence the power to which 10 must be raised to give such a number must lie between -1 and 0, *i.e.*, = -1 + a positive proper fraction. Hence such numbers have the characteristic of their logarithms = -1.

A decimal number with one zero immediately after the decimal point, like '078005, lies between '01 and '1 which are respectively equal to  $10^{-2}$  and  $10^{-1}$ .

Hence if  $10^x = \cdot 078005$ ,  $x$  must lie between -1 and -2 *i.e.*,  $x$  is of the form  $-1 \cdot \dots\dots$ . Writing the decimal part of  $x$  positively, in the form  $-2 + \dots\dots$ , we notice that the integral part of  $x$  *i.e.*, the characteristic of the logarithm of '078005 is -2.

Similarly the logarithms of numbers between '01 and '001 (*i.e.*,  $10^{-2}$  and  $10^{-3}$ ) which must have two zeroes after the decimal point, lie between -2 and -3 *i.e.*, are of the form  $-2 \cdot \dots\dots = -3 + \dots\dots$ , and so the characteristic in such cases is -3,

and so on.

Hence the result.

**Theorem II.** *All numbers, formed of the same digits in the same order, differing only in the positions of their decimal points, have the mantissæ of their logarithms same.*

This will be clear from an example. Let us take the numbers 835107, 835107000, 83'5107, '835107, '000835107 and 8351'07.

$$\begin{aligned}\text{Now } \log 835107000 &= \log (835107 \times 1000) \\ &= \log 835107 + \log 1000 \\ &= \log 835107 + 3.\end{aligned}$$

$$\begin{aligned}\text{Again, } \log 83'5107 &= \log \frac{835107}{10000} \\ &= \log 835107 - \log 10000 \\ &= \log 835107 - 4.\end{aligned}$$

$$\log '835107 = \log \frac{835107}{1000000} = \log 835107 - 6.$$

$$\log '000835107 = \log \frac{835107}{10^9} = \log 835107 - 9.$$

$$\log 8351'07 = \log \frac{835107}{100} = \log 835107 - 2.$$

Thus the logarithms of all the numbers here differ from the logarithm of 835107 by a whole number in each case and so must have their decimal parts *i.e.*, their mantissæ the same as that of  $\log 835107$ .

In fact, numbers formed of the same digits in the same order differing only in the position of their decimal points, must have their ratios equal to an integral power of 10 and so must have their logarithms differing only by a whole number.

Hence the result.

The two theorems above given show that (i) the characteristic of the logarithm of a number can be found by a simple glance at the number and (ii) that for the mantissa part of the logarithm of a number, we need only take into

account the digits of which the number is formed, without taking any notice of the position of the decimal point in it.

In logarithmic tables, only the mantissæ of the logarithms of numbers are therefore given.

These constitute the special advantages of the common system of logarithms.

### 78. Examples worked out.

**Ex. 1.** *Simplify*

$\log \frac{5^{\frac{1}{2}} \cdot 2^{\frac{1}{10}}}{\sqrt[3]{18 \cdot \sqrt{2}}}$ , and find its value, given

$\log 2 = \cdot 30103$  and  $\log 3 = \cdot 4771213$ .

The given exp. =  $\log \frac{5^{\frac{1}{2}} \cdot 2^{\frac{1}{10}}}{(18 \cdot 2^{\frac{1}{2}})^{\frac{1}{3}}}$

$$\begin{aligned} &= \log \frac{10^{\frac{1}{2}} \cdot 2^{\frac{1}{10}}}{2^{\frac{1}{3}} (2 \cdot 3^2 \cdot 2^{\frac{1}{2}})^{\frac{1}{3}}} = \log \frac{10^{\frac{1}{2}} \cdot 2^{\frac{1}{10}}}{2^{\frac{1}{3}} \cdot 2^{\frac{1}{3}} \cdot 3^{\frac{2}{3}} \cdot 2^{\frac{1}{6}}} \\ &= \log \frac{10^{\frac{1}{2}}}{2^{\frac{1}{3} + \frac{1}{3} + \frac{1}{6}} \cdot 3^{\frac{2}{3}}} = \log 10^{\frac{1}{2}} - \log (2^{\frac{1}{3} + \frac{1}{3} + \frac{1}{6}} \times 3^{\frac{2}{3}}) \\ &= \frac{1}{2} \log 10 - (\log 2^{\frac{1}{3} + \frac{1}{3} + \frac{1}{6}} + \log 3^{\frac{2}{3}}) \\ &= \frac{1}{2} \log 10 - \frac{1}{2} \log 2 - \frac{2}{3} \log 3 \end{aligned}$$

and its value is

$$\begin{aligned} &\frac{1}{2} \cdot 1 - \frac{1}{2} \cdot 3( \cdot 30103 ) - \frac{2}{3} ( \cdot 4771213 ) \\ &= \cdot 25 - \cdot 1956095 - \cdot 3180809 \\ &= -1 + \cdot 7362496 \\ &= \bar{1} \cdot 7362496. \end{aligned}$$

**Note.**  $\log 5 = \log 10 - \log 2 = 1 - \log 2$  and hence  $\log 5$  is deducible from  $\log 2$ .

**Ex. 2.** *Prove that*

$$7 \log \frac{10}{9} - 2 \log \frac{25}{12} + 3 \log \frac{81}{50} = \log 2.$$

The left-hand expression

$$\begin{aligned}
 &= \log \left(\frac{10}{9}\right)^7 - \log \left(\frac{25}{24}\right)^2 + \log \left(\frac{81}{80}\right)^3 \\
 &= \log \frac{\left(\frac{10}{9}\right)^7 \times \left(\frac{81}{80}\right)^3}{\left(\frac{25}{24}\right)^2} \\
 &= \log \left\{ \left(\frac{10}{9}\right)^7 \times \left(\frac{3^4}{10 \times 2^3}\right)^2 \times \left(\frac{3 \times 2^3 \times 2^3}{10^2}\right)^3 \right\} \\
 &= \log \left( \frac{10^7}{3^{14}} \times \frac{3^{12}}{10^5 \times 2^6} \times \frac{3^9 \times 2^{10}}{10^4} \right) \\
 &= \log 2.
 \end{aligned}$$

*Alternative method :*

Left side

$$\begin{aligned}
 &= 7(\log 10 - \log 9) - 2(\log 25 - \log 24) + 3(\log 81 - \log 80) \\
 &= 7\{\log (5 \times 2) - \log 3^2\} - 2\{\log 5^2 - \log (3 \times 2^3)\} \\
 &\quad + 3\{\log 3^4 - \log (5 \times 2^4)\} \\
 &= 7\{\log 5 + \log 2 - 2 \log 3\} - 2\{2 \log 5 - \log 3 - 3 \log 2\} \\
 &\quad + 3\{4 \log 3 - \log 5 - 4 \log 2\} \\
 &= \log 2.
 \end{aligned}$$

**Ex. 3.** Find the number of digits in  $4^{15}$ , having given  $\log 2 = .30103$ .

We have

$$\begin{aligned}
 \log 4^{15} &= \log 2^{30} = 30 \log 2 \\
 &= 30 \times .30103 = 9.0309.
 \end{aligned}$$

Hence since the characteristic of  $\log 4^{15}$  is 9,  $4^{15}$  must consist of 10 digits.

**Ex. 4.** Find approximately the 7<sup>th</sup> root of 35.28, having given  $\log 2 = .30103$ ,  $\log 3 = .4771213$ ,  $\log 7 = .8450980$  and  $\log 1197342 = 3.0782184$ .

$$\text{Let } x = (35.28)^{\frac{1}{7}} = \left( \frac{7^2 \times 3^2 \times 2^3}{10^2} \right)^{\frac{1}{7}}$$

$$\begin{aligned}
 \text{then } \log x &= \frac{1}{7}[2 \log 7 + 2 \log 3 + 3 \log 2 - 2 \log 10] \\
 &= \frac{1}{7}[2 \times .8450980 + 2 \times .4771213 + 3 \times .30103 - 2] \\
 &= .0782184 \text{ nearly.}
 \end{aligned}$$

Now  $\log 1197'342 = 3'0782184$ .

$\therefore \log 1'197342 = '0782184$ , having characteristic 0, but mantissa same as that of  $\log 1197'342$ .

Hence  $x = 1'197342$  approximately.

**Ex. 5.** Obtain an approximate numerical solution of  $2^x \cdot 3^{2x} = 100$ , having given  $\log 2 = '30103$ ,  $\log 3 = '47712$ .

We have

$$2^x \cdot 3^{2x} = 10^2.$$

$$\therefore \log (2^x \cdot 3^{2x}) = \log 10^2,$$

$$\text{i.e., } x \log 2 + 2x \log 3 = 2 \log 10 = 2.$$

$$\therefore x = \frac{2}{\log 2 + 2 \log 3} = \frac{2}{'30103 + 2 \times '47712} \\ = 1'5933 \text{ nearly.}$$

**Note.** Equations of this type are called **Exponential equations**.

### Examples XIII (a)

[ Use the values :  $\log 2 = '30103$ ,  $\log 3 = '4771213$ ,  
 $\log 7 = '8450980$  when required ]

1. Find the logarithm of (i) 1728 to the base  $2\sqrt{3}$   
 (ii)  $\cos^3 a$  to the base  $\sec a$ .

2. Find  $\log_{.01} 10000$ .

3. Show that  $\log_{10} 2$  lies between  $\frac{1}{3}$  and  $\frac{1}{2}$

[ C. U. I. 1926. ]

4. Prove that

$$(i) \log_a m \times \log_b n = \log_b m \times \log_a n.$$

$$(ii) \log_2 \log_2 \log_2 16 = 1.$$

(5.) If  $\log_e m + \log_e n = \log_e (m+n)$ , find  $m$  as a simple function of  $n$ .

6. Prove that if a series of numbers be in G.P., their logarithms are in A.P.

7. Prove that

$$2 \log a + 2 \log a^2 + 2 \log a^3 + \dots + 2 \log a^n = n(n+1) \log a.$$

8. If  $x$  is positive and less than unity, show that  
 $\log(1+x) + \log(1+x^2) + \log(1+x^4) + \log(1+x^8) + \dots$  to  $\infty$   
 $= -\log(1-x).$

9. Simplify

(i)  $\log_2 \sqrt{6} + \log_2 \sqrt{\frac{2}{3}}.$

(ii)  $\frac{\log \sqrt{27} + \log 8 - \log \sqrt{1000}}{\log 12}.$

10. Find  $\log(00225)^{\frac{1}{3}}$  and  $\log(7\frac{5}{2})^{-\frac{1}{8}}.$

11. Prove that

(i)  $\log_a b \times \log_b c \times \log_c a = 1.$

(ii)  $\log_a x = \log_b x \times \log_c b \times \log_d c \dots \times \log_n m \times \log_a n.$

12. Show that

(i)  $7 \log \frac{1}{16} + 5 \log \frac{2}{24} + 3 \log \frac{8}{36} = \log 2.$

(ii)  $7 \log \frac{1}{18} + 6 \log \frac{8}{3} + 5 \log \frac{2}{5} + \log \frac{3}{2} = \log 3.$

13. Extract the fifth root of 84 having given

$$\log 2425805 = 6.3848559.$$

14. Calculate  $(0020736)^{\frac{1}{7}}$ , having given

$$\log 41369 = 4.6166750.$$

15. Simplify

(i)  $\log \sqrt[7]{\frac{8^{\frac{1}{3}} \times 14^{\frac{1}{3}}}{\sqrt{72} \times \sqrt[5]{60}}}.$

(ii)  $\sqrt[3]{7.2 \times 6.3}$ , having given

$$\log 898665 = 5.9535977.$$

16. Find the value of  $64\{1 - (1.05)^{-20}\}$ , having given  $\log 24121 = 4.382394$ .

17. Find the number of digits in (i)  $2^{40}$ , (ii)  $3^{11}$ , (iii)  $(540)^9$ .

18. Find the number of zeroes after the decimal point before the first significant digit in the expressions

$$(i) (.024)^{15}, \quad (ii) \left(\frac{1}{1.05}\right)^4, \quad (iii) (.0259)^{50}.$$

19. Solve the equations

$$(i) 3^x = 2, \quad (ii) 3^{x-4} = 7.$$

$$(iii) 5^{6x} + 7^{x+2} = 3^{2x-3}.$$

$$(iv) \left. \begin{aligned} 2^x &= 3^y \\ 2^{y+1} &= 3^{x-1} \end{aligned} \right\} \quad (v) \left. \begin{aligned} 7^{x+y} \times 3^{2x+y} &= 9 \\ 3^{x-y} + 2^{x-2y} &= 3^x \end{aligned} \right\}$$

20. (i) If  $\log(x^2 y^3) = a$ ,  $\log\left(\frac{x}{y}\right) = b$ , find  $\log x$  and  $\log y$ .

(ii) If  $a^2 + b^2 = 7ab$ , show that

$$\log\left\{\frac{1}{3}(a+b)\right\} = \frac{1}{2}(\log a + \log b).$$

21. If  $\frac{\log x}{y-z} = \frac{\log y}{z-x} = \frac{\log z}{x-y}$  show that  $x^x y^y z^z = 1$ .

22. Why is  $\log(1+2+3) \neq \log 1 + \log 2 + \log 3$ ?

23. If  $a, b, c, \dots$  be in G.P., show that

$$\log_a x, \log_b x, \log_c x \dots \dots \dots \text{are in H.P.}$$

24. If  $xy^{l-1} = a$ ,  $xy^{m-1} = b$ ,  $xy^{n-1} = c$ , prove that

$$(m-n) \log a + (n-l) \log b + (l-m) \log c = 0.$$

25. If  $\frac{x(y+z-x)}{\log x} = \frac{y(z+x-y)}{\log y} = \frac{z(x+y-z)}{\log z}$ , show that

$$y^x z^y = z^x x^z = x^y y^x.$$



## 79. Tables of Logarithms and Trigonometrical ratios.

Several mathematical tables correct up to five places of decimals are given at the end of the book. An explanation of the tables is given below.

Table I gives the common logarithm of all numbers from 1 to 10000, *i.e.*, those which consist of 4 digits or less. The tabulated quantities are the mantissæ only, correct to five places, with the decimal point dropped. The characteristic is to be supplied according to the rule given in Art. 77. The main body of the table gives logarithms (mantissa part) of numbers of 3 digits, and the mean difference table at the side supplies the increment in the mantissa due to the fourth digit. This increment is written, in order to save space, giving the significant digits only, which are to be supplied with the necessary number of zeroes to make up 5 places (here the table being a five figure table). Thus '00024 will be written as 24 only in the difference table. As an example, to find  $\log 2'697$ , we notice from the table that the mantissa for  $\log 269$  is '42975, and along the same row, the difference table gives 115 under the heading 7. This means that for 7 in the fourth place of the number (*i.e.*, for the number 2697) the increment in the mantissa will be '00115. Hence  $\log 2697$  will have its mantissa '42975 + '00115 = '43090. Again  $\log 2'697$  has the same mantissa, but its characteristic is 0. Thus  $\log 2'697 = 0'43090$ .

Table II gives ordinary sines and cosines (usually referred to as *natural sines and cosines*) of all angles from  $0^\circ$  to  $90^\circ$  at intervals of  $1'$ , sines being given from the left side of the top towards the right and downwards, and cosines being given from the right side of the bottom towards the left and upwards. The table is arranged in such a way that the sine of any angle given is the same as the cosine of exactly the complementary angle, and it is on this arrangement that a single table serves as a sine as well as a cosine table. The main portion of the table gives sines or cosines of angles at interval of  $10'$ , and the difference

table at the side gives changes in the value of the sine or cosine for changes in minutes in the angles. It should be remembered that as an angle increases from  $0^\circ$  to  $90^\circ$ , its sine increases from 0 to 1 whereas its cosine decreases from 1 to 0. Hence *the changes given in the difference table are to be added in case of sines and subtracted in case of cosines* for the increased number of minutes in the angle. Moreover, as in Table I, the numbers in the difference table are to be made up to five places of decimals by supplying the requisite number of zeroes before it. For example, using the table,  $\sin 53^\circ 23' = .80212 + .00052 = .80264$  and  $\cos 29^\circ 42' = .86892 - .00029 = .86863$ .

Table III similarly gives natural tangents and cotangents of angles from  $0^\circ$  to  $90^\circ$ , obtained at intervals of  $1'$  with the help of the difference table. The quantities in the difference table, being made up into five figures, are to be *added in case of tangents and subtracted in case of cotangents* for increased number of minutes in the angle.

Table IV gives logarithmic sines and logarithmic cosines of all angles from  $0^\circ$  to  $90^\circ$  at intervals of  $1'$  (with the aid of the difference table). Logarithmic sine of angle  $\theta$ , written as  $L \sin \theta$  means  $10 + \log \sin \theta$ , and similarly logarithmic cosine of  $\theta$ , written as  $L \cos \theta$  means  $10 + \log \cos \theta$ . In taking logarithms of trigonometrical ratios of angles, it may be noted that sines and cosines of angles are numerically less than unity, and tangents of angles between  $0^\circ$  and  $45^\circ$  as also cotangents of angles between  $45^\circ$  and  $90^\circ$  are less than unity. Hence logarithms of these quantities are negative. To avoid using negative values in the tables, *logarithms of trigonometrical ratios are always tabulated after adding 10 to them*. Thus the table gives  $L \sin \theta$  and  $L \cos \theta$  (and not  $\log \sin \theta$  and  $\log \cos \theta$ ).

Table V gives logarithmic tangents (*i.e.*  $L \tan \theta = 10 + \log \tan \theta$ ) and logarithmic cotangents (*i.e.*  $L \cot \theta = 10 + \log \cot \theta$ ) of all angles from  $0^\circ$  to  $90^\circ$ , obtained at intervals of  $1'$  with the aid of the difference table.

### 80. Principle of Proportional Parts.

Suppose we find from table I the logarithms of the two numbers 6257 and 6258, and we want to find the logarithm of 6257.6; or that we find from table III,  $\tan 53^\circ 23'$  and  $\tan 53^\circ 24'$ , but we want to find  $\tan 53^\circ 23' 20''$ ; or similarly, from table IV, we get  $L \cos 37^\circ 42'$  and  $L \cos 37^\circ 43'$  but we want to find  $L \cos 37^\circ 42' 48''$ ; how are we to proceed?

In order to meet such cases, the 'Principle of Proportional Parts' may be used. The principle may be stated as follows:

*If the value of a quantity depending on a variable quantity  $x$  be tabulated for different values of  $x$  at regular small intervals, then in most cases, for a very small change in  $x$  (which is called the argument) the corresponding small change in the tabulated quantity, (called the function of the argument) is proportional to the change in  $x$ .*

We shall assume the truth of this principle; for a strict proof of it, with the proper restrictions under which it is true, depends on the use of Calculus. For the tables with which we are concerned, it is true for all practical purposes.

The application of the principle is illustrated in the following examples:

**Ex. 1.** *Given  $\log 63374 = 4.8019111$  and  $\log 63375 = 4.8019180$ , find  $\log 63374.3$  and find the number whose logarithm is  $2.8019136$ .*

Here  $\log 63375 = 4.8019180$   
and  $\log 63374 = 4.8019111$

Hence for an increase of 1 in the number, the increment in the logarithm is '0000069. (This is usually spoken as 'diff. for 1 is 69')

Therefore by the Principle of Proportional Parts, increase in the logarithm for an increase of '3 in the number is

$$\begin{aligned} '3 \times '0000069 &= '00000207 \\ &= 0000021, \text{ up to seven places.} \end{aligned}$$

$$\begin{aligned} \text{Hence } \log 63374'3 &= 4'8019111 + '0000021 \\ &= 4'8019132. \end{aligned}$$

$$\therefore \log 63'3743 = 1'8019132.$$

Again, 4'8019136 lies between 4'8019111 and 4'8019180, the difference from the former being '0000025. Hence 4'8019136 is the logarithm of a number lying between 63374 and 63375, say logarithm of  $63374 + x$ .

Then diff. for 1 being 69 (i.e., '0000069) and diff. for  $x$  being 25, (i.e., '0000025), by the Principle of Proportional Parts, we have

$$\begin{aligned} 69 : 25 &:: 1 : x \\ \text{or, } x &= \frac{25}{69} = '36\cdots\cdots \end{aligned}$$

$$\text{Hence } \log 63374'36\cdots = 4'8019136.$$

The required number whose logarithm is  $\bar{2}'8019136$ , having the same mantissa, must be formed of the same digits arranged in the same order, and its characteristic being -2, the number must be 06337436...

$$\begin{aligned} \text{Ex. 2. (i) Given } L \sin 37^\circ 43' 50'' &= 9'7867152 \\ L \sin 37^\circ 44' &= 9'7867424, \\ \text{find } L \sin 37^\circ 43' 56''. \end{aligned}$$

$$\begin{aligned} \text{(ii) Given } L \tan 79^\circ 51' 40'' &= 10'7475657 \\ L \tan 79^\circ 51' 50'' &= 10'7476872, \\ \text{find the angle whose } L \tan &\text{ is } 10'7476532. \end{aligned}$$

[ C. U. 1921. ]

In (i) diff. ( in the value of  $L \sin$  ) for  $10''$  (diff. in angle)  
 $= 272$  (i.e., '0000272)

hence diff. for  $6'' = \frac{6}{10} \times 272 = 163.2$  i.e., '00001632

and so  $L \sin 37^\circ 43' 56'' = 9.7867152 + '0000163$   
 $= 9.7867315.$

In (ii) the angle whose  $L \tan$  is  $19.7476532$  evidently lies between  $79^\circ 51' 40''$  and  $79^\circ 51' 50''$ .

Let the angle be  $79^\circ 51' 40'' + x''$ .

Now diff. (in the value of  $L \tan$ ) for  $10''$  (diff. in angle )  
 $= 1215$  (i.e., '0001215 )

and diff. for  $x'' = 875$

(i.e., '0000875, being  $10.7476532 - 10.7475657$  )

$\therefore \frac{x}{10} = \frac{875}{1215}$  or  $x = 7.2$  nearly.

Thus the required angle is  $79^\circ 51' 47''.2$ .

**Ex. 3.** Given  $\cos 58^\circ 17' = .5257191$  and diff. for  $1' = 2474$ ,  
 find  $\cos 58^\circ 17' 20''$ .

Here diff. for  $1'$  i.e.,  $60'' = 2474$ .

$\therefore$  diff. for  $20'' = \frac{20}{60} \times 2474 = 825$  ( nearly ).

As for increasing angle, cosine diminishes,

$\therefore \cos 58^\circ 17' 20'' = .5257191 - '0000825$   
 $= .5256366.$

### Examples XIII(b)

- Given  $\log 18.906 = 1.2765997$   
 and  $\log 18.907 = 1.2766226$ ,  
 find  $\log 1890.635$ .

- Given  $\log 69714 = 4.8433200$   
 $\log 69715 = 4.8433262$ ,  
 find  $\log (.000697145)^{\frac{1}{3}}$ .

3. Given  $\log 37602 = 4.5752109$   
 $\log 37601 = 4.5751994$ ,  
 find the number whose logarithm is  $1.5752086$ .
4. Given  $\log 3 = .4771213$   
 $\log 74008 = 4.8692787$   
 diff. for  $1' = 59$ ,  
 find  $(.09)^{\frac{1}{8}}$ .
5. Given  $\cos 32^\circ 16' = .8455726$   
 and  $\cos 32^\circ 17' = .8454172$ ,  
 find the value of  $\cos 32^\circ 16' 24''$   
 and find the angle whose cosine is  $.8455176$ .
6. Find  $\tan 38^\circ 24' 37.5''$  having given  
 $\tan 38^\circ 24' = .7925902$  and  $\tan 38^\circ 25' = .7930640$ .
7. Given  $L \sin 44^\circ 17' = 9.8439842$   
 and  $L \sin 44^\circ 18' = 9.8441137$ ,  
 find  $L \sin 44^\circ 17' 33''$ . Deduce the value of  
 $L \operatorname{cosec} 44^\circ 17' 33''$ .
8. Given  $L \sin 36^\circ 24' = 9.7733614$   
 $L \sin 36^\circ 25' = 9.7735327$ ,  
 find the angle whose  $L \sin$  is  $9.7734642$ .
9. If  $L \cot 53^\circ 13' = 9.8736937$   
 $L \cot 53^\circ 14' = 9.8734302$ ,  
 find  $\theta$  where  $L \cot \theta = 9.8734523$ .
10. Given  $L \tan 22^\circ 37' = 9.6197205$   
 diff. for  $1' = 3557$ ,  
 find the value of  
 $L \tan 22^\circ 37' 22''$   
 and the angle whose  $L \tan$  is  $9.6195283$ .

11. Prove that.  $\theta$  being any acute angle,

$$\begin{aligned} L \sin \theta + L \operatorname{cosec} \theta &= L \cos \theta + L \sec \theta \\ &= L \tan \theta + L \cot \theta = 20. \end{aligned}$$

12. Given  $L \cos 36^\circ 40' = 9.9042411$ , find  $L \sec 36^\circ 40'$ .

13. Given  $L \cos 34^\circ 44' = 9.9147729$

$$L \cos 34^\circ 45' = 9.9146852,$$

find the value of  $L \cos 34^\circ 44' 27''$ .

14. Given  $L \sin 36^\circ 40' = 9.7760897$

$$L \cos 36^\circ 40' = 9.9042411,$$

find  $L \tan 36^\circ 40'$ .

15. Prove that the difference of tabular logarithms of any two ratios is equal to the difference of the logarithms of those two ratios.

16. If  $\sin \theta = .8$ , find  $\theta$

$$\text{given } \log 2 = .3010300$$

$$L \sin 53^\circ 7' = 9.9030136$$

$$L \sec 36^\circ 52' = 10.0968916.$$

17. Find the value of

$$\frac{\sin 34^\circ 17' \times \cos 77^\circ 23'}{\tan 27^\circ 12'}$$

$$\text{given } L \sin 12^\circ 37' = 9.3391$$

$$L \cos 55^\circ 43' = 9.7501$$

$$L \tan 62^\circ 48' = 10.28$$

$$\text{and } \log 23.94 = 1.379$$

## CHAPTER XIV

### PROPERTIES OF TRIANGLES

81. In a triangle  $ABC$ , there are six parts, the three sides and the three angles. It is usual to denote the angles of the triangle by  $A, B, C$  and the corresponding opposite sides by  $a, b, c$ . The six parts are not independent of one another. The various relations existing among them are deduced in the following articles.

82. In any triangle, prove that

$$\frac{a}{\sin A} = \frac{b}{\sin B} = \frac{c}{\sin C}.$$




Fig. (i)




Fig. (ii)




Fig. (iii)

Let  $ABC$  be any triangle. From  $A$  draw  $AD$  perpendicular to  $BC$  or  $BC$  produced if necessary [ Fig. (ii). ]

[ In Fig. (i),  $C$  is an *acute* angle, in Fig. (ii),  $C$  is an *obtuse* angle in Fig. (iii),  $C$  is a *right* angle. ]

From  $\triangle ABD$ ,  $AD = AB \sin ABD = c \sin B$ .

From  $\triangle ACD$ ,  $AD = AC \sin ACD = b \sin C$  [ Fig. (i). ]

or,  $= b \sin (\pi - C)$  [Fig.(ii)]

i.e.,  $= b \sin C$ .



$$\therefore b \sin C = c \sin B, \text{ i.e., } \frac{b}{\sin B} = \frac{c}{\sin C}.$$

Similarly, by drawing a perpendicular from  $B$  upon  $CA$ , we have  $\frac{a}{\sin A} = \frac{c}{\sin C}$ .

In Fig. (iii),  $C$  is a *right angle* ;

$$\therefore \sin A = \frac{a}{c} ; \sin B = \frac{b}{c} ; \sin C = 1.$$

$$\therefore \frac{a}{\sin A} = \frac{b}{\sin B} = c = \frac{c}{\sin C}.$$


Hence, in all cases,

$$\frac{a}{\sin A} = \frac{b}{\sin B} = \frac{c}{\sin C} \quad \dots (1)$$

Thus, in any triangle,

*the sides are proportional to the sines of the opposite angles.*

An alternative method of proof :



Let  $O$  be the centre and  $R$  be the radius of the circle circumscribing the triangle  $ABC$ .

Join  $BO$  and produce it to meet the circumference in  $D$ . Join  $CD$ . The  $\angle BCD$  is then a right angle.

$$\text{From } \triangle BCD, \sin BDC = \frac{BC}{BD} = \frac{a}{2R}.$$

But  $\angle BDC = \angle A$ , being in the same segment.

$$\therefore \frac{a}{2R} = \sin A, \text{ or, } \frac{a}{\sin A} = 2R.$$

Similarly, by joining  $AO$  and producing it to meet the circumference in  $E$ , and joining  $CE$ ,  $BE$ , it can be shown that

$$\frac{b}{\sin B} = 2R \text{ and } \frac{c}{\sin C} = 2R.$$

$$\therefore \frac{a}{\sin A} = \frac{b}{\sin B} = \frac{c}{\sin C} = 2R. \quad \dots (2)$$

**Note 1.** If angle  $A$  be obtuse,  $A$  and  $D$  fall on opposite sides of  $BC$  and  $ABCD$  being cyclic,  $\sin BDC = \sin (180^\circ - A) = \sin A$ , and the same result follows. In case  $A$  is a right angle, evidently  $2R = a = a/\sin A$ , and we get the same result.

**Note 2.** It follows from the relation (2) that

$$a = 2R \sin A, b = 2R \sin B, c = 2R \sin C, \\ \sin A = \frac{a}{2R}, \sin B = \frac{b}{2R}, \sin C = \frac{c}{2R}.$$

**83.** In any triangle, to prove that

$$a^2 = b^2 + c^2 - 2bc \cos A, \text{ or } \cos A = \frac{b^2 + c^2 - a^2}{2bc}.$$

$$b^2 = c^2 + a^2 - 2ca \cos B, \text{ or } \cos B = \frac{c^2 + a^2 - b^2}{2ca}$$

$$c^2 = a^2 + b^2 - 2ab \cos C, \text{ or } \cos C = \frac{a^2 + b^2 - c^2}{2ab}.$$

Take the figures of Art. 82.

First, let  $C$  be an *acute* angle [ *Fig. (i)* ]; then from Geometry,

$$AB^2 = BC^2 + CA^2 - 2BC \cdot CD.$$

Now, from  $\triangle ACD$ ,  $CD = AC \cos C = b \cos C$ .

$$\therefore c^2 = a^2 + b^2 - 2ab \cos C.$$

Next, let the angle  $C$  be an *obtuse* angle [ *Fig. (ii)* ]; then from Geometry,

$$AB^2 = BC^2 + CA^2 + 2BC \cdot CD.$$

Now, from  $\triangle ACD$ ,  $CD = AC \cos ACD$   
 $= b \cos (\pi - C) = -b \cos C.$

$$\therefore c^2 = a^2 + b^2 - 2ab \cos C.$$

Lastly, let  $C$  be a *right angle* [ Fig. (iii) ]; then from Geometry,

$$AB^2 = BC^2 + CA^2,$$

$$\text{i.e., } c^2 = a^2 + b^2 = a^2 + b^2 - 2ab \cos C.$$

$$[ \because \cos C = \cos 90^\circ = 0. ]$$

Hence, for all values of  $C$ , we have

$$c^2 = a^2 + b^2 - 2ab \cos C,$$

$$\therefore \cos C = \frac{a^2 + b^2 - c^2}{2ab}.$$

Similarly, the other two relations can be established.

**Obs.** This theorem expresses the cosines of the angles of a triangle in terms of the sides.

**84.** In any triangle, to prove that

$$a = b \cos C + c \cos B.$$

$$b = c \cos A + a \cos C.$$

$$c = a \cos B + b \cos A.$$

Take the figures of Art. 82.

In Fig. (i), where  $C$  is an *acute angle*,

$$BC = BD + CD$$

$$= AB \cos ABD + AC \cos ACD.$$

$$\therefore a = c \cos B + b \cos C.$$

In Fig. (ii), where  $C$  is an *obtuse angle*,

$$BC = BD - CD$$

$$= AB \cos ABD - AC \cos ACD$$

$$= c \cos B - b \cos (180^\circ - C)$$

$$= c \cos B + b \cos C.$$

In Fig. (iii), where  $C$  is a *right angle*,

$$BC = AB \cos B.$$

$$\therefore a = c \cos B = c \cos B + b \cos C.$$

$$[\because \cos C = \cos 90^\circ = 0.]$$

Thus in all cases,

$$a = b \cos C + c \cos B.$$

Similarly, the other two relations can be established.

85. From Art. 83 and note of Art. 82, it follows that

$$\tan A = \frac{\sin A}{\cos A} = \frac{\frac{a}{2R}}{\frac{b^2 + c^2 - a^2}{2bc}} = \frac{abc}{R(b^2 + c^2 - a^2)}.$$

$$\text{Similarly, } \tan B = \frac{abc}{R(c^2 + a^2 - b^2)};$$

$$\tan C = \frac{abc}{R(a^2 + b^2 - c^2)}.$$

86. Trigonometrical ratios of half angles of a triangle in terms of the sides.

$$\begin{aligned} \text{We have, } 2 \sin^2 \frac{A}{2} &= 1 - \cos A = 1 - \frac{b^2 + c^2 - a^2}{2bc} \\ &= \frac{2bc - b^2 - c^2 + a^2}{2bc} = \frac{a^2 - (b^2 - 2bc + c^2)}{2bc} \\ &= \frac{a^2 - (b - c)^2}{2bc} = \frac{(a - b + c)(a + b - c)}{2bc}. \end{aligned}$$

Let  $s$  denote the semi-perimeter of the triangle ;

$$\text{then } 2s = a + b + c.$$

$$\begin{aligned} \text{Now, } a - b + c &= a + b + c - 2b = 2s - 2b = 2(s - b), \\ a + b - c &= a + b + c - 2c = 2s - 2c = 2(s - c). \end{aligned}$$

$$\text{Hence, } 2 \sin^2 \frac{A}{2} = \frac{2(s - b) \cdot 2(s - c)}{2bc}$$

$$\text{i.e., } \sin^2 \frac{A}{2} = \frac{(s - b)(s - c)}{bc}.$$

$$\therefore \sin \frac{A}{2} = \sqrt{\frac{(s - b)(s - c)}{bc}}.$$

The positive value of the square root must be taken ; for  $A$ , being an angle of a triangle, is less than  $180^\circ$  ; and hence  $\frac{1}{2}A < 90^\circ$  and consequently,  $\sin \frac{1}{2}A$  must always be positive.

$$\begin{aligned}\text{Again, } 2 \cos^2 \frac{A}{2} &= 1 + \cos A \\ &= 1 + \frac{b^2 + c^2 - a^2}{2bc} = \frac{2bc + b^2 + c^2 - a^2}{2bc} \\ &= \frac{(b+c)^2 - a^2}{2bc} = \frac{(b+c+a)(b+c-a)}{2bc}.\end{aligned}$$

$$\text{Now, } b+c-a = a+b+c-2a = 2s-2a = 2(s-a).$$

$$\therefore 2 \cos^2 \frac{A}{2} = \frac{2s \cdot 2(s-a)}{2bc}, \text{ i.e., } \cos^2 \frac{A}{2} = \frac{s(s-a)}{bc}.$$

$$\therefore \cos \frac{A}{2} = \sqrt{\frac{s(s-a)}{bc}}.$$

Here also the positive value of the square root must be taken, for  $\frac{1}{2}A$  being less than  $90^\circ$ ,  $\cos \frac{1}{2}A$  is always positive.

$$\begin{aligned}\text{Again, } \tan \frac{A}{2} &= \frac{\sin \frac{A}{2}}{\cos \frac{A}{2}} \\ &= \frac{\sqrt{(s-b)(s-c)}}{bc} + \frac{\sqrt{s(s-a)}}{bc} \\ &= \sqrt{\frac{(s-b)(s-c)}{s(s-a)}}.\end{aligned}$$

Similarly, the trigonometrical ratios of  $\frac{B}{2}$ ,  $\frac{C}{2}$  can be obtained in terms of the sides.

**Note.** Without assuming the values of  $\sin \frac{1}{2}A$ ,  $\cos \frac{1}{2}A$ , the value of  $\tan \frac{1}{2}A$  can be obtained by substituting the values of  $\cos A$  in terms of the sides from Art. 83 in the relation  $\tan^2 \frac{1}{2}A = \frac{1 - \cos A}{1 + \cos A}$  and then extracting the square root after simplification.

Thus, we have

$$\left. \begin{aligned} \sin \frac{A}{2} &= \sqrt{\frac{(s-b)(s-c)}{bc}} \\ \sin \frac{B}{2} &= \sqrt{\frac{(s-c)(s-a)}{ca}} \\ \sin \frac{C}{2} &= \sqrt{\frac{(s-a)(s-b)}{ab}} \end{aligned} \right\} \dots \dots (1)$$

$$\left. \begin{aligned} \cos \frac{A}{2} &= \sqrt{\frac{s(s-a)}{bc}} \\ \cos \frac{B}{2} &= \sqrt{\frac{s(s-b)}{ca}} \\ \cos \frac{C}{2} &= \sqrt{\frac{s(s-c)}{ab}} \end{aligned} \right\} \dots \dots (2)$$

$$\left. \begin{aligned} \tan \frac{A}{2} &= \sqrt{\frac{(s-b)(s-c)}{s(s-a)}} \\ \tan \frac{B}{2} &= \sqrt{\frac{(s-c)(s-a)}{s(s-b)}} \\ \tan \frac{C}{2} &= \sqrt{\frac{(s-a)(s-b)}{s(s-c)}} \end{aligned} \right\} \dots \dots (3)$$

87. Sine of an angle of a triangle in terms of the sides.

$$\begin{aligned} \sin A &= 2 \sin \frac{A}{2} \cos \frac{A}{2} \\ &= 2 \sqrt{\frac{(s-b)(s-c)}{bc}} \cdot \sqrt{\frac{s(s-a)}{bc}}. \text{ [Art. 86.]} \end{aligned}$$

$$\therefore \sin A = \frac{2}{bc} \sqrt{s(s-a)(s-b)(s-c)}.$$


$$\text{Similarly, } \sin B = \frac{2}{ca} \sqrt{s(s-a)(s-b)(s-c)}.$$

$$\sin C = \frac{2}{ab} \sqrt{s(s-a)(s-b)(s-c)}.$$

$\sqrt{s(s-a)(s-b)(s-c)}$ , being the expression for the area of the triangle [See Art. 88], is usually denoted by the Greek letter  $\Delta$ . Hence, the above formulæ may be written as

$$\sin A = \frac{2\Delta}{bc}, \quad \sin B = \frac{2\Delta}{ca}, \quad \sin C = \frac{2\Delta}{ab}.$$

### 88. Area of a triangle.



Let  $ABC$  be a triangle and let  $\Delta$  denote its area. Draw  $AD$  perpendicular to  $BC$ ; then from  $\triangle ACD$ ,

$$AD = AC \sin C = b \sin C.$$

$$\text{Now, } \Delta = \frac{1}{2}BC \cdot AD = \frac{1}{2}ab \sin C.$$

Similarly, by drawing perpendicular from  $B$  and  $C$  to the opposite sides, it can be shown that

$$\Delta = \frac{1}{2}bc \sin A = \frac{1}{2}ac \sin B.$$

$$\text{Otherwise, } \Delta = \frac{1}{2}ab \sin C$$

$$= \frac{1}{2}ac \sin B \quad [\because b \sin C = c \sin B]$$

$$= \frac{1}{2}bc \sin A \quad [\because a \sin B = b \sin A]$$

$$\text{Thus, } \Delta = \frac{1}{2}bc \sin A = \frac{1}{2}ca \sin B = \frac{1}{2}ab \sin C \quad \dots (i)$$

$= \frac{1}{2}(\text{product of two sides}) \times \text{sine of included angle}.$

$$\text{Again, } \Delta = \frac{1}{2}bc \sin A = bc \sin \frac{A}{2} \cos \frac{A}{2}$$

$$= bc \sqrt{\frac{(s-b)(s-c)}{bc}} \cdot \sqrt{\frac{s(s-a)}{bc}}$$

$$= \sqrt{s(s-a)(s-b)(s-c)}. \quad \dots (ii)$$

Substituting in the expression  $s = \frac{1}{2}(a + b + c)$ ,  
we get

$$\Delta = \frac{1}{4} \sqrt{(a + b + c)(b + c - a)(c + a - b)(a + b - c)}$$

$$= \frac{1}{4} \{2b^2c^2 + 2c^2a^2 + 2a^2b^2 - a^4 - b^4 - c^4\}^{\frac{1}{2}} \dots \text{(iii)}$$

Again,

$$\Delta = \frac{1}{2}bc \sin A = \frac{1}{2}bc \cdot \frac{a}{2R} [\text{Art. 82}] = \frac{abc}{4R} \dots \text{(iv)}$$

**Note.** In some text books,  $S$  is used to denote the area of a triangle; but to avoid confusion between  $S$  and  $s$  in writing, the symbol  $\Delta$  is preferable.

~~In any triangle, to prove that~~  

$$\tan \frac{B - C}{2} = \frac{b - c}{b + c} \cot \frac{A}{2}.$$

We have, in any triangle,

$$\frac{b}{c} = \frac{\sin B}{\sin C}.$$

$$\therefore \frac{b - c}{b + c} = \frac{\sin B - \sin C}{\sin B + \sin C} = \frac{2 \cos \frac{B + C}{2} \cdot \sin \frac{B - C}{2}}{2 \sin \frac{B + C}{2} \cdot \cos \frac{B - C}{2}}$$

$$= \cot \frac{B + C}{2} \tan \frac{B - C}{2}$$

$$= \tan \frac{A}{2} \tan \frac{B - C}{2} \left[ \because \frac{A}{2} + \frac{B + C}{2} = 90^\circ \right]$$

$$\tan \frac{B - C}{2} = \frac{b - c}{b + c} \frac{1}{\tan \frac{A}{2}} = \frac{b - c}{b + c} \cot \frac{A}{2}.$$

Similarly,

$$\tan \frac{C - A}{2} = \frac{c - a}{c + a} \cot \frac{B}{2}; \quad \tan \frac{A - B}{2} = \frac{a - b}{a + b} \cot \frac{C}{2}.$$

90. The three sets of formulæ in Arts. 82, 83, 84 have been established directly from the figures. These three sets,



however, are not independent, for, from any one set, the other two sets can be deduced.

For example, let us deduce the formulæ of Art. 83 from those of Art. 84.

$$\begin{aligned}\text{By Art. 84,} \quad a &= b \cos C + c \cos B \\ b &= c \cos A + a \cos C \\ c &= a \cos B + b \cos A.\end{aligned}$$

Multiplying these in succession by  $a$ ,  $b$  and  $c$ , and subtracting the first result from the sum of the other two, we have,

$$\begin{aligned}b^2 + c^2 - a^2 &= b(c \cos A + a \cos C) + c(a \cos B + b \cos A) \\ &\quad - a(b \cos C + c \cos B) = 2bc \cos A.\end{aligned}$$

$$\therefore \cos A = \frac{b^2 + c^2 - a^2}{2bc}; \text{ similarly for } \cos B, \cos C.$$

**Note.** For other cases, see *Appendix*.

91. In working out identities which involve both the sides and angles of a triangle, it is sometimes convenient to express the sides in terms of the angles, or the angles in terms of the sides.

Also, it is sometimes found convenient to express the values of  $\tan \frac{A}{2}$ ,  $\tan \frac{B}{2}$ ,  $\tan \frac{C}{2}$  in a form in which the denominator is constant and numerator is free from radical. Thus, multiplying the numerator and the denominator of the value of  $\tan \frac{A}{2}$  by  $\sqrt{(s-b)(s-c)}$  and noting that

$$\sqrt{s(s-a)(s-b)(s-c)} = \Delta, \text{ we have}$$

$$\tan \frac{A}{2} = \frac{(s-b)(s-c)}{\Delta}; \text{ similarly, } \tan \frac{B}{2} = \frac{(s-c)(s-a)}{\Delta};$$

$$\tan \frac{C}{2} = \frac{(s-a)(s-b)}{\Delta}.$$

Again, multiplying the numerator and the denominator of the value of  $\cot \frac{A}{2}$  by  $\sqrt{s(s-a)}$ , we have

$$\cot \frac{A}{2} = \frac{s(s-a)}{\Delta}.$$

$$\text{Similarly, } \cot \frac{B}{2} = \frac{s(s-b)}{\Delta}; \cot \frac{C}{2} = \frac{s(s-c)}{\Delta}.$$

**Ex. 1.** Show that in any triangle

$$a(\sin B - \sin C) + b(\sin C - \sin A) + c(\sin A - \sin B) = 0.$$

$$\begin{aligned} \text{Left side} &= (a \sin B - b \sin A) + (b \sin C - c \sin B) \\ &\quad + (c \sin A - a \sin C) \\ &= 0 + 0 + 0 \quad \left[ \because \text{ by Art. 82, } \frac{a}{\sin A} = \frac{b}{\sin B} = \frac{c}{\sin C} \right] \\ &= 0. \end{aligned}$$

**Ex. 2.** Show that in any triangle

$$a \sin (B - C) + b \sin (C - A) + c \sin (A - B) = 0.$$

$$\begin{aligned} a &= 2R \sin A \text{ [by Art. 82]} = 2R \sin (B + C), \text{ [ } \because A + B + C = \pi \text{ ]} \\ \therefore \text{ 1st term of left side} &= 2R \sin (B + C) \sin (B - C) \\ &= 2R (\sin^2 B - \sin^2 C). \\ &\quad \text{[ by Ex. 2, Art. 35 ]} \end{aligned}$$

$$\begin{aligned} \text{Similarly, } \quad \text{2nd term} &= 2R (\sin^2 C - \sin^2 A) \\ \text{3rd term} &= 2R (\sin^2 A - \sin^2 B). \end{aligned}$$

Now, adding together the three terms, the required result follows.

**Ex. 3.** In any triangle, prove that

$$(b-c) \cot \frac{A}{2} + (c-a) \cot \frac{B}{2} + (a-b) \cot \frac{C}{2} = 0.$$

Substituting the values of  $\cot \frac{A}{2}$ ,  $\cot \frac{B}{2}$ ,  $\cot \frac{C}{2}$ , as given in Art. 91 we have, the left side

$$\begin{aligned} &= (b-c) \cdot \frac{s(s-a)}{\Delta} + (c-a) \cdot \frac{s(s-b)}{\Delta} + (a-b) \cdot \frac{s(s-c)}{\Delta} \\ &= \frac{s}{\Delta} \left[ (b-c)(s-a) + (c-a)(s-b) + (a-b)(s-c) \right] \\ &= \frac{s}{\Delta} \cdot 0 = 0. \end{aligned}$$

**Ex. 4.** *If the cosines of two of the angles of a triangle are inversely proportional to the opposite sides, show that the triangle is either isosceles or right-angled.*

We have, by the question,

$$\frac{\cos A}{\cos B} = \frac{b}{a} = \frac{\sin B}{\sin A} \quad [\text{by Art. 82.}]$$

$$\therefore \sin A \cos A = \sin B \cos B \text{ or } \sin 2A = \sin 2B.$$

$$\therefore \sin 2A - \sin 2B = 0.$$

$$\therefore 2 \cos (A+B) \sin (A-B) = 0.$$

$$\therefore \text{either } \cos (A+B) = 0, \text{ i.e., } (A+B) = 90^\circ.$$

$$\therefore \text{the triangle is right-angled ;}$$

$$\text{or, } \sin (A-B) = 0, \text{ i.e., } A-B=0, \text{ i.e., } A=B.$$

$$\therefore \text{the triangle is isosceles.}$$

**Ex. 5.** *If the sides of a triangle are in A.P., show that  $\cot \frac{A}{2}$ ,  $\cot \frac{B}{2}$ ,  $\cot \frac{C}{2}$  are also in A.P.*

$$\cot \frac{A}{2}, \cot \frac{B}{2}, \cot \frac{C}{2} \text{ are in A. P.,}$$

$$\text{if } \cot \frac{B}{2} - \cot \frac{A}{2} = \cot \frac{C}{2} - \cot \frac{B}{2}.$$

$$\text{i.e., if } \frac{s(s-b)}{\Delta} - \frac{s(s-a)}{\Delta} = \frac{s(s-c)}{\Delta} - \frac{s(s-b)}{\Delta},$$

$$\text{i.e., if } (s-b) - (s-a) = (s-c) - (s-b)$$

$$\text{i.e., if } a - b = b - c,$$

$$\text{i.e., if } a, b, c \text{ are in A.P.}$$

**Ex. 6.** Show that

$$b^2 \sin 2C + c^2 \sin 2B = 4\Delta.$$

$$\text{Left side} = b^2 \cdot 2 \sin C \cos C + c^2 \cdot 2 \sin B \cos B$$

$$= 2b \sin C \cdot b \cos C + 2c \sin B \cdot c \cos B$$

$$= 2b \sin C (b \cos C + c \cos B)$$

$$[ \because c \sin B = b \sin C ]$$

$$= 2ab \sin C \quad [ \text{by Art. 84} ]$$

$$= 4 \cdot \frac{1}{2} ab \sin C = 4\Delta. \quad [ \text{by Art. 88} ]$$

### Examples XIV(a)

In any triangle, prove that (Ex. 1 to 21) :—

$$1. \sin \frac{B-C}{2} = \frac{b-c}{a} \cos \frac{A}{2}.$$

$$2. \cos \frac{B-C}{2} = \frac{b+c}{a} \sin \frac{A}{2}.$$

$$3. (b+c) \cos A + (c+a) \cos B + (a+b) \cos C = a+b+c.$$

$$4. \frac{a+b}{a-b} = \tan \frac{A+B}{2} \cot \frac{A-B}{2}.$$

$$5. a^2 + b^2 + c^2 = 2(bc \cos A + ca \cos B + ab \cos C).$$

$$6. (b+c-a) \tan \frac{A}{2} = (c+a-b) \tan \frac{B}{2} = (a+b-c) \tan \frac{C}{2}.$$

7.  $\frac{a \sin (B-C)}{b^2-c^2} = \frac{b \sin (C-A)}{c^2-a^2} = \frac{c \sin (A-B)}{a^2-b^2}.$
8.  $a^2(\sin^2 B - \sin^2 C) + b^2(\sin^2 C - \sin^2 A) + c^2(\sin^2 A - \sin^2 B) = 0.$
9.  $a^2(\cos^2 B - \cos^2 C) + b^2(\cos^2 C - \cos^2 A) + c^2(\cos^2 A - \cos^2 B) = 0.$
10.  $\frac{a^2 \sin (B-C)}{\sin B + \sin C} + \frac{b^2 \sin (C-A)}{\sin C + \sin A} + \frac{c^2 \sin (A-B)}{\sin A + \sin B} = 0.$
11.  $a \sin \frac{A}{2} \sin \frac{B-C}{2} + b \sin \frac{B}{2} \sin \frac{C-A}{2} + c \sin \frac{C}{2} \sin \frac{A-B}{2} = 0.$
12.  $\frac{b^2-c^2}{a^2} \sin 2A + \frac{c^2-a^2}{b^2} \sin 2B + \frac{a^2-b^2}{c^2} \sin 2C = 0.$
13.  $a^3 \sin (B-C) + b^3 \sin (C-A) + c^3 \sin (A-B) = 0.$
14.  $a^3 \cos (B-C) + b^3 \cos (C-A) + c^3 \cos (A-B) = 3abc.$
15.  $\frac{a^2 \sin (B-C)}{\sin A} + \frac{b^2 \sin (C-A)}{\sin B} + \frac{c^2 \sin (A-B)}{\sin C} = 0.$
16.  $(b^2-c^2) \cot A + (c^2-a^2) \cot B + (a^2-b^2) \cot C = 0.$
17.  $\frac{b^2-c^2}{\cos B + \cos C} + \frac{c^2-a^2}{\cos C + \cos A} + \frac{a^2-b^2}{\cos A + \cos B} = 0.$
18.  $(s-a) \tan \frac{A}{2} = (s-b) \tan \frac{B}{2} = (s-c) \tan \frac{C}{2}.$
19.  $\frac{b-c}{a} \cos^2 \frac{A}{2} + \frac{c-a}{b} \cos^2 \frac{B}{2} + \frac{a-b}{c} \cos^2 \frac{C}{2} = 0.$
20.  $bc \cos^2 \frac{A}{2} + ca \cos^2 \frac{B}{2} + ab \cos^2 \frac{C}{2} = s^2.$
21.  $\frac{1}{a} \cos^2 \frac{A}{2} + \frac{1}{b} \cos^2 \frac{B}{2} + \frac{1}{c} \cos^2 \frac{C}{2} = \frac{s^2}{abc}.$
22. If  $A$  be  $60^\circ$ , show that  $b+c-2a \cos \frac{B-C}{2}.$

23. Show that a triangle having its sides equal to 3, 5, 7 is an obtuse-angled triangle and determine the obtuse angle.

24. Given  $(a+b+c)(b+c-a) = 3bc$ , find  $A$ .

25. If  $c^4 - 2(a^2 + b^2)c^2 + a^4 + a^2b^2 + b^4 = 0$ , prove that  $C = 60^\circ$ , or,  $120^\circ$ .

26. If  $a^4 + b^4 + c^4 = 2c^2(a^2 + b^2)$ , prove that  $C = 45^\circ$ , or,  $135^\circ$ .

27. The sides of a triangle are  $2x+3$ ,  $x^2+3x+3$ ,  $x^2+2x$ ; show that the greatest angle is  $120^\circ$ .

28. If  $\frac{1}{a+c} + \frac{1}{b+c} = \frac{3}{a+b+c}$ , show that  $C = 60^\circ$ .

29. If  $a = 2b$  and  $A = 3B$ , find the angles of the triangle.

30. If the cosines of two of the angles of a triangle are proportional to the opposite sides, show that the triangle is isosceles.

31. If  $\cos A = \frac{\sin B}{2 \sin C}$ , show that the triangle is isosceles.

32. If  $(a^2 + b^2) \sin(A-B) = (a^2 - b^2) \sin(A+B)$ , prove that the triangle is either isosceles or right-angled.

33. If  $\cos A + 2 \cos C : \cos A + 2 \cos B = \sin B : \sin C$ , prove that the triangle is either isosceles or right-angled.

34. If  $a^2, b^2, c^2$  be in A.P., prove that  $\cot A, \cot B, \cot C$  are also in A.P.

35. If  $a \cos^2 \frac{C}{2} + c \cos^2 \frac{A}{2} = \frac{3b}{2}$ , show that the sides of the triangle are in A.P.

36. If  $\sin A : \sin C = \sin(A-B) : \sin(B-C)$ , show that  $a^2, b^2, c^2$  are in A.P.

37. If  $a, b, c$  are in A.P., show that  
 $\cos A \cot \frac{1}{2}A, \cos B \cot \frac{1}{2}B, \cos C \cot \frac{1}{2}C$  are in A.P.

$$[\cos A \cot \frac{1}{2}A = (1 - 2 \sin^2 \frac{1}{2}A) \cot \frac{1}{2}A = \cot \frac{1}{2}A - \sin A.]$$

38. Assuming  $\Delta = \frac{1}{2}bc \sin A$  and using the value of  $\cos A$  in terms of sides, show that

$$\Delta = \sqrt{s(s-a)(s-b)(s-c)}.$$

39. Find the area of the triangle whose sides are

$$\frac{y+z}{z}, \frac{z+x}{x}, \frac{x+y}{y}.$$

40. In a triangle, if  $a = 13, b = 14, c = 15$ , find its area.

Prove that in any triangle :

$$41. \frac{a^2 - b^2}{2} \frac{\sin A \sin B}{\sin(A-B)} = \Delta.$$

$$42. 4\Delta (\cot A + \cot B + \cot C) = a^2 + b^2 + c^2.$$

$$43. a \cos A + b \cos B + c \cos C = 4R \sin A \sin B \sin C.$$

$$44. a \sin B \sin C + b \sin C \sin A + c \sin A \sin B = \frac{3\Delta}{R}.$$

$$45. (a \sin A + b \sin B + c \sin C)^2 \\ = (a^2 + b^2 + c^2)(\sin^2 A + \sin^2 B + \sin^2 C).$$

$$46. \frac{\cos B \cos C}{bc} + \frac{\cos C \cos A}{ca} + \frac{\cos A \cos B}{ab} = \frac{1}{4R^2}.$$

[ Use  $\Sigma \cot B \cot C = 1$  ; ex. 2, Ex. X. ]

$$47. \frac{b^2 - c^2}{a} \cos A + \frac{c^2 - a^2}{b} \cos B + \frac{a^2 - b^2}{c} \cos C = 0.$$

$$48. \frac{\cos A}{a} + \frac{a}{bc} = \frac{\cos B}{b} + \frac{b}{ca} = \frac{\cos C}{c} + \frac{c}{ab}.$$

$$49. 4\Delta = a^2 \cot A + b^2 \cot B + c^2 \cot C.$$

$$50. \left( \frac{a^2}{\sin A} + \frac{b^2}{\sin B} + \frac{c^2}{\sin C} \right) \sin \frac{A}{2} \sin \frac{B}{2} \sin \frac{C}{2} = \Delta.$$

### 92. Circum-radius of a triangle.


From Art. 82, we have

$$\frac{a}{\sin A} = \frac{b}{\sin B} = \frac{c}{\sin C} = 2R. \quad \dots (i)$$

$$\text{again, } R = \frac{a}{2 \sin A} = \frac{abc}{2bc \sin A} = \frac{abc}{4\Delta}. \quad \dots (ii)$$

### 93. In-radius of a triangle.

Let  $I$  be the centre and  $r$  the radius of the circle inscribed in the triangle  $ABC$ ; let  $D, E, F$  be the points of contact of the in-circle with the sides  $BC, CA, AB$  respectively.



Then,  $ID = IE = IF = r$ .

Join  $IA, IB, IC$ .

$$\begin{aligned} \triangle ABC &= \triangle IBC + \triangle ICA + \triangle IAB \\ &= \frac{1}{2}BC.ID + \frac{1}{2}CA.IE + \frac{1}{2}AB.IF \\ &= \frac{1}{2}ar + \frac{1}{2}br + \frac{1}{2}cr \\ &= \frac{1}{2}r(a + b + c) = rs. \end{aligned}$$

Thus,  $\Delta = rs$ .

$$\therefore r = \frac{\Delta}{s}. \quad \dots (i)$$

Again,  $a = BC = BD + DC$

$$= r \cot \frac{1}{2}B + r \cot \frac{1}{2}C, \text{ from } \triangle IBD, ICD,$$

$$= r \left[ \frac{\cos \frac{1}{2}B}{\sin \frac{1}{2}B} + \frac{\cos \frac{1}{2}C}{\sin \frac{1}{2}C} \right]$$

$$= r \left[ \frac{\cos \frac{1}{2}B \sin \frac{1}{2}C + \sin \frac{1}{2}B \cos \frac{1}{2}C}{\sin \frac{1}{2}B \sin \frac{1}{2}C} \right]$$

$$= r \frac{\sin (\frac{1}{2}B + \frac{1}{2}C)}{\sin \frac{1}{2}B \sin \frac{1}{2}C} = r \frac{\cos \frac{1}{2}A}{\sin \frac{1}{2}B \sin \frac{1}{2}C}.$$

$$[\because \frac{1}{2}A + \frac{1}{2}B + \frac{1}{2}C = 90^\circ, \therefore \sin (\frac{1}{2}B + \frac{1}{2}C) = \sin (90^\circ - \frac{1}{2}A) = \cos \frac{1}{2}A.]$$



$$\therefore r = a \sin \frac{1}{2}B \sin \frac{1}{2}C \sec \frac{1}{2}A = a \frac{\sin \frac{1}{2}B \sin \frac{1}{2}C}{\cos \frac{1}{2}A}.$$

Since by Art. 92 (i),  $a = 2R \sin A = 4R \sin \frac{1}{2}A \cos \frac{1}{2}A$ ,

$$\therefore r = 4R \sin \frac{1}{2}A \sin \frac{1}{2}B \sin \frac{1}{2}C. \quad \dots (ii)$$

Since, from the figure,  $AF = AE$ ,  $BD = BF$ ,  $CD = CE$  and since the sum of these six quantities is equal to the perimeter,

$$\therefore AF + BD + CD = \text{semi-perimeter} = s,$$

$$\text{i.e., } AF + BC, \text{ or, } AF + a = s.$$

$$\therefore AF = s - a = AE.$$

Similarly,  $BF = s - b = BD$ ;  $CF = s - c = CD$ .

From  $\triangle AIF$ ,  $IF = AF \tan IAF$ .

$$\therefore r = (s - a) \tan \frac{1}{2}A.$$

$$\left. \begin{array}{l} \text{Similarly, } r = (s - b) \tan \frac{1}{2}B, \\ \text{and } r = (s - c) \tan \frac{1}{2}C. \end{array} \right\} \quad \dots (iii)$$

**Note.** Distances of the In-centre from the vertices.


From  $\triangle AIF$ ,  $IA = IF \operatorname{cosec} IAF$ .  $\therefore IA = r \operatorname{cosec} \frac{1}{2}A$

Similarly,  $IB = r \operatorname{cosec} \frac{1}{2}B$  and  $IC = r \operatorname{cosec} \frac{1}{2}C$ .

#### 94. Ex-radii of a triangle.

Let  $I_1$  be the centre and  $r_1$  the radius of the escribed circle (opposite to the angle  $A$ ) of the  $\triangle ABC$ ; let  $D, E, F$  be the points of contact of the circle with the sides  $BC$ , and  $AC$  and  $AB$  produced.

Let  $r_2, r_3$  denote the radii of the escribed circles opposite to the angles  $B$  and  $C$  respectively.



Now,  $I_1D = I_1E = I_1F = r_1$ ; join  $AI_1, BI_1, CI_1$ .

$$\begin{aligned}\triangle ABC &= \triangle I_1AB + \triangle I_1AC - \triangle I_1BC \\ &= \frac{1}{2}AB \cdot I_1F + \frac{1}{2}AC \cdot I_1E - \frac{1}{2}BC \cdot I_1D \\ &= \frac{1}{2}cr_1 + \frac{1}{2}br_1 - \frac{1}{2}ar_1 \\ &= \frac{1}{2}r_1(b+c-a) = \frac{1}{2}r_1(b+c+a-2a) \\ &= \frac{1}{2}r_1(2s-2a) \\ &= r_1(s-a).\end{aligned}$$

Thus,  $\triangle = r_1(s-a)$ ,

$$\therefore r_1 = \frac{\triangle}{s-a}.$$

$$\left. \begin{array}{l} \text{Similarly,} \\ \text{and} \end{array} \right\} \begin{array}{l} r_2 = \frac{\triangle}{s-b} \\ r_3 = \frac{\triangle}{s-c} \end{array} \quad \dots (i)$$

Again,  $a = BC = BD + CD$ ,

$$\begin{aligned}&= r_1 \cot I_1BD + r_1 \cot I_1CD, \\ &\quad \text{from } \triangle^s I_1BD, I_1CD \\ &= r_1 \cot (90^\circ - \tfrac{1}{2}B) + r_1 \cot (90^\circ - \tfrac{1}{2}C),\end{aligned}$$

because,  $\angle I_1ED = \frac{1}{2}(180^\circ - B) = 90^\circ - \frac{1}{2}B$ ,

and  $\angle I_1CD = \frac{1}{2}(180^\circ - C) = 90^\circ - \frac{1}{2}C$ .

$$\begin{aligned}\therefore a &= r_1(\tan \tfrac{1}{2}B + \tan \tfrac{1}{2}C) \\ &= r_1 \left[ \frac{\sin \tfrac{1}{2}B}{\cos \tfrac{1}{2}B} + \frac{\sin \tfrac{1}{2}C}{\cos \tfrac{1}{2}C} \right] \\ &= r_1 \left[ \frac{\sin \tfrac{1}{2}B \cos \tfrac{1}{2}C + \sin \tfrac{1}{2}C \cos \tfrac{1}{2}B}{\cos \tfrac{1}{2}B \cos \tfrac{1}{2}C} \right] \\ &= r_1 \frac{\sin (\tfrac{1}{2}B + \tfrac{1}{2}C)}{\cos \tfrac{1}{2}B \cos \tfrac{1}{2}C} \\ &= r_1 \frac{\cos \tfrac{1}{2}A}{\cos \tfrac{1}{2}B \cos \tfrac{1}{2}C}, \text{ as in Art. 93.}\end{aligned}$$

$$\therefore r_1 = a \cos \tfrac{1}{2}B \cos \tfrac{1}{2}C \sec \tfrac{1}{2}A.$$

Putting  $a = 2R \sin A = 4R \sin \frac{1}{2}A \cos \frac{1}{2}A$ ,

$$\left. \begin{aligned} r_1 &= 4R \sin \frac{1}{2}A \cos \frac{1}{2}B \cos \frac{1}{2}C. \\ \text{Similarly, } r_2 &= 4R \cos \frac{1}{2}A \sin \frac{1}{2}B \cos \frac{1}{2}C, \\ \text{and } r_3 &= 4R \cos \frac{1}{2}A \cos \frac{1}{2}B \sin \frac{1}{2}C. \end{aligned} \right\} \dots (ii)$$

Again,  $AE = AC + CE = b + CD$  [  $\because CE = CD$  ]

and  $AF = AB + BF = c + BD$  [  $\because BF = BD$  ]

But  $AE = AF$ ; therefore, by addition, we get

$$2AE = b + c + BD + CD = b + c + a = 2s.$$

$$\therefore AE = s.$$

Again, from  $\triangle AI_1E$ ,  $I_1E = AE \tan I_1AE$ .

$$\left. \begin{aligned} \therefore r_1 &= s \tan \frac{1}{2}A. \\ \text{Similarly, } r_2 &= s \tan \frac{1}{2}B, \\ \text{and } r_3 &= s \tan \frac{1}{2}C. \end{aligned} \right\} \dots (iii)$$

**Note.** Distances of Ex-centres from the vertices.

From  $\triangle AI_1F$ ,  $I_1A = I_1F \operatorname{cosec} I_1AF$ .

$$\begin{aligned} \therefore I_1A &= r_1 \operatorname{cosec} \frac{1}{2}A \\ &= 4R \cos \frac{1}{2}B \cos \frac{1}{2}C. \end{aligned} \quad [ \text{by formula (vi)} ]$$

From  $\triangle BI_1F$ ,  $I_1B = I_1F \operatorname{cosec} I_1BF$ .

$$\therefore I_1B = r_1 \sec \frac{1}{2}B \quad [ \because \angle I_1BF = (90^\circ - \frac{1}{2}B) ]$$

Similarly,  $I_1C = r_1 \sec \frac{1}{2}C$ .

In the same way,  $I_2B = r_2 \operatorname{cosec} \frac{1}{2}B$ ,  $I_2C = r_2 \operatorname{cosec} \frac{1}{2}C$ .

**Ex. 1.** Prove that  $\frac{1}{r_1} + \frac{1}{r_2} + \frac{1}{r_3} = \frac{1}{r}$ .

By formula (i), Art. 94,

$$\begin{aligned} \text{left side} &= \frac{s-a}{\Delta} + \frac{s-b}{\Delta} + \frac{s-c}{\Delta} \\ &= \frac{3s - (a+b+c)}{\Delta} = \frac{3s - 2s}{\Delta} = \frac{s}{\Delta} = \frac{1}{r}. \end{aligned}$$

**Ex. 2.** Prove that  $4 \cos \frac{1}{2}A \cos \frac{1}{2}B \cos \frac{1}{2}C = \frac{s}{R}$ .

$$\begin{aligned} \text{Left side} &= 4 \cdot \sqrt{\frac{s(s-a)}{bc}} \cdot \sqrt{\frac{s(s-b)}{ca}} \cdot \sqrt{\frac{s(s-c)}{ab}} \\ &= \frac{4s}{abc} \cdot \sqrt{s(s-a)(s-b)(s-c)} \\ &= \frac{4s}{abc} \cdot \Delta = s \cdot \frac{4\Delta}{abc} = \frac{s}{R} \text{ by formula (ii), Art. 92.} \end{aligned}$$

**Ex. 3.** Show that

$$\frac{bc - r_2 r_3}{r_1} = \frac{ca - r_3 r_1}{r_2} = \frac{ab - r_1 r_2}{r_3}.$$

$$r_2 r_3 = \frac{\Delta^2}{(s-b)(s-c)} = s(s-a).$$

$$\begin{aligned} \therefore bc - r_2 r_3 &= \frac{1}{4}[4bc - 2s(2s - 2a)] \\ &= \frac{1}{4}[4bc - (a+b+c)(b+c-a)] \\ &= \frac{1}{4}[4bc + a^2 - (b+c)^2] = \frac{1}{4}[a^2 - (b-c)^2] \\ &= \frac{1}{4}[(a+b-c)(a-b+c)] = (s-b)(s-c). \end{aligned}$$

$$\begin{aligned} \therefore \frac{bc - r_2 r_3}{r_1} &= \frac{(s-b)(s-c)}{r_1} = \frac{(s-a)(s-b)(s-c)}{\Delta} \\ &= \frac{\Delta}{s} = r. \end{aligned}$$

Similarly the other ratios are equal to the same quantity.

**Ex. 4.** Prove that in any triangle

$$r_1 + r_2 + r_3 - r = 4R.$$

$$\begin{aligned} \text{Left side} &= \left( \frac{\Delta}{s-a} + \frac{\Delta}{s-b} \right) + \left( \frac{\Delta}{s-c} - \frac{\Delta}{s} \right) \\ &= \Delta \cdot \frac{2s - (a+b)}{(s-a)(s-b)} + \Delta \cdot \frac{c}{s(s-c)} \\ &= \Delta c \left[ \frac{1}{(s-a)(s-b)} + \frac{1}{s(s-c)} \right] \quad [\because 2s = a+b+c.] \\ &= \Delta c \left[ \frac{s(s-c) + (s-a)(s-b)}{s(s-a)(s-b)(s-c)} \right] \end{aligned}$$

$$\begin{aligned}\text{Now, Numerator} &= 2s^2 - s(a+b+c) + ab \\ &= 2s^2 - s.2s + ab = ab.\end{aligned}$$

$$\text{Denominator} = \Delta^2.$$

$$\therefore \text{Left side} = \frac{abc}{\Delta} = 4R.$$

**Ex. 5.** If  $r_1 = r_2 + r_3 + r$ , prove that the triangle is right-angled.

From the given relation, we have

$$r_1 - r = r_2 + r_3$$

$$\text{or, } \frac{\Delta}{s-a} - \frac{\Delta}{s} = \frac{\Delta}{s-b} + \frac{\Delta}{s-c},$$

$$\text{or, } \frac{\Delta \cdot a}{s(s-a)} = \frac{\Delta \cdot a}{(s-b)(s-c)}.$$

$$\therefore s(s-a) = (s-b)(s-c).$$

$$\therefore \tan^2 \frac{1}{2}A = \frac{(s-b)(s-c)}{s(s-a)} = 1. \quad \therefore \tan \frac{1}{2}A = 1.$$

$$\therefore \frac{1}{2}A = 45^\circ. \quad \therefore A = 90^\circ.$$

**Note.** Although we got  $\tan \frac{1}{2}A = \pm 1$ , we reject the negative value because  $\frac{1}{2}A$  is an acute angle.

### Examples XIV(b)

Prove that in any triangle (Ex. 1 to 14) :—

$$1. \quad \sin A + \sin B + \sin C = \frac{s}{R}.$$

$$2. \quad \cos A + \cos B + \cos C = 1 + \frac{r}{R}.$$

$$[ \text{Use } \cos A + \cos B + \cos C = 1 + 4 \sin \frac{1}{2}A \sin \frac{1}{2}B \sin \frac{1}{2}C. ]$$

$$3. \quad \frac{b-c}{r_1} + \frac{c-a}{r_2} + \frac{a-b}{r_3} = 0.$$

$$(4.) \quad r_2 r_3 + r_3 r_1 + r_1 r_2 = s^2.$$

$$5. \quad r = R (\cos A + \cos B + \cos C - 1).$$

$$6. \quad r_1 = R (\cos B + \cos C - \cos A + 1).$$

$$[ \text{Use } \cos B + \cos C - \cos A = -1 + 4 \sin \frac{1}{2} A \cos \frac{1}{2} B \cos \frac{1}{2} C ]$$

$$7. \quad a \cos B \cos C + b \cos C \cos A + c \cos A \cos B = \frac{\Delta}{R}.$$

$$8. \quad a \cot A + b \cot B + c \cot C = 2(R + r).$$

$$\left[ a \cot A = \frac{a}{\sin A} \cos A = 2R \cos A. \quad \text{Then use Ex. 2.} \right]$$

$$9. \quad R = \frac{1}{4} \frac{(r_2 + r_3)(r_3 + r_1)(r_1 + r_2)}{r_2 r_3 + r_3 r_1 + r_1 r_2}.$$

$$10. \quad \Delta = \sqrt{rr_1 r_2 r_3} = r^2 \cot \frac{1}{2} A \cot \frac{1}{2} B \cot \frac{1}{2} C.$$

$$11. \quad \left(1 - \frac{1}{r}\right) \left(1 - \frac{1}{r_1}\right) \left(1 - \frac{1}{r_2}\right) \left(1 - \frac{1}{r_3}\right) = \frac{4R}{r^2 s^2} = \frac{16R}{r^2 (a+b+c)^2}.$$

[ A. I. 1938. ]

$$12. \quad \left(1 + \frac{1}{r_1} + \frac{1}{r_2} + \frac{1}{r_3}\right)^2 = 4 \left(1 + \frac{1}{r_1} + \frac{1}{r_2} + \frac{1}{r_3}\right).$$

$$13. \quad r_1(r_2 + r_3) \operatorname{cosec} A = r_2(r_3 + r_1) \operatorname{cosec} B \\ = r_3(r_1 + r_2) \operatorname{cosec} C.$$

$$14. \quad \frac{bc}{r_1} + \frac{ca}{r_2} + \frac{ab}{r_3} = 2R \left\{ \frac{b}{a} + \frac{c}{a} + \frac{c}{b} + \frac{a}{b} + \frac{a}{c} + \frac{b}{c} - 3 \right\}.$$

$$15. \quad \text{In a triangle } a = 13, b = 14, c = 15; \text{ find } r \text{ and } R.$$

$$16. \quad \text{If } a, b, c \text{ are in A.P., show that } r_1, r_2, r_3 \text{ are in H.P.}$$

$$17. \quad \text{If in a triangle } 3R = 4r, \text{ show that } \\ 4(\cos A + \cos B + \cos C) = 7.$$

$$18. \quad \text{If the diameter of an ex-circle be equal to the perimeter of the triangle, show that the triangle is right-angled.}$$

$$[ \text{Use } r_1 = s \tan \frac{1}{2} A. ]$$

19. If  $\left(1 - \frac{r_1}{r_2}\right)\left(1 - \frac{r_1}{r_3}\right) = 2$ , show that the triangle must be right-angled.

20. If  $8R^2 = a^2 + b^2 + c^2$ , show that the triangle is right-angled.

21. If  $S$  be the area of the in-circle and  $S_1, S_2, S_3$  the areas of the escribed circles, then

$$\frac{1}{\sqrt{S}} = \frac{1}{\sqrt{S_1}} + \frac{1}{\sqrt{S_2}} + \frac{1}{\sqrt{S_3}}.$$

22. In any triangle, prove that the area of the in-circle is to the area of the triangle as  $\pi : \cot \frac{1}{2}A \cot \frac{1}{2}B \cot \frac{1}{2}C$ .

23. If  $p_1, p_2, p_3$  are the perpendiculars from the angular points of a triangle to the opposite sides, show that

$$\frac{1}{p_1} + \frac{1}{p_2} + \frac{1}{p_3} = \frac{1}{r_1} + \frac{1}{r_2} + \frac{1}{r_3}.$$

24. If  $x, y, z$  be the lengths of the perpendiculars from the circum-centre on the sides  $BC, CA, AB$  of the triangle  $ABC$ , prove that

$$\frac{a}{x} + \frac{b}{y} + \frac{c}{z} = \frac{abc}{4xyz}.$$

25. If  $x, y, z$  are respectively equal to  $IA, IB, IC$ , and  $\alpha, \beta, \gamma$  are respectively equal to  $I_1A, I_2B, I_3C$ , show that

$$\begin{aligned} \text{(i)} \quad \frac{xyz}{abc} &= \frac{r}{s} & \text{(ii)} \quad \frac{x}{\alpha} + \frac{y}{\beta} + \frac{z}{\gamma} &= 1. \\ \text{(iii)} \quad \frac{bc}{\alpha^2} + \frac{ca}{\beta^2} + \frac{ab}{\gamma^2} &= 1. & \text{(iv)} \quad ax^2 + by^2 + cz^2 &= abc. \end{aligned}$$

[ Use Notes of Arts. 93 and 94. ]

## CHAPTER XV

### SOLUTION OF TRIANGLES

95. In a triangle, there are six parts, the three sides and the three angles. These are not independent, but are connected by the relations between the sides and angles of the triangle, which have been established in Chapter XIV. In fact, if three of the parts are given, the remaining three can, in general, be determined, and the corresponding triangle completely known. The cases that can arise are the following :

- (1) three sides may be given
- (2) three angles may be given
- (3) two sides and the included angle may be given
- (4) two angles and one side may be given
- (5) two sides and an opposite angle may be given.

We shall discuss these cases one by one.

#### 96. Three sides given.

Let the three sides  $a, b, c$  of a triangle  $ABC$  be given. Now, provided the sum of any two of these given sides is greater than the third, the triangle  $ABC$  with the three given sides can be geometrically constructed and the triangle is unique ; in other words, its angles are definite. To determine any angle  $A$  say, we may use the rigorous formula,

$$\cos A = \frac{b^2 + c^2 - a^2}{2bc},$$



and thereby determine  $\cos A$ , and then from the cosine-table find out the angle with this cosine. It is clear that the angle, being an angle of a triangle, lies between 0 and  $\pi$  and within this range an angle with a given cosine has got only one value. Thus the angle is definitely known.

Here we want to make one point clear. Though the formula used is rigorous, the cosine-table, by means of which we determine the angle with a given cosine, gives only approximate values. Now it is a principle proved in books on higher mathematics (with the aid of calculus), that *when an angle is determined by using an approximate table, the best result is obtained by using the Logarithmic tangent-table*, and an angle determined from its  $L \tan$ , using a four-figure table is more accurate than that determined by using even a seven-figure sine-table or cosine-table. If a suitable tangent formula is available therefore, we should make use of it.

Hence, for practical purposes, in this case, to determine  $A$ , we use the formula

$$\tan \frac{1}{2}A = \sqrt{\frac{(s-b)(s-c)}{s(s-a)}}.$$

where  $s = \frac{1}{2}(a+b+c)$ , which is known.

Taking logarithm, and adding 10, we get the value of  $L \tan \frac{1}{2}A$  and therefore  $A$  is known.

Similarly  $B$  and  $C$  are determined.

In case  $\tan \frac{1}{2}A$  happens to be equal to the tangent of a standard angle,  $\frac{1}{2}A$  is at once known and the use of logarithm is not wanted.

**Ex.** The sides of a triangle are 2, 3, 4. Find the greatest angle, having given

$$\log 2 = '30103, \quad \log 3 = '4771213$$

$$L \tan 55^\circ 14' = 10'1108395, \quad L \tan 52^\circ 15' = 10'1111004.$$

$$\text{Here} \quad s = \frac{2+3+4}{2} = \frac{9}{2}.$$

The greatest side 4 being denoted by 'a', the greatest angle  $A$  (which is opposite to the greatest side) is given by

$$\tan \frac{1}{2}A = \sqrt{\frac{(\frac{9}{2}-2)(\frac{9}{2}-3)}{\frac{9}{2}(\frac{9}{2}-4)}} = \sqrt{\frac{5.3}{9.1}} = \sqrt{\frac{10}{2.3}}.$$

$$\begin{aligned} \therefore L \tan \frac{1}{2}A &= 10 + \frac{1}{2}(\log 10 - \log 2 - \log 3) \\ &= 10 + \frac{1}{2}(1 - '30103 - '4771213) \\ &= 10'1109244. \end{aligned}$$

Now  $L \tan \frac{1}{2}A$  lies between  $L \tan 52^\circ 14'$  and  $L \tan 52^\circ 15'$ .

Hence,  $\frac{1}{2}A$  lies between  $52^\circ 14'$  and  $52^\circ 15'$ .

$$\text{Let} \quad \frac{1}{2}A = 52^\circ 14' x''.$$

Then diff. for  $x''$  is '0000849.

and diff. for  $1'$  i.e.  $60''$  is '0002609.

$$\text{Hence,} \quad \frac{x}{60} = \frac{849}{2609}, \quad \text{or,} \quad x = \frac{60 \times 849}{2609} = 19.5 \text{ nearly.}$$

$$\text{Hence,} \quad \frac{1}{2}A = 52^\circ 14' 19''.5,$$

$$\text{or,} \quad A = 104^\circ 28' 39'' \text{ nearly.}$$

### 97. Three angles given.

In this case the triangle cannot be solved, for there are innumerable triangles with the same three angles. All

these triangles, being equiangular, are similar, and the ratio of their sides can be determined from the formula,

$$\frac{a}{\sin A} = \frac{b}{\sin B} = \frac{c}{\sin C},$$

$$\text{or, } a : b : c = \sin A : \sin B : \sin C.$$

**Ex.** *The angles of a triangle are in the ratio 2 : 3 : 7. Prove that the sides are in the ratio of  $\sqrt{2} : 2 : (\sqrt{3} + 1)$ .*

The angles being in the ratio of 2 : 3 : 7, and their sum being  $180^\circ$ , the angles are evidently  $30^\circ$ ,  $45^\circ$  and  $105^\circ$  respectively. Hence the ratio of the sides will be

$$\sin 30^\circ : \sin 45^\circ : \sin 105^\circ$$

$$\text{i.e., } \frac{1}{2} : \frac{1}{\sqrt{2}} : \frac{\sqrt{3}+1}{2\sqrt{2}}$$

$$\text{or, } \sqrt{2} : 2 : (\sqrt{3} + 1).$$

### Examples XV(a)

1. The sides of a triangle are 24, 22, 14 ; find the least angle, given  $L \tan 17^\circ 33' = 9.500042$ , diff. for  $1' = 439$ .

2. The sides of a triangle are 50, 36 and 28 ; find the greatest angle, having given

$$\log 19 = 1.2787536, \quad \log 29 = 1.4623980$$

$$L \tan 51^\circ 0' = 10.0916308, \quad L \tan 51^\circ 1' = 10.0918891.$$

3. The sides of a triangle are 9, 10 and 11 ; find the angle opposite to the side 10, given

$$L \tan 29^\circ 30' = 9.7526420, \quad L \tan 29^\circ 29' = 9.7523472$$

$$\log 2 = .30103. \quad [C. U. 1943.]$$

4. The sides of a triangle are 2, 3, 4. Find all the angles correctly to degrees and minutes by the help of mathematical tables.

5. (i) The sides of a triangle are 15, 19, 24 ; find the greatest angle of the triangle.

$$\text{Given } \log 5.7 = .75587, L \cos 88^\circ 59' = 8.24903$$

$$\text{diff. for } 1' = 718. \quad [C. U. 1936.]$$

(ii) Find the greatest angle in degrees, minutes and seconds in a triangle whose sides are 5, 6, 7, having given

$$\log 6 = .7781513$$

$$L \cos 39^\circ 14' = 9.8890644, \text{ diff. for } 60'' = .0001032.$$

6. (i) The sides of a triangle are 7, 8, 9 ; solve the triangle. [ C. U. 1938. ]

(ii) If  $a = 32$ ,  $b = 40$ ,  $c = 66$ , determine the greatest angle. [ C. U. 1945. ]

[ Use Mathematical Tables ]

7. Given  $a = \sqrt{6}$ ,  $b = 2$ ,  $c = \sqrt{3} - 1$  ; solve the triangle.

8. Given  $a = 2$ ,  $b = \sqrt{2}$ ,  $c = \sqrt{3} + 1$  ; solve the triangle.

9. If  $a = 7$ ,  $b = 5$ ,  $c = 8$ , solve the triangle.

$$\text{Given } \cos 38^\circ 11' = \frac{1}{4}.$$

10. If  $a = 3 + \sqrt{3}$ ,  $b = 2\sqrt{3}$ ,  $c = \sqrt{3}$ , solve the triangle.

11. The angles of a triangle are  $105^\circ$ ,  $60^\circ$  and  $15^\circ$  ; find the ratio of the sides.

12. If  $A = 45^\circ$ ,  $B = 60^\circ$ , show that  $c : a = \sqrt{3} + 1 : 2$ .

13. The angles of a triangle are as  $1 : 2 : 7$  ; find the ratio of the greatest side to the least side.

14. If  $\cos A = \frac{1}{2}$ ,  $\cos B = \frac{2}{3}$ , find  $a : b : c$ .

15. If the angles adjacent to the base of a triangle are  $22\frac{1}{2}^\circ$  and  $112\frac{1}{2}^\circ$ , show that the altitude is half the base.

16. If the sides of a triangle are 4, 5, 6, show that the greatest angle is double the least.

**98. Two sides and the included angle given.**

Let the two sides  $b, c$  and the included angle  $A$  of a triangle  $ABC$  be given. It is easy to construct the triangle geometrically, and there will be only one definite triangle with the given parts. To find the other angles  $B$  and  $C$ , we notice that

$$B + C = 180^\circ - A,$$

$$\text{i.e., } \frac{B + C}{2} = 90^\circ - \frac{A}{2}.$$

Again,

$$\tan \frac{B - C}{2} = \frac{b - c}{b + c} \cot \frac{A}{2}.$$

$$\begin{aligned} L \tan \frac{B - C}{2} &= 10 + \log \left( \frac{b - c}{b + c} \cot \frac{A}{2} \right) \\ &= \log \left( \frac{b - c}{b + c} \right) + L \cot \frac{A}{2} \end{aligned}$$

$b, c$  and  $A$  being given, the right-hand side is known and thus  $L \tan \frac{B - C}{2}$  is known, whence  $\frac{B - C}{2}$  is known.

Now  $\frac{B + C}{2}$  and  $\frac{B - C}{2}$  being both known, by addition and subtraction, we get  $B$  and  $C$  respectively.

The reason of using tangent formula to determine  $\frac{B - C}{2}$  is already explained in Art. 96.

When  $B$  and  $C$  are known, the third side  $a$  is easily obtained from

$$\frac{a}{\sin A} = \frac{b}{\sin B} \quad \text{or, } = \frac{c}{\sin C}.$$

**Ex.** In a triangle,  $b = 2.25$ ,  $c = 1.75$ ,  $A = 54^\circ$ , find  $B$  and  $C$ , having given

$$\begin{aligned} \log 2 &= .301030, & L \tan 63^\circ &= 10.292834 \\ L \tan 13^\circ 47' &= 9.389724, & L \tan 13^\circ 48' &= 9.390270 \end{aligned}$$

[C. U. 1931.]

Here,

$$\frac{B+C}{2} = 90^\circ - \frac{A}{2} = 90^\circ - 27^\circ = 63^\circ. \quad \dots (i)$$

Again,

$$\begin{aligned} \tan \frac{B-C}{2} &= \frac{b-c}{b+c} \cot \frac{A}{2} = \frac{5}{4} \cot 27^\circ \\ &= \frac{1}{8} \tan 63^\circ. \end{aligned}$$

$$\begin{aligned} \therefore L \tan \frac{B-C}{2} &= L \tan 63^\circ - 3 \log 2 \\ &= 10.292834 - .903090 \\ &= 9.389744. \end{aligned}$$

Now  $L \tan 13^\circ 47' = 9.389724$

and  $L \tan 13^\circ 48' = 9.390270.$

Hence,  $\frac{B-C}{2}$  being  $13^\circ 47' x''$

we get, diff. for  $x'' = .000020$

and diff. for  $1' \text{ i.e., } 60'' = .000546.$

$$\therefore \frac{x}{60} = \frac{20}{546}, \text{ or, } x = \frac{20 \times 60}{546} = 2.2 \text{ nearly.}$$

Hence,  $\frac{B-C}{2} = 13^\circ 47' 2'' 2 \text{ nearly.}$

Combining with (i),  $B = 76^\circ 47' 2'' 2$  and  $C = 49^\circ 12' 57'' 8.$

### 99. Two angles and a side given.

Let any side  $a$  of a triangle  $ABC$ , and any two of its angles be given. The sum of the three angles being  $180^\circ$ , the third angle is also known. To find the other two sides  $b$  and  $c$ , we use the formula

$$\frac{a}{\sin A} = \frac{b}{\sin B} = \frac{c}{\sin C}.$$

**Ex.** In a triangle  $ABC$ ,  $A = 38^\circ 20'$ ,  $B = 45^\circ$  and  $b = 64 \text{ ft.}$   
Find  $c$ , having given  $\log 2 = .30103$ ,  $L \sin 83^\circ 20' = 9.99705$   
and  $\log .089896 = \bar{2}.95374.$

Here,

$$\begin{aligned} C &= 180^\circ - (A + B) \\ &= 180^\circ - 83^\circ 20'. \end{aligned}$$

Now,

$$\frac{c}{\sin C} = \frac{b}{\sin B},$$

$$\text{or, } \frac{c}{\sin (180^\circ - 83^\circ 20')} = \frac{64}{\sin 45^\circ} = \frac{64}{1/\sqrt{2}} = 64\sqrt{2}.$$

$$\therefore c = 2^{\frac{1}{2}} \sin 83^\circ 20'.$$

$$\begin{aligned} \therefore \log c &= \frac{1}{2} \log 2 + L \sin 83^\circ 20' - 10 \\ &= \frac{1}{2} (.30103) + 9.99705 - 10 = 1.95374. \end{aligned}$$

Thus  $\log c$  has the same mantissa as  $\log .089896$ , but has its characteristic 1. Hence,  $c = 89.896$  feet.

### Examples XV(b)

- Two sides of a triangle are 3 and 5 feet and the included angle is  $120^\circ$ ; find the other angles, having given

$$\log 4.8 = .6812412$$

$$L \tan 8^\circ 12' = 9.1586706, \text{ diff. for } 60'' = 8940.$$

[C. U. 1949.]

- If  $b = 1300$ ,  $c = 1400$  and  $A = 60^\circ$ , find  $B$  and  $C$ .

$$\text{Given } \log 3 = .4771213,$$

$$L \tan 3^\circ 40' = 8.8067422, \text{ diff. for } 10'' = 3306.$$

- If  $a = 21$ ,  $b = 11$ ,  $C = 34^\circ 42' 30''$ , find  $A$  and  $B$ .

$$\text{Given } \log 2 = .30103,$$

$$\text{and } L \tan 72^\circ 38' 45'' = 10.50515.$$

- If the sides  $a$  and  $b$  are in the ratio  $7 : 3$  and the included angle  $C$  is  $60^\circ$ , find  $A$  and  $B$ , given

$$\log 2 = .3010300$$

$$\log 3 = .4771213$$

$$L \tan 34^\circ 42' = 9.8403776, \text{ diff. for } 1' = 2699.$$

5. Two sides of a plane triangle are 14 and 11 and the included angle is  $60^\circ$ . Find the remaining angles, having given  $L \tan 11^\circ 44' = 9.3174299$ ,  $L \tan 11^\circ 45' = 9.3180640$ .

[ C. U. 1922. ]

6. (i) Two sides of a triangle are 80 and 100 ft. and the included angle is  $60^\circ$ . Find the other angles.

[ C. U. 1946. ]

(ii) If  $a = 5$ ,  $b = 3$ ,  $C = 70^\circ 30'$ , find the remaining angles.

(iii) If  $a = 39.9$ ,  $b = 43.2$ ,  $C = 38^\circ 14'$ , solve the triangle.

[ Use Mathematical Tables ]

7. (i) In a plane triangle,  $b = 540$ ,  $c = 420$  and  $A = 52^\circ 6'$ ; find  $B$  and  $C$ , having given

$$L \tan 26^\circ 3' = 9.6891430,$$

$$L \tan 14^\circ 20' = 9.4074189, L \tan 14^\circ 21' = 9.4079453.$$

[ C. U. 1934. ]

(ii) Given  $a = 70$ ,  $b = 35$ ,  $C = 36^\circ 52' 12''$ ,  $\log 3 = 0.4771213$ ,  $L \cot 18^\circ 26' 6'' = 10.4771213$ . Calculate the other two angles  $A$  and  $B$ .

[ C. U. 1935, '37. ]

8. If  $a = 2\sqrt{6}$ ,  $c = 6 - 2\sqrt{3}$ ,  $B = 75^\circ$ , solve the triangle.

9. Two sides of a triangle are  $\sqrt{3} + 1$  and  $\sqrt{3} - 1$  and the included angle is  $60^\circ$ ; solve the triangle.

10. (i) If  $a = 2$ ,  $b = 1 + \sqrt{3}$ ,  $C = 60^\circ$ , solve the triangle.

(ii) If  $a = 2$ ,  $b = 4$ ,  $C = 60^\circ$ , find  $A$  and  $B$ .

11. If  $a = 19$ ,  $B = 52^\circ 28'$  and  $C = 93^\circ 40'$ , find  $b$ , having given  $\log 27038 = 4.4319746$ ;  $\log 19 = 1.2787536$ ;

$$\log 27037 = 4.4319585;$$

$$L \sin 52^\circ 28' = 9.8992727, L \sin 33^\circ 52' = 9.7460595.$$



12. If  $B = 45^\circ$ ,  $C = 10^\circ$  and  $a = 200$  ft., find  $b$ , having  
given  $\log 2 = .30103$ ,  $L \sin 55^\circ = 9.9133645$

$$\log 1726.4 = 3.2371414, \quad \log 1726.5 = 3.2371666.$$

[ *C. U. 1947.* ]

13. If  $A = 41^\circ 13' 22''$ ,  $B = 71^\circ 19' 5''$ , and  $a = 55$ , find  $b$ ,  
given  $\log 55 = 1.7403627$ ,  $\log 79063 = 4.8979775$

$$L \sin 41^\circ 13' 22'' = 9.8188779$$

$$L \sin 71^\circ 19' 5'' = 9.9764927.$$

14. (i) If  $B = 70^\circ 30'$ ,  $C = 78^\circ 10'$ ,  $a = 102$ , solve the triangle.

(ii) If  $a = 39$ ,  $A = 81^\circ 35'$ ,  $B = 27^\circ 55'$ , solve the triangle.

(iii) If  $A = 37^\circ 15'$ ,  $B = 72^\circ 5'$ ,  $a = 75.2$ , find  $b$  and  $c$ .

[ *Mathematical tables should be used* ]

15. If  $A = 75^\circ$ ,  $B = 30^\circ$ ,  $b = \sqrt{8}$ , solve the triangle.

16. If  $A = 30^\circ$ ,  $B = 45^\circ$ ,  $b = 2$ , solve the triangle.

17. In a triangle in which each base angle is double of the third angle, the base is 2 ; solve the triangle.

18. Given  $a = \sqrt{57}$ ,  $A = 60^\circ$ ,  $\Delta = 2\sqrt{3}$ , find  $b$  and  $c$ .

#### 100. Two sides and an opposite angle given.

Let the two sides  $b$  and  $c$  in a triangle  $ABC$ , and the angle  $B$  opposite to the side  $b$  be given.

In this case, we get the angle  $C$  from the formula,

$$\frac{\sin C}{c} = \frac{\sin B}{b} \quad \text{or,} \quad \sin C = \frac{c \sin B}{b}.$$

Now three cases may arise, namely,

- (i)  $c \sin B > b$ . In this case  $\sin C$  is greater than 1, and so  $C$  cannot be obtained. In fact in this case no triangle is possible.

- (ii)  $c \sin B = b$ . Here  $\sin C$  becomes 1 and therefore  $C = 90^\circ$ . Thus  $A = 90^\circ - B$ . We thus get a right-angled triangle with right angle at  $C$ , and the side  $a$  will be obtained from

$$c^2 = a^2 + b^2, \text{ or, } a = \sqrt{c^2 - b^2}.$$

- (iii)  $c \sin B < b$ . In this case  $\sin C$  is less than 1, and hence  $C$  can be determined. Now sines of supplementary angles are known to be equal, and an angle of a triangle may be acute or obtuse. We therefore get two supplementary values of  $C$  having the same value for  $\sin C$ . Three sub-cases now arise :

*Sub-case A.* If of the two given sides,  $b > c$ , then  $B > C$ , and therefore the obtuse value of  $C$  becomes inadmissible, for otherwise  $B$  is also obtuse and two angles  $B$  and  $C$  of the triangle become both obtuse. Thus the only admissible solution is the acute value of  $C$ . Now  $B$  and  $C$  being both known,  $A$  is obtained from  $A + B + C = 180^\circ$ . The side  $a$  will be known from

$$\frac{a}{\sin A} = \frac{b}{\sin B} \text{ or, } \frac{a}{\sin C}.$$

Thus the triangle is uniquely solved.

*Sub-case B.* If  $b = c$ , then  $B = C$ , and here also the obtuse value of  $C$  is inadmissible ; with the acute value of  $C$  the triangle is uniquely solved exactly as in the above case.

*Sub-case C.* If  $b < c$ , then  $B < C$ , so that  $C$  may be either acute or obtuse. Both the supplementary values of  $C$  being admissible now, there will be two possible triangles with the three given parts  $b, c, B$ . Corresponding to each value of  $C$ , the value of  $A$  is determined from  $A + B + C = 180^\circ$ , and then  $a$  is obtained from the formula,

$$\frac{a}{\sin A} = \frac{b}{\sin B} \text{ or, } \frac{a}{\sin C}.$$

As there are two solutions of the triangle in this case, each equally admissible, this sub-case in the solution of a

triangle in which  $b, c, B$  are given and  $b > c \sin B$  but  $< c$ , is referred to as the **Ambiguous Case** in the solution of triangles.

We may sum up the result as follows :

When in a triangle,  $b, c, B$  are given,

- (i) if  $b < c \sin B$ , no triangle is possible ;
- (ii) if  $b = c \sin B$ , we get a definite right-angled triangle as solution ;
- (iii) if  $b > c$  and therefore necessarily  $> c \sin B$ , we get one definite solution having  $C$  acute ;
- (iv) if  $b = c$  and therefore necessarily  $> c \sin B$ , we get one definite solution having  $C$  acute.
- (v) if  $b > c \sin B$  but  $< c$ , there are two solutions, and this case is the *ambiguous case*.

### 101. Geometrical treatment of the Ambiguous Case.

To make the ideas clear, we proceed to construct geometrically the triangle in which two sides and an opposite angle, viz.,  $b, c$  and  $B$  are given.




Fig. (i)




Fig. (ii)




Fig. (iii)




Fig. (iv)

Let  $ABX$  be the given angle  $B$ , and along one arm of it, take  $BA = c$ . Let  $AN$  be the perpendicular from  $A$  on  $BX$ . Then  $\frac{AN}{AB} = \sin B$ , so that  $AN = AB \sin B = c \sin B$ .

With centre  $A$  and radius  $b$  draw a circle.

*Case (i).* If  $b < c \sin B$ , i.e.,  $< AN$ , the circle does not meet the side  $BX$  at all and no triangle is therefore obtained. See fig. (i).

*Case (ii).* If  $b = c \sin B$ , i.e.,  $= AN$ , the circle touches the side  $BX$  at  $C$  coincident with  $N$ , as in fig. (ii). Hence a right-angled triangle is formed, in which the sides  $AB$ ,  $AC$  and the angle  $B$  have the given values  $c$ ,  $b$ ,  $B$ . Thus  $ABC$  is the required triangle.

*Case (iii).* If  $b > c$ , i.e.,  $> AB$ , the circle cuts  $BX$  at two points  $C$  and  $C'$  on opposite sides of  $B$  as in fig. (iii). The triangle  $ABC'$ , though it has the sides  $AB$ ,  $AC$  equal to the given quantities  $c$  and  $b$ , has the angle  $B$  not equal to the given angle, but equal to its supplement. Hence it is not the solution required. In this case the triangle  $ABC$  is the only solution.

*Case (iv).* If  $b = c$ , i.e.,  $= AB$ , the point  $C'$  of the above case coincides with  $B$ , and only one triangle  $ABC$  is obtained as the required solution.

*Case (v).* If  $b > c \sin B$ , i.e.,  $> AN$  but less than  $c$  (or  $AB$ ), the circle cuts  $BX$  at two points  $C_1$  and  $C_2$  on the same side of  $B$  as in fig. (iv). Both the triangles  $ABC_1$  and  $ABC_2$  have the same three given parts and both are possible solutions. This is therefore the *Ambiguous* case.

**Note.** By considering the equation

$$b^2 = c^2 + a^2 - 2ca \cos B$$

in which  $b, c, B$  are given, we may first of all determine  $a$ , instead of trying to determine  $C$ .

Considering the equation as a quadratic in  $a$ , viz.,

$$a^2 - 2c \cos B \cdot a + c^2 - b^2 = 0,$$

and by solving it, we get

$$a = c \cos B \pm \sqrt{b^2 - c^2 \sin^2 B}.$$

(i) If  $b < c \sin B$ ,  $b^2 - c^2 \sin^2 B$  is negative and thus the two values of  $a$  are imaginary. (No solution)

(ii) If  $b = c \sin B$ ,  $b^2 - c^2 \sin^2 B = 0$  and thus the two values of  $a$  are real and coincident.

(one solution ; one triangle right-angled at  $C$ , since  $b = c \sin B$ )

(ii) If  $b > c \sin B$ ,  $b^2 - c^2 \sin^2 B$  is positive, so two values of  $a$  are real and distinct, but they are not always admissible.

(a) When  $b > c$ , { i.e.,  $b^2 > c^2 (\sin^2 B + \cos^2 B)$  },  $b^2 - c^2 \sin^2 B > c^2 \cos^2 B$ , i.e.,  $\sqrt{b^2 - c^2 \sin^2 B} > c \cos B$  and hence one value of  $a$  is positive and the other negative. (one solution)

(b) When  $b = c$ ,  $b^2 - c^2 \sin^2 B = c^2 - c^2 \sin^2 B = c^2 \cos^2 B$  and hence one value of  $a$  is zero. (one solution)

(c) When  $b < c$ , i.e.,  $b^2 < c^2 (\sin^2 B + \cos^2 B)$ ,  $b^2 - c^2 \sin^2 B < c^2 \cos^2 B$ , i.e.,  $\sqrt{b^2 - c^2 \sin^2 B} < c \cos B$ .

So both the values of  $a$  are real and positive. (two solutions)

This is known as the *algebraical discussion* of the ambiguous case.

An example illustrating the algebraic method is added below.

**Ex. 1.** In a triangle,  $b = 15$  ft.,  $c = 10$  ft.,  $B = 60^\circ$ . Find  $a$  and  $A$  having given  $\sin 84^\circ 44' = .99578$ .

We have  $b^2 = c^2 + a^2 - 2ca \cos B$ , giving here

$$225 = 100 + a^2 - 20a \cos 60^\circ ;$$

$$\text{or, } a^2 - 10a - 125 = 0 \text{ whence}$$

$$a = 5 \pm 5\sqrt{6}.$$

Rejecting the negative value for  $a$  as inadmissible, the only possible value of  $a = 5(\sqrt{6} + 1)$  ft. There is thus one solution and there is no ambiguity. In fact this is case (iii) of the previous article.

$$\begin{aligned}\text{Again, } \sin A &= \frac{a}{b} \sin B = \frac{5(\sqrt{6} + 1)}{15} \cdot \frac{\sqrt{3}}{2} = \frac{3\sqrt{2} + \sqrt{3}}{6} \\ &= \frac{3 \times 1.41421 \dots + 1.73205}{6} = .99578 \dots\end{aligned}$$

$$\text{so } A = 84^\circ 44'.$$

**Ex. 2.** In a triangle,  $a = 73.4$ ,  $b = 64.9$  and  $B = 48^\circ 13' 25''$ ; find  $A$  having given

$$\log 734 = 2.8656961, \quad \log 649 = 2.8122447$$

$$L \sin 48^\circ 13' 25'' = 9.8725936$$

$$L \sin 57^\circ 30' = 9.9260292 \quad (\text{diff. for } 1' = 804)$$

Is the case ambiguous?

Here

$$\sin A = \frac{a \sin B}{b} = \frac{734}{649} \sin 48^\circ 13' 25''.$$

$$\begin{aligned}\therefore L \sin A &= \log 734 - \log 649 + L \sin 48^\circ 13' 25'' \\ &= 2.8656961 - 2.8122447 + 9.8725936 \\ &= 9.9260450.\end{aligned}$$

Now diff. of this from  $L \sin 57^\circ 30' = 158$  (i.e., .0000158) and diff. for  $1'$  (or  $60''$ ) = 804 (i.e., .0000804).

Hence,  $A = 57^\circ 30' x''$  where

$$\frac{x}{60} = \frac{158}{804} \text{ whence } x = 11.8 \text{ nearly.}$$

Thus  $A = 57^\circ 30' 11.8''$  or its supplement  $122^\circ 29' 48.2''$  which has also the same sine, and so the same  $L$  sine.

Now in this case  $a > b$  and so  $A > B$  and thus both values of  $A$  are admissible. The case, is therefore, the *ambiguous case* and will have two solutions.

## Examples XV(c)

1. Given (i)  $A = 30^\circ$ ,  $a = 6$ ,  $b = 4$ .  
 (ii)  $A = 60^\circ$ ,  $a = 7$ ,  $b = 8$ .  
 (iii)  $A = 45^\circ$ ,  $a = 2$ ,  $b = 8$ .  
 (iv)  $A = 30^\circ$ ,  $a = 3$ ,  $b = 6$ .

Find in which case the solution is ambiguous, in which case there is one solution, and in which case there is no solution.

2. (i) If  $b = 2$ ,  $c = \sqrt{3} + 1$  and  $B = 45^\circ$ , solve the triangle.

(ii) If  $a = 3$ ,  $b = 3\sqrt{3}$ ,  $A = 30^\circ$ , find  $B$ .

3. If  $a = 2$ ,  $b = \sqrt{6}$ ,  $B = 60^\circ$ , solve the triangle.

4. If  $a = 2$ ,  $b = 5$ ,  $A = 30^\circ$ , solve the triangle.

5. If  $b$ ,  $c$ ,  $B$  are given and if  $b < c$ , show that

$$(a_1 - a_2)^2 + (a_1 + a_2)^2 \tan^2 B = 4b^2$$

$a_1$  and  $a_2$  being the two possible values of  $a$ .

6. In the ambiguous case, given  $a$ ,  $b$  and  $A$ , prove that the difference between the two values of  $c$  is

$$2\sqrt{a^2 - b^2 \sin^2 A}.$$

7. If  $a$ ,  $b$ ,  $A$  are given and if  $c_1$ ,  $c_2$  are the values of the third side in the ambiguous case, prove that if  $c_1 > c_2$

$$(i) \quad c_1 - c_2 = 2a \cos B. \quad [B. H. U. I. 1928.]$$

$$(ii) \quad c_1^2 + c_2^2 - 2c_1 c_2 \cos 2A = 4a^2 \cos^2 A.$$

[B. H. U. I. 1935; Pat. I. 1936.]

$$(iii) \quad \cos \frac{C_1 - C_2}{2} = \frac{b \sin A}{a}. \quad [A. I. 1941.]$$

8. If  $b = 16$ ,  $c = 25$  and  $B = 33^\circ 15'$ , find the other angles; given

$$L \sin 33^\circ 15' = 9.7390129, \quad \log 2 = .30103,$$

$$L \sin 58^\circ 57' = 9.9328376 \quad L \sin 58^\circ 56' = 9.9327616.$$

9. If  $a=5$ ,  $b=4$ ,  $A=45^\circ$ , find  $B$  and  $C$ ; given  
 $\log 2 = .30103$ ,  $L \sin 34^\circ 27' = 9.75257$ .

10. If  $a=30$ ,  $b=300$ , find  $A$  in order that  $B$  may be a right angle, having given that

$$L \sin 5^\circ 44' = 8.9995595, \text{ diff. for } 1' = 12565.$$

11. If  $a=16$ ,  $c=25$  and  $C=60^\circ$ , find the other angles.  
 Given

$$\log 2 = .30103, \quad \log 3 = .4771213$$

$$L \sin 33^\circ 39' = 9.7436024, \text{ diff. for } 1' = 1897.$$

12. If  $b=165$ ,  $c=258$ , and  $B=35^\circ 10'$ , find the angles  $A$  and  $C$ .

$$\text{Given } \log 1.65 = .21749, \quad \log 2.58 = .41162$$

$$L \sin 35^\circ 10' = 9.76039, \quad L \sin 64^\circ 14' = 9.95452.$$

13. If  $2b=3a$  and  $\tan^2 A = \frac{8}{5}$ , prove that there are two values of the third side, one of which is double the other.

14. If  $A_1$ ,  $B_1$  and  $A_2$ ,  $B_2$  are the angles of the two triangles in the ambiguous case, then

$$\frac{\sin A_1}{\sin B_1} + \frac{\sin A_2}{\sin B_2} = 2 \cos C.$$

15. Show that in the case that admits of two solutions the two values of  $C$  satisfy the equation

$$\frac{(a+b)^2}{1+\cos C} + \frac{(b-a)^2}{1-\cos C} = \frac{2a^2}{\sin^2 A} \quad [B. H. U. I. 1942.]$$

16. If  $\log b + 10 = \log c + L \sin B$ , can the triangle be ambiguous?



## Miscellaneous Examples II

In any triangle  $ABC$ , prove that (Ex. 1 to 8):—

$$1. \quad \frac{1}{a} \cos A + \frac{1}{b} \cos B + \frac{1}{c} \cos C = \frac{a^2 + b^2 + c^2}{2abc}.$$

$$2. \quad (b^2 + c^2 - a^2) \tan A = (c^2 + a^2 - b^2) \tan B \\ = (a^2 + b^2 - c^2) \tan C.$$

$$3. \quad b^2 + c^2 - 2bc \cos (A + 60^\circ) = c^2 + a^2 - 2ca \cos (B + 60^\circ) \\ = a^2 + b^2 - 2ab \cos (C + 60^\circ).$$

$$4. \quad (\cot \tfrac{1}{2}A - \tan \tfrac{1}{2}B - \tan \tfrac{1}{2}C)^{\frac{1}{2}} \\ + (\cot \tfrac{1}{2}B - \tan \tfrac{1}{2}C - \tan \tfrac{1}{2}A)^{\frac{1}{2}} + (\cot \tfrac{1}{2}C - \tan \tfrac{1}{2}A - \tan \tfrac{1}{2}B)^{\frac{1}{2}} \\ = (\cot \tfrac{1}{2}A + \cot \tfrac{1}{2}B + \cot \tfrac{1}{2}C)^{\frac{1}{2}}.$$

$$5. \quad a \sin (B - C) \cos (B + C - A) + b \sin (C - A) \\ \times \cos (C + A - B) + c \sin (A - B) \cos (A + B - C) = 0.$$

$$6. \quad \frac{a \sin A + b \sin B + c \sin C}{a \cos A + b \cos B + c \cos C} = \frac{R}{abc} (a^2 + b^2 + c^2).$$

$$7. \quad (b + c - 2a) \sin \tfrac{1}{2}A \sin \tfrac{1}{2}(B - C) \\ + (c + a - 2b) \sin \tfrac{1}{2}B \sin \tfrac{1}{2}(C - A) \\ + (a + b - 2c) \sin \tfrac{1}{2}C \sin \tfrac{1}{2}(A - B) = 0.$$

$$8. \quad a \cos A \cos 2A + b \cos B \cos 2B + c \cos C \cos 2C \\ + 4 \cos A \cos B \cos C (a \cos A + b \cos B + c \cos C) = 0.$$

$$9. \quad \text{If in a triangle, } a^2, b^2, c^2 \text{ are in A.P., show that} \\ \tan A, \tan B, \tan C \text{ are in H. P.}$$

$$10. \quad \text{If in a triangle, } \sin A, \sin B, \sin C \text{ are in H. P.,} \\ \text{show that } 1 - \cos A, 1 - \cos B, 1 - \cos C \text{ are in H.P.}$$

11. Determine the triangle whose sides are three consecutive terms in the series of natural numbers and whose largest angle is double the least.

12. If in a triangle,  $\cos 3A + \cos 3B + \cos 3C = 1$ , show that one angle must be  $120^\circ$ .

13. If in a triangle,  $\sin A, \sin B, \sin C$  be in A.P., show that  $\tan \frac{1}{2}A \tan \frac{1}{2}C = \frac{1}{3}$ .

14. If  $a=5, b=7$  and  $A=30^\circ$ , find  $B$  in degrees and minutes, having given

$$\sin 44^\circ = 0.6947, \sin 45^\circ = 0.7071.$$

15. In the *ambiguous case*, the area of one of the triangles is  $n$  times that of the other; show that if  $b$  be the greater of the given sides and  $c$  the less,  $\frac{b}{c}$  is less than  $\frac{n+1}{n-1}$ .

16. In the *ambiguous case*, show that the circum-circles of the two triangles are equal.

17. Prove that

$$(i) \tan^{-1} \left( \frac{x \cos \phi}{1 - x \sin \phi} \right) - \tan^{-1} \left( \frac{x - \sin \phi}{\cos \phi} \right) = \phi.$$

$$(ii) \tan^{-1} \frac{t_1 - t_2}{1 + t_1 t_2} + \tan^{-1} \frac{t_2 - t_3}{1 + t_2 t_3} + \dots + \tan^{-1} \frac{t_{n-1} - t_n}{1 + t_{n-1} t_n} \\ = \tan^{-1} t_1 - \tan^{-1} t_n.$$

18. If the sum of four angles be  $180^\circ$ , prove that the sum of the products of their cosines taken two and two together is equal to the sum of the products of their sines taken similarly.

$$19. \text{ Prove that } \cos^2 A + \cos^2 \left( A + \frac{\pi}{3} \right) + \cos^2 \left( A - \frac{\pi}{3} \right) = \frac{3}{2}.$$

[ C. U. 1933. ]

20. In a triangle  $ABC$ , if  $\tan \frac{A}{2}, \tan \frac{B}{2}, \tan \frac{C}{2}$  be in Arithmetical progression, then  $\cos A, \cos B, \cos C$  are also in Arithmetical progression.

21. Give in general terms the solutions of the following equation :

$$\tan(x+b) \tan(x+c) + \tan(x+c) \tan(x+a) + \tan(x+a) \tan(x+b) = 1.$$

22. If  $A+B+C=180^\circ$ , prove that

$$\begin{aligned} \left(1 + \tan \frac{A}{4}\right) \left(1 + \tan \frac{B}{4}\right) \left(1 + \tan \frac{C}{4}\right) \\ = 2 \left(1 + \tan \frac{A}{4} \tan \frac{B}{4} \tan \frac{C}{4}\right). \end{aligned}$$

23. Prove that

$$\begin{aligned} \sin^2 x + \sin^2 y + \sin^2 z + \sin^2(x+y+z) \\ = 2 - 2 \cos(x+y) \cos(y+z) \cos(z+x). \end{aligned}$$

24. Solve the following equation :

$$\tan x + \tan\left(x + \frac{\pi}{3}\right) + \tan\left(x + \frac{2\pi}{3}\right) = 3.$$

[ Left side reduces to  $3 \tan 3x$ . ]

25. Prove that in a triangle  $ABC$ ,

$$\Delta = \frac{(a+b+c)^2}{4 \left( \cot \frac{A}{2} + \cot \frac{B}{2} + \cot \frac{C}{2} \right)}.$$

26. Prove that

$$\log \sin 8x = 3 \log 2 + \log \sin x + \log \cos x + \log \cos 2x + \log \cos 4x.$$

27. Show that in any triangle  $ABC$ ,

$$\log \tan \frac{A}{2} = \frac{1}{2} [\log(s-b) + \log(s-c) - \log s - \log(s-a)].$$

28. Prove that (i)  $x^{\log y} = y^{\log x}$ .

$$(ii) x^{\log y - \log z} \times y^{\log z - \log x} \times z^{\log x - \log y} = 1.$$


29. In any right-angled triangle  $ABC$ ,  $C$  being the right-angle, show that  $R+r = \frac{1}{2}(a+b)$ .

30. Show how to solve a triangle having given the three perpendiculars from the vertices on the opposite sides.

# CHAPTER XVI

## GRAPHS OF TRIGONOMETRICAL FUNCTIONS

**102. Changes in the Trigonometrical ratios of an angle as the angle increases from  $0^\circ$  to  $360^\circ$ .**



Suppose an angle traced out by a revolving line starting from  $OX$ , changes gradually from  $0^\circ$  to  $360^\circ$ .

Take a circle with centre  $O$  of any radius. It is clear that in determining the trigonometrical ratios of an angle  $XOP_1$  in its different positions, we can keep the hypotenuse  $OP_1$  always the same, equal to the radius of the circle.

### (i) Changes in sine.

When the angle  $N_1OP_1$  ( $= \theta$  say) is zero, its sine is zero. As the angle increases from  $0^\circ$  to  $90^\circ$ , the hypotenuse  $OP_1$  remaining the same, the opposite side  $P_1N_1$  is positive and gradually increases, as is evident by comparing the triangles  $N_1OP_1$  and  $N_2OP_2$ .

Hence,  $\sin \theta = \frac{P_1 N_1}{OP_1}$  gradually increases, until when  $\theta = 90^\circ$ ,  $P_2 N_2$  and  $OP_2$  both coincide with  $OY$  and  $\sin \theta$  attains its greatest value 1.

As  $\theta$  still further increases, from  $90^\circ$  to  $180^\circ$ , the hypotenuse  $OP_3$  retains the same value, but  $P_3 N_3$  remaining positive, now gradually diminishes from  $OY$  to zero, and so  $\sin \theta$  diminishes from 1 to 0. In the third quadrant, as  $\theta$  increases from  $180^\circ$  to  $270^\circ$ ,  $P_4 N_4$  is negative and numerically increases from zero to  $OY'$ , the hypotenuse remaining always positive and unaltered.  $\sin \theta$  is therefore negative and numerically increases from 0 to -1; in other words, it diminishes gradually from 0 to -1. In the fourth quadrant, as  $\theta$  increases from  $270^\circ$  to  $360^\circ$ ,  $P_5 N_5$  remaining negative numerically diminishes from  $OY'$  to 0, and  $\sin \theta$  therefore remaining negative numerically diminishes from -1 to 0; in other words, it increases from -1 to 0. The results are therefore as follows :

In the *first* quadrant, as  $\theta$  increases from  $0^\circ$  to  $90^\circ$ ,

*$\sin \theta$  increases from 0 to 1.*

In the *second* quadrant, as  $\theta$  increases from  $90^\circ$  to  $180^\circ$ ,

*$\sin \theta$  diminishes from 1 to 0.*

In the *third* quadrant, as  $\theta$  increases from  $180^\circ$  to  $270^\circ$ ,

*$\sin \theta$  diminishes from 0 to -1.*

In the *fourth* quadrant, as  $\theta$  increases from  $270^\circ$  to  $360^\circ$ ,

*$\sin \theta$  increases from -1 to 0.*

## (ii) Changes in cosine.

In the first quadrant as the angle  $XOP_1$  increases,  $ON_1$  diminishes, from the value of  $OX$  at  $\theta = 0^\circ$  to the value 0 at  $\theta = 90^\circ$ , and is always positive.

In the second quadrant, as  $\theta$  goes on increasing from  $90^\circ$  to  $180^\circ$ ,  $ON_2$  increases numerically from 0 to  $OX'$  but is

negative. In the third quadrant,  $ON_4$  remains negative, but diminishes numerically from  $OX'$  to 0. In the fourth quadrant,  $ON_5$  is positive and increases from 0 to  $OX$  again.

The hypotenuse remains always positive and is equal to  $OX$  or  $OX'$  in magnitude.

We thus come to the conclusions :

As  $\theta$  increases from  $0^\circ$  to  $90^\circ$ ,

$\cos \theta$  diminishes from 1 to 0.

As  $\theta$  increases from  $90^\circ$  to  $180^\circ$ ,

$\cos \theta$  diminishes from 0 to  $-1$ .

As  $\theta$  increases from  $180^\circ$  to  $270^\circ$ ,

$\cos \theta$  increases from  $-1$  to 0.

As  $\theta$  increases from  $270^\circ$  to  $360^\circ$ ,

$\cos \theta$  increases from 0 to 1.

### (iii) Changes in tangent.

As  $\theta$  goes on increasing from  $0^\circ$  to  $90^\circ$  in the first quadrant,  $P_1N_1$  increases from 0 to  $OY$  and simultaneously  $ON_1$  decreases from  $OX$  to 0, both remaining positive; hence  $\tan \theta = \frac{P_1N_1}{ON_1}$  increases from the value  $\frac{0}{OX} = 0$  to  $\frac{ON}{0} = \infty$ .

In the second quadrant,  $P_2N_2$  diminishes from  $OY$  to 0 while  $ON_2$ , becoming negative, numerically increases from 0 to  $OX'$ . Hence,  $\tan \theta = \frac{P_2N_2}{ON_2}$  is negative but numerically diminishes from  $\infty$  to 0, i.e. increases from  $-\infty$  to 0.

Immediately before  $90^\circ$ ,  $\tan \theta$  is positive and very large, while immediately after  $90^\circ$ ,  $\tan \theta$  is negative and numerically very large. In fact, here, as  $\theta$  passes through the value  $90^\circ$  from the first to the second quadrant, there is a

sudden break or discontinuity in the value of  $\tan \theta$ , which suddenly passes from a very large positive value to a very large negative value, i.e. from the positive to the negative side in passing through infinity.

In the third quadrant, both  $P_4N_4$  and  $ON_4$  are negative and  $P_4N_4$  increases numerically from 0 to  $OY'$  while  $ON_4$  decreases numerically from  $OX'$  to 0. Hence,  $\tan \theta = \frac{P_4N_4}{ON_4}$  is positive, and increases from 0 to  $\infty$ .

In the fourth quadrant,  $P_5N_5$  is negative and numerically diminishes from  $OY'$  to 0 while  $ON_5$  is positive and increases from 0 to  $OX$ . Hence,  $\tan \theta = \frac{P_5N_5}{ON_5}$  is negative and numerically diminishes from  $\infty$  to 0 i.e., increases from  $-\infty$  to 0.

In passing through  $270^\circ$ , there is another discontinuity,  $\tan \theta$  suddenly passing from the positive to the negative side through infinity.

The results are therefore as follows :

As  $\theta$  increases from  $0^\circ$  to  $90^\circ$ ,  $\tan \theta$  increases from 0 to  $\infty$

As  $\theta$  passes through  $90^\circ$ ,  $\tan \theta$  suddenly changes from  
 $+\infty$  to  $-\infty$

As  $\theta$  increases from  $90^\circ$  to  $180^\circ$ ,  $\tan \theta$  increases from  
 $-\infty$  to 0

As  $\theta$  increases from  $180^\circ$  to  $270^\circ$ ,  $\tan \theta$  increases from  
 0 to  $\infty$

As  $\theta$  passes through  $270^\circ$ ,  $\tan \theta$  suddenly changes from  
 $+\infty$  to  $-\infty$

As  $\theta$  increases from  $270^\circ$  to  $360^\circ$ ,  $\tan \theta$  increases from  
 $-\infty$  to 0.

**(iv) Changes in cotangent.**

From the changes in the value of the tangent, the changes in  $\cot \theta = \frac{1}{\tan \theta}$  are traced as follows :

$\theta$  increasing from  $0^\circ$  to  $90^\circ$ ,  $\cot \theta$  diminishes from  $\infty$  to 0

$\theta$  increasing from  $90^\circ$  to  $180^\circ$ ,  $\cot \theta$  diminishes from 0 to  $-\infty$

As  $\theta$  passes through  $180^\circ$ , there is a sudden change in  $\cot \theta$  from  $-\infty$  to  $+\infty$

$\theta$  increasing from  $180^\circ$  to  $270^\circ$ ,  $\cot \theta$  diminishes from  $\infty$  to  $+\infty$

$\theta$  increasing from  $270^\circ$  to  $360^\circ$ ,  $\cot \theta$  diminishes from 0 to  $-\infty$

As  $\theta$  passes through  $360^\circ$ ,  $\cot \theta$  again suddenly changes from  $-\infty$  to  $+\infty$ .

**(v) Changes in secant.**

For  $\sec \theta = \frac{1}{\cos \theta}$ , the results are as follows :

From  $0^\circ$  to  $90^\circ$  for  $\theta$ ,  $\sec \theta$  increases from 1 to  $\infty$ .

Here there is a sudden change from  $+\infty$  to  $-\infty$ .

Then from  $90^\circ$  to  $180^\circ$ ,  $\sec \theta$  increases from  $-\infty$  to  $-1$ .

From  $180^\circ$  to  $270^\circ$ ,  $\sec \theta$  diminishes from  $-1$  to  $-\infty$ .

Here again there is a sudden change from  $-\infty$  to  $+\infty$ .

Then from  $270^\circ$  to  $360^\circ$ ,  $\sec \theta$  diminishes from  $\infty$  to 1.

**(vi) Changes in cosecant.**

For  $\operatorname{cosec} \theta = \frac{1}{\sin \theta}$ , the results are as follows :

From  $0^\circ$  to  $90^\circ$ ,  $\operatorname{cosec} \theta$  diminishes from  $\infty$  to 1.

From  $90^\circ$  to  $180^\circ$ ,  $\operatorname{cosec} \theta$  increases from 1 to  $\infty$ .

Here  $\operatorname{cosec} \theta$  suddenly changes from  $+\infty$  to  $-\infty$ .



Then from  $180^\circ$  to  $270^\circ$ ,  $\operatorname{cosec} \theta$  increases from

$-\infty$  to  $-1$ .

From  $270^\circ$  to  $360^\circ$ ,  $\operatorname{cosec} \theta$  diminishes from  $-1$  to  $-\infty$ .

As  $\theta$  passes through  $360^\circ$ ,  $\operatorname{cosec} \theta$  again suddenly

changes from  $-\infty$  to  $+\infty$ .

**Note.** As  $\theta$  increases by complete multiples of  $2\pi$  (i.e.,  $360^\circ$ ) we know that all the Trigonometrical ratios remain unaltered. Hence after  $360^\circ$ , as  $\theta$  goes on increasing, the same series of values for the ratios are repeated over and over again for each complete revolution of the revolving line. The trigonometrical ratios are therefore all of them periodic functions having the same period  $2\pi$ ,\* after each of which, the same cycle of values is repeated.

The changes traced out above, of the trigonometrical ratios, may be much more clearly demonstrated to the eye from a study of their graphs.

### 103. Graphs of Trigonometrical Functions.

Just like algebraic functions, trigonometrical functions (i.e.,  $\sin x$ ,  $\cos x$ ,  $\sin^2 2x + \tan \frac{x}{2}$ , etc.) may be conveniently represented by means of graphs, showing their changes with the change in the values of the angles.

The method is the same as for graphs in Algebra. Two straight lines  $XOX'$  and  $YOY'$ , intersecting at right angles are taken as axes of co-ordinates. Along the  $x$ -axis, the angles are represented on a suitably chosen scale, positive angles along  $OX$ , and negative angles along  $OX'$ . Along the  $y$ -axis, the values of the trigonometrical functions corresponding to the angles are represented on a suitably chosen scale, positive values being measured upwards (along  $OY$ ), and negative values downwards (along  $OY'$ ). Thus the *abscissa* and *ordinate* of a point stand respectively for an angle and its trigonometrical function.

---

\* $\tan \theta$  and  $\cot \theta$  have a period  $\pi$ .

Plotting a number of points in this way and joining them free-hand, we get the required graph of a given trigonometrical function.

#### 104. Graph of $\sin x$ or *sinc-graph*.

Let  $y = \sin x$ .

Using the table of natural sines, the corresponding values of  $x$  and  $y$  are tabulated corresponding to the values of  $x$  differing by  $10^\circ$  (the values of  $y$  being correct up to two places of decimals) as follows :—

$x$	$-90^\circ$	$-80^\circ$	$-70^\circ$	$-60^\circ$	$-50^\circ$	$-40^\circ$	$-30^\circ$	$-20^\circ$	$-10^\circ$	$0^\circ$
$y$ or $\sin x$	1	.98	.94	.87	.77	.64	.50	.34	.17	0


$x$	$10^\circ$	$20^\circ$	$30^\circ$	$40^\circ$	$50^\circ$	$60^\circ$	$70^\circ$	$80^\circ$	$90^\circ$	$100^\circ$	$110^\circ$	$120^\circ$	etc.
$y$ or $\sin x$	.17	.34	.50	.64	.77	.87	.94	.98	1	.98	.94	.87	etc.

Now let the scale be so chosen that 1 small division along  $OX$  represents  $10^\circ$ , and 10 small divisions along  $OY$  represent unity.\*

The points corresponding to the tabulated values are plotted on the graph paper according to the scale chosen and joined free-hand.

The graph is as shown on the next page (drawn here between the range  $x = -180^\circ$  to  $x = +360^\circ$ ).

\*According to the graph paper supplied and the range within which the graph is to be drawn, the scale should be suitably chosen in each individual case separately.



Sine-Graph

**Note 1.** In the table of natural sines, sines of angles from  $0^\circ$  to  $90^\circ$  only are available. With the help of the formulæ  $\sin(-\theta) = -\sin \theta$ ,  $\sin(180^\circ - \theta) = \sin \theta$ ,  $\sin(180^\circ + \theta) = -\sin \theta$  etc. of Chapter IV, however, the tabulation for  $\sin \theta$  shown above, outside the range of  $0^\circ$  to  $90^\circ$ , is effected.

Similar is the case of tabulation for other graphs in the following pages.

**Note 2.** *Peculiarities of the sine-graph.*

From the figure, the following features will be apparent :—(i) The graph is continuous, and wavy in form ; (ii) The maximum value of  $\sin x$  is  $+1$  and the minimum value is  $-1$ , these values being attained for values of  $x$  which are odd multiples of  $90^\circ$  ; (ii)  $\sin x$  is 0 at the origin and at points for which  $x$  is an even multiple of  $90^\circ$  i.e., any multiple of  $180^\circ$  ; (iii) that  $\sin\left(\frac{\pi}{2} - x\right) = \sin\left(\frac{\pi}{2} + x\right)$ ,  $\sin(\pi - x) = \sin x$ ,  $\sin(-x) = -\sin x$ ,  $\sin(\pi + x) = -\sin x$  etc. ; (v) since  $\sin(2\pi + x) = \sin x$ , the portion between 0 to  $2\pi$  is repeated over and over again on either side.


### 105. Graph of $\cos x$ or cosine-graph.

Let  $y = \cos x$ .

Using the table of natural cosines (see Note 1 of the previous Article), the corresponding values of  $x$  and  $y$  are tabulated at intervals of  $10^\circ$  for  $x$  as follows :—

$x$	$-90^\circ$	$-80^\circ$	$-70^\circ$	$-60^\circ$	$-50^\circ$	$-40^\circ$	$-30^\circ$	$-20^\circ$	$-10^\circ$
$y$ or $\cos x$	0	.17	.34	.50	.64	.77	.87	.94	.98

$x$	$0^\circ$	$10^\circ$	$20^\circ$	$30^\circ$	$40^\circ$	$50^\circ$	$60^\circ$	$70^\circ$	$80^\circ$	$90^\circ$	$100^\circ$	$110^\circ$	etc.
$y$ or $\cos x$	1	.98	.94	.87	.77	.64	.50	.34	.17	0	-.17	-.34	etc.



Cosine-graph

Now choosing the scale such that 1 small division along  $OX$  represents  $10^\circ$ , and 10 small divisions along  $OY$  represent unity, the points corresponding to the above tabulated values are plotted and joined free-hand.

We then get the required graph, which is shown on the annexed page (shown here between the range  $-\pi$  to  $+2\pi$  of  $x$ ).

**Note.** It is apparent from the figure, that the *cosine-graph* is *exactly the same as the sine-graph only shifted whole-sale backwards (to the left) through a space of  $90^\circ$ .*

This is due to the fact that  $\sin(90^\circ + x) = \cos x$ , or  $\sin x = \cos(x - 90^\circ)$  so that the ordinate in the sine-graph corresponding to any value of  $x$  = the ordinate of the cosine-graph corresponding to a value of  $x$  which is  $90^\circ$  less than before.

### 106. Graph of $\tan x$ or *tangent-graph*.


Let  $y = \tan x$ .

Using the table of natural tangents, the corresponding values of  $x$  and  $y$  are tabulated at intervals of  $10^\circ$  of  $x$  as follows :—

$x$	$-20^\circ$	$-10^\circ$	$0^\circ$	$10^\circ$	$20^\circ$	$30^\circ$	$40^\circ$	$50^\circ$	$60^\circ$	$70^\circ$	$80^\circ$	$90^\circ$	$100^\circ$	etc.
$y$ or $\tan x$	$-.36$	$-.18$	$0$	$.18$	$.36$	$.58$	$.84$	$1.19$	$1.73$	$2.75$	$5.67$	$\infty$	$-5.67$	etc.

Now choosing the scale such that 1 small division along  $OX$  represents  $10^\circ$ , and 3 small divisions along  $OY$  represent unity, the points corresponding to the above tabulated values are plotted and joined free-hand.

The graph is as shown on the next page (shown here between the range  $-\pi$  to  $+2\pi$  for  $x$ ).



Tangent-Graph

**Note. Peculiarities of the tangent-graph.**

From the figure, the following points will be apparent : (i) That the curve is not continuous, but consists of separate branches or portions, the points of discontinuities being the values of  $x$  corresponding to the odd multiples of  $\frac{\pi}{2}$ . (ii) As  $x$  passes through these points from the left to the right, the value of  $\tan x$  suddenly changes from very large positive values on the left to very large negative values on the right. (iii) The lines parallel to  $y$ -axis corresponding to the odd multiple of  $\frac{\pi}{2}$  are continuously approached by the graph on either side, but never actually met. Such lines are called *asymptotes* to the curve. (iv) Since  $\tan (n\pi + x) = \tan x$ , each branch, is simply a repetition of the branch from  $-\frac{\pi}{2}$  to  $+\frac{\pi}{2}$ .

**107. Graph of  $\cot x$  or *cotangent-graph*.**

As before the values of  $x$  and  $y$  ( $= \cot x$ ) are tabulated, and with the same scale as in the tangent-graph the points are plotted and joined free-hand.

The graph is shown on the next page (between  $x = -\pi$  to  $x = +2\pi$ ).


This graph also, like the tangent-graph, is *discontinuous*, the points of discontinuity being  $x=0$  and  $x=n\pi$ . The portion between  $x=0$  and  $x=\pi$  is repeated over and over again on either side, as is consequent from the formula  $\cot (n\pi + x) = \cot x$ .

**108. Graph of  $\operatorname{cosec} x$  or *cosecant-graph*.**


The corresponding values of  $x$  and  $y$  are tabulated at intervals of  $10^\circ$  of  $x$  as follows :—

$x$	$-20^\circ$	$-10^\circ$	$0^\circ$	$10^\circ$	$20^\circ$	$30^\circ$	etc.	$80^\circ$	$90^\circ$	$100^\circ$	$110^\circ$	etc.
$y$ or $\operatorname{cosec} x$	-2.92	-5.76	$\infty$	5.76	2.92	2	etc.	1.02	1	1.02	1.06	etc.





Cotangent-Graph



Cosecant-Graph

[ If the table of natural cosecants be not available, the table of natural sines may be used and the values of  $\operatorname{cosec} x = \frac{1}{\sin x}$  may be calculated for different values of  $x$ . ]

The scale is so choosen that 1 small division along  $OX$  represents  $10^\circ$ , and 3 small divisions along  $OY$  represent unity.

The tabulated points are now plotted and joined free-hand.

The graph is shown on the previous page (between the range  $x = -\pi$  to  $x = 2\pi$ ).

**Note.** This graph also consists of *detached branches*, the points of discontinuity being  $x=0$  and  $x=\pi$ . The value of  $y$  never lies between  $\pm 1$ , being always greater than 1 or less than  $-1$ . The lines  $x=\pi$  are asymptotes. The portion between  $x=0$  to  $x=2\pi$  is repeated on either side, over and over again.

### 109. Graph of $\sec x$ or *secant-graph*.


The corresponding values of  $x$  and  $y$  ( $=\sec x$ ) are tabulated as in the caso of cosecant graph, by making use of the table of cosines, if a table of secants be not available.

With the same scale as in the cosecant-graph, the tabulated points have been plotted and joined free-hand.

The graph is shown in the adjoining page (between the range  $x = -\pi$  to  $x = 2\pi$ ).

**Note.** It is apparent from the figure that the *secant-graph* is *exactly the same as the cosecant-graph, only shifted backwards (to the left) through a space of  $90^\circ$ .*

This is due to the fact that  $\operatorname{cosec}(90^\circ + x) = \sec x$ . [ See note below Art. 105. ]



Secant-Graph

### 110. Graphs of other Trigonometrical Expressions.

Graphs of other Trigonometrical functions may be obtained in a similar manner.

We illustrate this by an example.

**Ex.** Draw the graph of  $y = \sin x + \cos x$  between the range  $x = 0$  to  $x = 2\pi$ , and find from the graph the values of  $x$  for which (i)  $y = 0$ , (ii)  $y$  is maximum, (iii)  $y$  is minimum.


[C. U. 1934.]

From the table of natural sines and cosines, corresponding to each value of  $x$ , the values of  $\sin x$  and  $\cos x$  may be separately obtained and then added to give  $y$ ; or else we may write  $y = \sin x + \cos x = \sqrt{2} (\sin x \cos \frac{1}{2}\pi + \cos x \sin \frac{1}{2}\pi) = \sqrt{2} \sin (x + \frac{1}{2}\pi)$ , and corresponding to any value of  $x$ ,  $\sin (x + \frac{1}{2}\pi)$  may be deduced from the sine-table, and this multiplied by  $\sqrt{2} (= 1.414)$  will give  $y$ .

The corresponding values of  $x$  and  $y$  are tabulated at intervals of  $10^\circ$  of  $x$ , between the interval  $x = 0$  to  $x = 2\pi$  as follows :—

$x$	$0^\circ$	$10^\circ$	$20^\circ$	$30^\circ$	$40^\circ$	$50^\circ$	$60^\circ$	$70^\circ$	$80^\circ$	$90^\circ$	$100^\circ$
$y$	1	1.15	1.27	1.37	1.41	1.41	1.37	1.27	1.15	1	.81

$x$	$110^\circ$	$120^\circ$	$130^\circ$	$140^\circ$	$150^\circ$	$160^\circ$	$170^\circ$	$180^\circ$	$190^\circ$	$200^\circ$
$y$	.59	.37	.18	-.13	-.37	-.59	-.81	-1	-1.15	-1.27



Graph of  $\sin x + \cos x$

$x$	210°	220°	230°	240°	250°	260°	270°	280°
$y$	-1·37	-1·41	-1·41	-1·37	-1·27	-1·15	-1	-.81

$x$	290°	300°	310°	320°	330°	340°	350°	360°
$y$	-.59	-.37	-.13	.13	.37	.59	.81	1

The scale is chosen so that 1 small division along  $OX$  represents  $10^\circ$ , and 10 small divisions along  $OY$  represent unity.

The tabulated points are now plotted and joined. The graph is as shown on the previous page.

From the graph we find that (i)  $y=0$  when  $x=135^\circ$  and  $x=315^\circ$ , (ii)  $y$  is maximum when  $x=45^\circ$ , (iii)  $y$  is minimum when  $x=225^\circ$ .

### 111. Graphical Solution of Equations.

Trigonometrical equations, just like algebraic equations, may be solved graphically. In fact in many practical cases particularly where the solutions are not obtained in terms of the standard angles, the graphical method of solution is the only one which is found convenient and is actually adopted. The method is illustrated by the following examples.


**Ex. 1.** Solve graphically the equation  $2 \sin^2 x = \cos 2x$ , giving only those solutions of  $x$  which lie between  $-\frac{\pi}{2}$  and  $\frac{3\pi}{2}$ .  
[ C. U. 1938, '46, '48. ]

We draw two graphs, namely

$$y = 2 \sin^2 x \quad (= 1 - \cos 2x)$$

$$\text{and } y = \cos 2x$$

by tabulating the corresponding values of  $x$  and  $y$  for the two cases separately, making use of the table of natural



Graphical solution of  $2 \sin^2 x = \cos 2x$



cosines, for the range  $x = -\frac{\pi}{2}$  to  $x = \frac{3\pi}{2}$ , at intervals of  $10^\circ$  or  $15^\circ$  of  $x$ .

With the same scale, namely, 1 small division along  $OX$  representing  $10^\circ$ , and 10 small divisions along  $OY$  representing unity, we plot the tabulated values for the two cases in the same graph paper, and joining them, we get the two graphs, as shown in the adjoining page.

We find that the two graphs intersect, and thus have the same abscissæ and ordinates at the points for which

$$x = -\frac{\pi}{6}, \frac{\pi}{6}, \frac{5\pi}{6}, \frac{7\pi}{6}.$$

Thus  $2 \sin^2 x = \cos 2x$  is satisfied for the values of  $x$  given by

$$x = -\frac{\pi}{6}, \frac{\pi}{6}, \frac{5\pi}{6} \text{ and } \frac{7\pi}{6}$$

which are the required solutions within the range

$$-\frac{\pi}{2} \text{ to } \frac{3\pi}{2}.$$


**Ex. 2.** Solve graphically the equation  $\tan x = 2x$  between  $x = 0$  and  $x = \frac{\pi}{2}$ . [C. U. 1939.]

Here,  $x$  is supposed to be measured in radians.

First of all we draw separately the two graphs, namely

$$y = 2x \quad \dots \quad \dots \quad (i)$$

$$\text{and} \quad y = \tan x \quad \dots \quad \dots \quad (ii)$$



Graphical solution of  $\tan x = 2x$ .

The corresponding values of  $x$  and  $y$  within the range  $x=0$  and  $x=\frac{\pi}{2}$  are tabulated in case (i) as follows :

$x$ (in radians)	0	$\frac{\pi}{6}$	$\frac{\pi}{3}$	$\frac{\pi}{2}$
$y$ (i.e. $2x$ ) (numerical value)	0	1.05	2.10	3.15

and in case (ii) as follows :

$x$ (in radians)	0	$\frac{\pi}{18}$	$\frac{2\pi}{18}$	$\frac{3\pi}{18}$	$\frac{4\pi}{18}$	$\frac{5\pi}{18}$	$\frac{6\pi}{18}$	$\frac{7\pi}{18}$	$\frac{8\pi}{18}$	$\frac{\pi}{2}$
$y$ (i.e. $\tan x$ ) (numerical value)	0	.18	.36	.57	.84	1.19	1.73	2.75	5.67	$\infty$

Now choosing the same scale, namely 5 small divisions along  $OX$  to represent  $\frac{\pi}{18}$  radians, and 10 small divisions along  $OY$  to represent unity, we plot the tabulated points for the two cases in the same graph paper and joining them we get the two graphs within the range  $x=0$  and  $x=\frac{\pi}{2}$ , as shown in the adjoining page.

We find that the two graphs intersect at the point given by  $x=0$  and also at the point corresponding to 33.5 small divisions along  $OX$ , which, from our chosen scale, represents  $x = \frac{33.5}{5} \times \frac{\pi}{18}$  radians = 1.17 radians (approximately).

Hence the given equation  $\tan x = 2x$  is satisfied between  $x=0$  and  $x=\frac{\pi}{2}$  by the values of  $x$ , namely  $x=0$  and  $x=1.17$  (approximately), which are the required solutions in radians.

### Examples XVI

1. Draw the graphs of

(i)  $\sin 3x$  between  $x=0^\circ$  to  $x=180^\circ$ .

(ii)  $\tan \frac{2}{3}x$  between  $x = -\frac{1}{2}\pi$  to  $x = \pi$ .

(iii)  $\sin \theta \cos \theta$  between  $\theta = -\pi$  to  $\theta = +\pi$ .

(iv)  $\frac{1}{\cos^2 \theta - \sin^2 \theta}$  between  $\theta = -\frac{\pi}{2}$  to  $+\frac{\pi}{2}$ .

(v)  $\cos (\pi \sin x)$  between  $x=0$  to  $x=\frac{1}{2}\pi$ .

(vi)  $\sin \theta - \sqrt{3} \cos \theta$  between  $\theta=0$  to  $\theta=\pi$ .

(vii)  $\frac{1}{2} \operatorname{cosec} \frac{1}{2}x$  between  $x=0$  to  $x=2\pi$ .

2. (i) Trace the changes in the sign of  $\cos \theta - \sin \theta$  as  $\theta$  changes from  $0^\circ$  to  $360^\circ$ . Verify your conclusions by a graph.

(ii) Trace the changes in sign and magnitude of  $\frac{2 \sin \theta - \sin 2\theta}{2 \sin \theta + \sin 2\theta}$ . [ *B. H. U. 1931.* ]

3. Draw the graph of  $y = \sin (x + \frac{1}{2}\pi)$  between the limits  $x = -\pi$  and  $x = +\pi$ .

4. Draw the graphs of  $\sin \theta$  and  $\cos \theta$  between  $\theta=0$  and  $\theta=\pi$ . Find the points where the graphs intersect.

[ *C. U. 1936, '46.* ]

5. Construct the graphs of  $\tan x$  and  $\cos x$  between 0 and  $\frac{1}{2}\pi$  for  $x$ , making a tabulation of the values of  $y$  dividing the interval into 9 equal parts.

If  $\tan x = \cos x$ , find approximately the value of  $x$  from the above two graphs. [ *C. U. 1943.* ]

6. Obtain graphically a solution of the equation  $\tan x = 1$ , between  $x=0$  and  $x=\frac{1}{2}\pi$ . [ *C. U. 1937.* ]

[ *Draw the graphs of  $y = \tan x$  and  $y = 1$ .* ]

7. Draw the graph of  $\cos x - \sin 2x$  for values of  $x$  lying between  $0^\circ$  and  $90^\circ$  and hence obtain the least value of  $\cos x - \sin 2x$  in this range.

8. Solve graphically the equations :

(i)  $x - \tan x = 0$ , between  $x = 0$  and  $x = \frac{1}{2}\pi$ .

[ C. U. 1945. ]

(ii)  $5 \sin \theta + 2 \cos \theta = 5$ , between  $\theta = 0^\circ$  to  $\theta = 270^\circ$ .

[ Draw the graphs of  $y = 5 \sin \theta + 2 \cos \theta$  and  $y = 5$  and find the common points. ] [ C. U. 1947. ]

(iii)  $\cot \theta - \tan \theta = 2$ , between  $\theta = 0$  to  $\theta = \pi$ .

[ C. U. 1949. ]

(iv)  $\operatorname{cosec} x = \cot x + \sqrt{3}$ , between  $x = 0$  to  $x = \pi$ .

(v)  $\cos x = \sin 2x + \frac{1}{2}$ , between  $x = -\frac{1}{2}\pi$  to  $x = +\frac{1}{2}\pi$ .

(vi)  $5 - \tan x = 2x$ , between 0 and  $2\pi$ .

(vii)  $2 \sin x + x - 3 = 0$ .

(viii)  $x^2 = \cos x$ .

(ix)  $x = \cos^2 x$ .

[ Draw the graphs of  $y = \cos 2x$  and  $y = 2x - 1$ . ]

9. Represent by a graph the displacement given by  
 $s = 2 \sin t + \sin 3t$ .

10. Show graphically that the equation  $2 \sin x + \cos 2x = \frac{1}{2}x$  has only three real roots.

11. Sketch the graphs of :

$y = x$ ,  $y = \sin x$ ,  $y = \tan x$  in  $(-\frac{1}{2}\pi, \frac{1}{2}\pi)$ . From the nature of graphs near the origin, can you suggest any relation among them at the origin ? [ C. U. 1952. ]

## CHAPTER XVII

### MISCELLANEOUS THEOREMS AND EXAMPLES

#### Sec. A

#### HARDER PROBLEMS ON HEIGHTS AND DISTANCES

**112.** Some simple practical applications of Trigonometry, dealing with easy problems on determination of heights and distances, have already been discussed in Chapter V. The problems in the present section are of a more general character, requiring for their solutions, the general relations between the sides and angles of a triangle, as also some geometrical skill.

**113.** *To find the height and the distance of an inaccessible object standing on a horizontal plane.*




Fig. (i)




Fig. (ii)

Let  $CD$  be the object, which is observed from a point  $A$  on a horizontal ground,  $\alpha$  being the observed elevation of its top  $C$ . Let  $h$  be the required height  $CD$  and  $d$  the required distance  $AD$  of the object from  $A$ .

*Case I.* If possible, measure off any suitable distance  $AB (= c)$  from  $A$  directly towards the object, and from  $B$  let the observed elevation of  $C$  be  $\beta$ .

Then from fig. (i),

$$c = AD - BD = h \cot \alpha - h \cot \beta$$

$$h \left( \frac{\cos \alpha}{\sin \alpha} - \frac{\cos \beta}{\sin \beta} \right) = \frac{h \sin (\beta - \alpha)}{\sin \alpha \sin \beta}$$

$$\therefore h = c \sin \alpha \sin \beta \operatorname{cosec} (\beta - \alpha).$$

$$\text{Also } d = AD = h \cot \alpha = c \cos \alpha \sin \beta \operatorname{cosec} (\beta - \alpha).$$

**Note.** Each of the above expressions for  $h$  and  $d$  is in a suitable form for logarithmic computation.

**Case II.** If however it is not convenient to measure the length  $AB$  directly towards the object, we may proceed as follows :

Measure off the length  $AB (= c)$  in *any* suitable direction from  $A$ . From  $A$  let the observed elevation of  $C$  be  $\alpha$  as before. The angles  $CAB$  and  $CBA$  are also observed from  $A$  and  $B$  respectively. Let these be  $\theta$  and  $\phi$ .

We get from fig. (ii) in this case,

$$\text{in } \triangle ABC, \quad \frac{AC}{\sin \phi} = \frac{AB}{\sin C}$$

$$\text{i.e., } AC = \frac{c}{\sin (180^\circ - \theta - \phi)} = \frac{c}{\sin (\theta + \phi)}.$$

$$\therefore AC = c \sin \phi \operatorname{cosec} (\theta + \phi).$$

$$\therefore h = AC \sin \alpha = c \sin \alpha \sin \phi \operatorname{cosec} (\theta + \phi)$$


$$\text{and } d = AD = AC \cos \alpha = c \cos \alpha \sin \phi \operatorname{cosec} (\theta + \phi).$$

**Note.** Here also, the expressions for  $h$  and  $d$  are suitable for calculation by logarithm.

**114.** To find the distance between two visible but inaccessible objects.

Let  $P$  and  $Q$  be the objects whose distance apart is required.

Take two suitable points  $A$  and  $B$  for observation, the distance between which is measured, say  $c$ .



At  $A$ , measure the angles  $PAQ$ ,  $PAB$ , and  $QAB$  (the second observation being unnecessary if all the four points  $P$ ,  $A$ ,  $B$ ,  $Q$  are in the same plane, for in that case,  $\angle PAB = \angle PAQ + \angle QAB$ ). Let these be  $\theta$ ,  $\alpha$  and  $\beta$  respectively.

At  $B$  measure the angles  $PBA$  and  $QBA$ , and let them be  $\gamma$  and  $\delta$ .

$$\text{From } \triangle PAB, \quad \frac{PA}{\sin \gamma} = \frac{c}{\sin (180^\circ - \alpha - \gamma)} = \frac{c}{\sin (\alpha + \gamma)},$$

$$\text{whence,} \quad PA = c \sin \gamma \operatorname{cosec} (\alpha + \gamma).$$

Similarly, from  $\triangle QAB$ ,

$$QA = c \sin \delta \operatorname{cosec} (\beta + \delta).$$

Lastly, from  $\triangle PAQ$ ,

$$PQ^2 = PA^2 + QA^2 - 2PA \cdot QA \cdot \cos \theta.$$


Thus  $PQ$  is determined.

115. Some more illustrative examples of harder problems on heights and distances are worked out below.

**Ex. 1.** *A flagstaff is fixed on the top of a tower standing on a horizontal plane. An observer walking directly towards the foot of the tower, observes the angle subtended*



by the flagstaff from two positions on his path to be the same namely  $\theta$ . The distance between these two positions is  $d$ , and the angle subtended by the tower at his first position is  $\alpha$ . Find the height of the tower, and the length of the flagstaff.



Let  $CD$  be the tower,  $PC$  the flagstaff, whose heights required are  $h$  and  $l$  respectively.  $A$  and  $B$  are the points of observation.

$\therefore \angle PAC = \angle PBC = \theta$ , the points  $P, A, B, C$  are concyclic.

$$\therefore \angle CBD = \angle APC = 90^\circ - \angle PAD = 90^\circ - (\theta + \alpha).$$

$$\begin{aligned} \text{Now } d &= AD - BD = h \cot \alpha - h \cot (CBD) \\ &= h \{ \cot \alpha - \tan (\theta + \alpha) \} \\ &= h \left\{ \frac{\cos \alpha}{\sin \alpha} - \frac{\sin (\theta + \alpha)}{\cos (\theta + \alpha)} \right\} = h \frac{\cos (\theta + 2\alpha)}{\sin \alpha \cos (\theta + \alpha)}. \end{aligned}$$


$$\therefore h = d \sin \alpha \cos (\theta + \alpha) \sec (\theta + 2\alpha).$$

Again, from  $\triangle APC$ ,

$$\frac{l}{\sin \theta} = \frac{AC}{\sin APC} = \frac{h}{\sin \alpha \cos (\theta + \alpha)} = \frac{d}{\cos (\theta + 2\alpha)}.$$

$$\therefore l = d \sin \theta \sec (\theta + 2\alpha).$$

**Ex. 2.** A man walking towards a building, on which a flagstaff is fixed, observes the angle subtended by the flagstaff to be greatest, when he is at a distance  $d$  from the building. If  $\theta$  be the observed greatest angle, find the length of the flagstaff, and the height of the building.



Let  $QB$  be the building, and  $PQ$  the flagstaff. It is easily seen from Geometry that the point of contact  $A$  of a circle through  $P$  and  $Q$  touching the path of the observer on the ground, is the point at which the angle subtended by  $PQ$  is greatest.

Thus  $\angle QAB = \angle APQ = \alpha$  say.

Then,  $\angle PAB + \angle APB = 90^\circ$ ,  
or,  $\theta + 2\alpha = 90^\circ$ . ... (i)

Now,  $PQ = PB - QB = d \tan (\theta + \alpha) - d \tan \alpha$

$$\begin{aligned} &= d \left\{ \frac{\sin (\theta + \alpha)}{\cos (\theta + \alpha)} - \frac{\sin \alpha}{\cos \alpha} \right\} \\ &= d \frac{\sin \theta}{\cos (\theta + \alpha) \cos \alpha} = \frac{2d \sin \theta}{\cos (\theta + 2\alpha) + \cos \theta} \\ &= 2d \tan \theta. \quad [\text{from (i)}] \end{aligned}$$


Also,  $QB = d \tan \alpha = d \tan \left( \frac{\pi}{4} - \frac{\theta}{2} \right)$ .

**Ex. 8.** The angle of elevation of a light at the top of a distant tower from a point 12 ft. above a lake is  $24^\circ 55'$ , and the angle of depression of its reflection in the lake is  $35^\circ 5'$ . Find the height of the tower correct to two decimal places, having given

$$\log 2 = .30103, \quad \log 3 = .47712$$

$$\log 588 = 2.76938, \quad \log 589 = 2.77012$$

$$L \sin 10^\circ 10' = 9.24677.$$



Let  $L$  be the light at the top of the tower  $LM$ ,  $LRO$  the ray from  $L$ , which reflected in the lake at  $R$ , reaches the observer  $O$ , so that  $OR$  is the direction in which the reflexion is seen, and thus, from the laws of reflexion,  $\angle ORA = \angle LRM = \phi$  (say) which is evidently equal to the angle of depression of the reflexion, i.e.,  $35^\circ 5'$ .

Let  $\theta$  denote the angle of elevation of  $L$  from  $O$ , i.e.,  $24^\circ 55'$ .

Now from the figure, in  $\triangle ORL$ ,

$$\frac{RL}{\sin(\theta + \phi)} = \frac{OR}{\sin(\phi - \theta)} = \frac{12}{\sin \phi \sin(\phi - \theta)} \text{ ft.}$$

$$\begin{aligned} \therefore h \equiv LM &= RL \sin \phi = 12 \frac{\sin(\theta + \phi)}{\sin(\phi - \theta)} = 12 \frac{\sin 60^\circ}{\sin(10^\circ 10')} \\ &= \frac{6\sqrt{3}}{\sin(10^\circ 10')} = \frac{2.9^{\frac{8}{5}}}{\sin(10^\circ 10')} \end{aligned}$$

$$\begin{aligned}
 \text{Hence, } \log h &= \log(2.3^{\frac{3}{2}}) - \log \sin(10^\circ 10') \\
 &= \log 2 + \frac{3}{2} \log 3 + 10 - L \sin(10^\circ 10') \\
 &= '30103 + \frac{3}{2}('47712) + 10 - 9'24677 \\
 &= 1'76994.
 \end{aligned}$$

From the given data, it is seen that

$$\log h \text{ lies between } \log 58'8 \text{ and } \log 58'9.$$

$$\begin{aligned}
 \text{Hence, if } h &= 58'8 + x, \text{ diff. for } '1 = 1'77012 - 1'76938 \\
 &= '00074,
 \end{aligned}$$

$$\text{and diff. for } x = 1'76994 - 1'76938 = '00056.$$

$\therefore$  by the theory of proportional parts,

$$\frac{x}{'1} = \frac{56}{74} = '75. \quad \therefore x = '075 = '08 \text{ approximately.}$$

$$\text{Thus, } h = 58'88 \text{ ft.}$$

### Examples XVII (a)

1. The angles of elevation of the top of a palm tree standing on horizontal ground, observed from two points  $A$  and  $B$ , distant 40 and 30 feet from the foot, and in the same straight line with it are found to be complementary. Prove that the height of the tree is nearly 35 feet, and that the angle subtended at the top of the tree by the line  $AB$  is  $\sin^{-1}\frac{1}{7}$ .

2. The angles of elevation of an aeroplane from two places one mile apart and from a point half way between them are found to  $60^\circ$ ,  $30^\circ$  and  $45^\circ$  respectively. Show that the height of the aeroplane is  $440\sqrt{6}$  yards.

3. A building with ten storeys, each of equal height  $x$  ft., stands on one side of a wide street, and from a point

on the other side of the street directly opposite the building, it is observed that the three uppermost storeys together and the two lowest storeys together subtend equal angles. Show that the width of the street is  $x\sqrt{140}$  ft.

4. A two storeyed building has the height of its lower storey 12 ft. and that of the upper storey 13 ft. Find at what distance the two storeys subtend equal angles to an observer's eye at a height 5 feet from the ground.

5. A vertical rod is erected in a horizontal rectangular field  $ABCD$ . The angular elevation of its top from  $A, B, C, D$  are  $\alpha, \beta, \gamma, \delta$ . Show that

$$\cot^2 \alpha - \cot^2 \beta = \cot^2 \delta - \cot^2 \gamma.$$

6. The angles of elevation of a bird flying in a horizontal straight line, from a fixed point at four successive observations are  $\alpha, \beta, \gamma, \delta$ , the observations being taken at equal intervals of time. Assuming that the speed of the bird is uniform, show that

$$\cot^2 \alpha - \cot^2 \delta = 3(\cot^2 \beta - \cot^2 \gamma).$$

7. A man on a hill observes that three towers on a horizontal plane subtend equal angles at his eye and that the angles of depression of their bases are  $\alpha, \beta, \gamma$ . If  $a, b, c$  are the heights of the towers, prove that

$$\frac{\sin(\beta - \gamma)}{a \sin \alpha} + \frac{\sin(\gamma - \alpha)}{b \sin \beta} + \frac{\sin(\alpha - \beta)}{c \sin \gamma} = 0.$$

8. A gun is fired from a fort  $F$  at a distance  $d$  from a station  $O$ , and from two stations  $A$  and  $B$  in a straight line with  $O$  and distant  $a$  and  $b$  respectively from  $O$ , the intervals between seeing the flash and hearing the report are  $t$  and  $t'$ . Show that the velocity of sound is

$$\sqrt{\frac{(d^2 - ab)(a - b)}{at'^2 - bt^2}}.$$

9. A person observes the elevation of the top of a telegraph post which is E. S. E. of him to be  $45^\circ$ , and at noon, the extremity of its shadow is to the N. E. of him; if the length of the shadow be  $x$ , shew that the height of the post is  $x\sqrt{2 - \sqrt{2}}$ .

10. A straight tree on the horizontal ground leans towards the North ; at two points due South and distant  $a, b$  respectively from the foot, the angular elevations of the top of the tree are  $\alpha$  and  $\beta$ . Show that the inclination of the tree to the horizon is

$$\tan^{-1} \left( \frac{a - b}{a \cot \beta - b \cot \alpha} \right).$$

11. An observer on a carriage moving with a speed  $V$  along a straight road observes in one position that two distant trees are in the same line with him which is inclined at an angle  $\theta$  to the road. After a time  $t$ , he observes that the trees subtend their greatest angle  $\phi$  ; show that the distance between the trees is

$$2Vt \sin \theta \sin \phi / (\cos \theta + \cos \phi).$$

12. A train travelling on one of two straight intersecting railways subtends at a certain station on the other line, angles  $\alpha$  and  $\beta$ , when the front of the first carriage and the end of the last carriage reach the junction respectively. Show that the angle of intersection of the two lines is

$$\tan^{-1} \frac{2 \sin \alpha \sin \beta}{\sin (\alpha - \beta)}.$$

13. Two vessels are sailing in parallel directions, and at one instant the bearing of one from the other is  $\alpha^\circ$  N. of E. After an hour's sailing the bearing of the first from the second is  $\beta^\circ$  N. of E., and after another hour the bearing is  $\gamma^\circ$  N. of E. Show that the vessels are sailing in a direction  $\theta^\circ$  N. of E., where

$$\sin (\alpha - \theta) \sin (\gamma - \beta) = \sin (\beta - \alpha) \sin (\gamma - \theta).$$

14. A rod of given length can turn in a vertical plane passing through the sun, one end being fixed on the ground ; if the longest shadow it can cast is  $3\frac{1}{2}$  times the length of the rod, calculate the altitude of the sun, having given

$$\log 3 = \cdot 47712, L \cos 72^\circ 32' 30'' = 9 \cdot 47712.$$

15. A ship sailing N. E. is, at a particular moment, in a line with two light-houses, one of which is situated 5 miles

due N. of the other. In 3 minutes and also in 21 minutes the light-houses are found to subtend a right angle at the ship. Prove that the ship is sailing at the rate of 10 miles an hour, and that the light-houses subtend their greatest angle at the ship at the end of  $3\sqrt{7}$  minutes.

16. A parachute was observed in the N. E. at the elevation  $45^\circ$ ; ten minutes afterwards it was found to be due N. at an elevation  $22\frac{1}{2}^\circ$ . The parachute was descending at the rate of 6 miles per hour, and was all along drifted uniformly towards the west by the wind. Show that wind blows at the rate of 6 miles per hour.

17. A person wishing to determine the height of a distant temple observes the elevation of its top from a point on the horizontal ground through its base to be  $30^\circ$ . On walking a distance  $80\sqrt{3}$  ft. in a certain direction, he finds the elevation of the top to be the same as before, and then on walking a distance 80 ft. at right angles to the former direction, the elevation is found to be  $45^\circ$ . Show that the height of the temple is 80 ft.

18. The shadow of a telegraph post is observed to be half the known height of the post, and sometime afterwards it is equal to the known height; how much will the sun have gone down in the interval, given

$$\log 2 = \cdot 30103, \quad \therefore \tan 63^\circ 26' = 10 \cdot 3009994$$

$$\text{and diff. for } 1' = 3159 ?$$

19. The side of a hill faces due S, and is inclined to the horizon at an angle  $\alpha$ . A straight railway upon it is inclined at an angle  $\beta$  to the horizon; show that the bearing of the railway is

$$\cos^{-1} (\cot \alpha \tan \beta) \text{ E. of N.}$$

20. A spherical time-ball of diameter  $d$  at the top of a tower subtends an angle  $2\alpha$  at a point on the ground from which the elevation of its centre is  $\theta$ ; prove that the height of the centre of the ball above the ground is  $\frac{1}{2}d \sin \theta \operatorname{cosec} \alpha$ .

## Sec. B—SUMMATION OF FINITE SERIES

## 116. Method of Difference.

When the  $r$ th term of a trigonometrical series can be expressed as the difference of two quantities, one of which is the same function of  $r$  as the other is of  $(r+1)$ , the sum of the series may be readily found as illustrated in the Examples 1 and 2 below.

**Ex. 1.** Find the sum of the series.

$$(i) \operatorname{cosec} \theta + \operatorname{cosec} 2\theta + \operatorname{cosec} 2^2\theta + \cdots + \operatorname{cosec} 2^{n-1}\theta.$$

$$(ii) \frac{\sin x}{\sin 2x \sin 3x} + \frac{\sin x}{\sin 3x \sin 4x} + \frac{\sin x}{\sin 4x \sin 5x} + \cdots \text{ to } n \text{ terms.}$$

$$\begin{aligned} (i) \text{ We have } \operatorname{cosec} \theta &= \frac{1}{\sin \theta} = \frac{\sin \frac{1}{2}\theta}{\sin \frac{1}{2}\theta \sin \theta} \\ &= \frac{\sin (\theta - \frac{1}{2}\theta)}{\sin \frac{1}{2}\theta \sin \theta} \\ &= \frac{\sin \theta \cos \frac{1}{2}\theta - \cos \theta \sin \frac{1}{2}\theta}{\sin \frac{1}{2}\theta \sin \theta} \\ &= \cot \frac{1}{2}\theta - \cot \theta. \end{aligned}$$

$$\text{Thus,} \quad \operatorname{cosec} \theta = \cot \frac{1}{2}\theta - \cot \theta.$$

$$\text{Similarly,} \quad \operatorname{cosec} 2\theta = \cot \theta - \cot 2\theta.$$

$$\operatorname{cosec} 2^2\theta = \cot 2\theta - \cot 2^2\theta$$

$$\operatorname{cosec} 2^{n-1}\theta = \cot 2^{n-2}\theta - \cot 2^{n-1}\theta.$$

$\therefore$  by addition, the required sum

$$= \cot \frac{1}{2}\theta - \cot 2^{n-1}\theta.$$



(ii) Here,  $r$ th term

$$\begin{aligned}
 &= \frac{\sin x}{\sin (r+1)x \sin (r+2)x} \\
 &= \frac{\sin \{(r+2) - (r+1)\}x}{\sin (r+1)x \sin (r+2)x} \\
 &= \frac{\sin (r+2)x \cos (r+1)x - \cos (r+2)x \sin (r+1)x}{\sin (r+1)x \sin (r+2)x} \\
 &= \cot (r+1)x - \cot (r+2)x.
 \end{aligned}$$

Putting  $r = 1, 2, 3, \dots, n$  and adding, the sum of the required series would be found to be

$$\cot 2x - \cot (n+2)x.$$

**Ex. 2.** Find the sum of the series

$$\begin{aligned}
 \tan^{-1} \frac{x}{1+1.2x^2} + \tan^{-1} \frac{x}{1+2.3x^2} + \dots \\
 + \tan^{-1} \frac{x}{1+n(n+1)x^2}
 \end{aligned}$$

$$\begin{aligned}
 \text{Here, } r\text{th term} &= \tan^{-1} \frac{x}{1+r(r+1)x^2} \\
 &= \tan^{-1} \frac{(r+1)x - rx}{1+(r+1)x \cdot rx} \\
 &= \tan^{-1}(r+1)x - \tan^{-1}rx.
 \end{aligned}$$

putting  $r = 1, 2, 3, \dots, n$ , we have

$$\tan^{-1} \frac{x}{1+1.2x^2} = \tan^{-1}2x - \tan^{-1}x$$

$$\tan^{-1} \frac{x}{1+2.3x^2} = \tan^{-1}3x - \tan^{-1}2x$$

.....

$$\tan^{-1} \frac{x}{1+n(n+1)x^2} = \tan^{-1}(n+1)x - \tan^{-1}nx.$$

by addition, the required sum

$$= \tan^{-1}(n+1)x - \tan^{-1}x.$$

117. Sometimes the  $r$ th term of a series, *when multiplied by a factor*, can be expressed as the difference of two quantities one of which is the same function of  $r$  as the other is of  $(r+1)$ . It is illustrated in the following two cases.

**(I) Sum of sines of  $n$  angles in A. P.**

Let the angles in A.P. be  $\alpha, \alpha + \beta, \alpha + 2\beta, \dots, \{\alpha + (n-1)\beta\}$ , the first term being  $\alpha$ , and the common difference,  $\beta$ .

Let  $S$  denote the sum of the series,

$$\sin \alpha + \sin (\alpha + \beta) + \sin (\alpha + 2\beta) + \dots + \sin \{\alpha + (n-1)\beta\}.$$

Multiplying each term of the above series by  $2 \sin$  (*half difference*), i.e., by  $2 \sin \frac{1}{2}\beta$ , we have,

$$\begin{aligned} 2 \sin \alpha \sin \frac{1}{2}\beta &= \cos \left(\alpha - \frac{1}{2}\beta\right) - \cos \left(\alpha + \frac{1}{2}\beta\right) \\ 2 \sin (\alpha + \beta) \sin \frac{1}{2}\beta &= \cos \left(\alpha + \frac{1}{2}\beta\right) - \cos \left(\alpha + \frac{3}{2}\beta\right) \\ 2 \sin (\alpha + 2\beta) \sin \frac{1}{2}\beta &= \cos \left(\alpha + \frac{3}{2}\beta\right) - \cos \left(\alpha + \frac{5}{2}\beta\right) \\ &\dots \quad \dots \quad \dots \quad \dots \\ 2 \sin \{\alpha + (n-1)\beta\} \sin \frac{1}{2}\beta \\ &= \cos \left(\alpha + \frac{2n-3}{2}\beta\right) - \cos \left(\alpha + \frac{2n-1}{2}\beta\right). \end{aligned}$$

Adding vertically, we have

$$\begin{aligned} 2 \sin \frac{1}{2}\beta \cdot S &= \cos \left(\alpha - \frac{\beta}{2}\right) - \cos \left(\alpha + \frac{2n-1}{2}\beta\right) \\ &= 2 \sin \left(\alpha + \frac{n-1}{2}\beta\right) \sin \frac{n\beta}{2}. \end{aligned}$$

$$\therefore S = \frac{\sin \frac{n\beta}{2}}{\sin \frac{\beta}{2}} \cdot \sin \left(\alpha + \frac{n-1}{2}\beta\right).$$

**Cor.** Putting  $\beta = \alpha$ , we get

$$\sin \alpha + \sin 2\alpha + \sin 3\alpha + \dots + \sin n\alpha$$

$$= \frac{\sin \frac{n\alpha}{2}}{\sin \frac{\alpha}{2}} \sin \frac{n+1}{2}\alpha.$$

**(II) Sum of cosines of  $n$  angles in A.P.**

As before, let  $S$  denote the sum of the series

$$\cos a + \cos (a + \beta) + \cos (a + 2\beta) + \dots + \cos \{a + (n-1)\beta\}.$$

Multiplying each term of the above series by

$2 \sin (\text{half difference}),$  we have,

$$\begin{aligned} 2 \cos a \cdot \sin \frac{1}{2}\beta &= \sin \left(a + \frac{1}{2}\beta\right) - \sin \left(a - \frac{1}{2}\beta\right) \\ 2 \cos (a + \beta) \cdot \sin \frac{1}{2}\beta &= \sin \left(a + \frac{3}{2}\beta\right) - \sin \left(a + \frac{1}{2}\beta\right) \\ 2 \cos (a + 2\beta) \cdot \sin \frac{1}{2}\beta &= \sin \left(a + \frac{5}{2}\beta\right) - \sin \left(a + \frac{3}{2}\beta\right) \\ &\dots \qquad \dots \qquad \dots \qquad \dots \\ 2 \cos \{a + (n-1)\beta\} \cdot \sin \frac{1}{2}\beta & \end{aligned}$$

$$= \sin \left(a + \frac{2n-1}{2}\beta\right) - \sin \left(a + \frac{2n-3}{2}\beta\right).$$

Adding vertically, we have

$$\begin{aligned} 2 \sin \frac{1}{2}\beta \cdot S &= \sin \left(a + \frac{2n-1}{2}\beta\right) - \sin \left(a - \frac{\beta}{2}\right) \\ &= 2 \cos \left(a + \frac{n-1}{2}\beta\right) \sin \frac{n\beta}{2}. \end{aligned}$$

$$\therefore S = \frac{\sin \frac{n\beta}{2}}{\sin \frac{\beta}{2}} \cdot \cos \left(a + \frac{n-1}{2}\beta\right).$$

**Cor.** Putting  $\beta = a$ , we get

$$\cos a + \cos 2a + \cos 3a + \dots + \cos na = \frac{\sin \frac{na}{2}}{\sin \frac{a}{2}} \cdot \cos \frac{n+1}{2}a.$$

**Note.** The sum of the cosine series may be deduced from that of the sine series by writing  $a + \frac{\pi}{2}$  for  $a$ .

As an aid to memory, the two formulæ of this article may be expressed in language as follows :

$$\text{since, } \alpha + \frac{n-1}{2}\beta = \alpha + \frac{\alpha + (n-1)\beta}{2},$$

$\therefore$  Sum of sines of  $n$  angles in A.P.

$$= \frac{\sin \frac{n \cdot \text{diff.}}{2}}{\sin \frac{\text{diff.}}{2}} \sin \frac{\text{first angle} + \text{last angle}}{2}.$$

Sum of cosines of  $n$  angles in A.P.

$$= \frac{\sin \frac{n \cdot \text{diff.}}{2}}{\sin \frac{\text{diff.}}{2}} \cos \frac{\text{first angle} + \text{last angle}}{2}.$$

**Note.** From the above formulæ, it is clear that if  $\sin \frac{n\beta}{2} = 0$ , then the sum of the sine series as also the sum of the cosine series is zero.

Now, if  $\sin \frac{n\beta}{2} = 0$ , then  $\frac{n\beta}{2} = k\pi$  or  $\beta = \frac{2k\pi}{n}$ , where  $k$  is an integer.

Thus, the sum of the sines and the sum of the cosines of  $n$  angles in A. P. are each equal to zero when the common difference of the angles is an even multiple of  $\frac{\pi}{n}$ .

**Ex. 1.** Find the sum of  $n$  terms of the series

$$\sin \alpha - \sin (\alpha + \beta) + \sin (\alpha + 2\beta) - \dots$$

Since,  $\sin (\pi + \theta) = -\sin \theta$  ;  $\sin (2\pi + \theta) = \sin \theta$  etc.

$\therefore$  the series is equal to

$$\sin \alpha + \sin (\pi + \alpha + \beta) + \sin (2\pi + \alpha + 2\beta) + \dots$$

i.e., equal to a series in which the common difference of the angles is  $\beta + \pi$  and the last angle is  $\alpha + (n-1)(\beta + \pi)$ .

$$S = \frac{\sin \frac{n(\beta + \pi)}{2}}{\sin \frac{\beta + \pi}{2}} \sin \left\{ \alpha + \frac{(n-1)(\beta + \pi)}{2} \right\}.$$

**Ex. 2.** Find the sum of the series

$$\sin^2 \theta + \sin^2 2\theta + \sin^2 3\theta + \dots + \sin^2 n\theta.$$

Since,  $\sin^2 \theta = \frac{1}{2}(1 - \cos 2\theta)$  ;  $\sin^2 2\theta = \frac{1}{2}(1 - \cos 4\theta)$  ; &c.

$\therefore$  the given series

$$\begin{aligned} &= \frac{1}{2}(1 - \cos 2\theta) + \frac{1}{2}(1 - \cos 4\theta) + \dots + \frac{1}{2}(1 - \cos 2n\theta) \\ &= \frac{n}{2} - \frac{1}{2}(\cos 2\theta + \cos 4\theta + \dots + \cos 2n\theta) \\ &= \frac{n}{2} - \frac{1}{2} \frac{\sin n\theta}{\sin \theta} \cos (n+1)\theta. \quad [\text{by Art. 117}] \end{aligned}$$

**Ex. 3.** Sum the series

$$\begin{aligned} &\cos \alpha + 2 \cos (\alpha + \beta) + 3 \cos (\alpha + 2\beta) + \dots \\ &\dots + n \cos \{\alpha + (n-1)\beta\}. \end{aligned}$$

Let  $u_r$  denote the  $r$ th term and  $S$  denote the sum of the given series.

$$\begin{aligned} \text{Now, } 2 \cos \beta \cdot u_r &= 2 \cos \beta \cdot r \cos \{\alpha + (r-1)\beta\} \\ &= r[\cos (\alpha + r\beta) + \cos \{\alpha + (r-2)\beta\}]. \end{aligned}$$

$\therefore$  putting  $r=1, 2, 3, \dots, n$  and adding together,

we get  $2 \cos \beta \cdot S$ .

Now, subtract  $2 \cos \beta \cdot S$  from  $2S$  ; then

$$\begin{aligned} 2S(1 - \cos \beta) &= (n+1) \cos \{\alpha + (n-1)\beta\} \\ &\quad - \cos (\alpha - \beta) - n \cos (\alpha + n\beta). \end{aligned}$$

Then, dividing by  $2(1 - \cos \beta)$ ,  $S$ , the sum of the required series would be obtained.

**Note.** Similarly the sum of the series

$\sin \alpha + 2 \sin (\alpha + \beta) + 3 \sin (\alpha + 2\beta) + \dots + n \sin \{\alpha + (n-1)\beta\}$   
would be obtained.

## Examples XVII(b)

Sum the following series to  $n$  terms :

1.  $\sin a + \sin \left( a - \frac{\pi}{n} \right) + \sin \left( a - \frac{2\pi}{n} \right) + \dots$
2.  $\cos a + \cos \left( a + \frac{2\pi}{n} \right) + \cos \left( a + \frac{4\pi}{n} \right) + \dots$
3.  $\sin a - \sin 2a + \sin 3a - \dots$
4.  $\cos^2 \theta + \cos^2 2\theta + \cos^2 3\theta + \dots$
5.  $\sin^3 a + \sin^3 3a + \sin^3 5a + \dots$
6.  $\sin^3 \theta - \sin^3 2\theta + \sin^3 3\theta - \sin^3 4\theta + \dots$
7.  $\sin^4 a + \sin^4 2a + \sin^4 3a + \dots$
8.  $\cos \theta - \sin 2\theta - \cos 3\theta + \sin 4\theta + \cos 5\theta - \sin 6\theta - \dots$
9.  $\sin a \sin 2a + \sin 2a \sin 3a + \sin 3a \sin 4a + \dots$
10.  $\cos a \cos 3a + \cos 3a \cos 5a + \cos 5a \cos 7a + \dots$

Find the sum of the following series :

11.  $\cos \frac{\pi}{19} + \cos \frac{3\pi}{19} + \cos \frac{5\pi}{19} + \dots + \cos \frac{17\pi}{19}$
12.  $\sin 5^\circ + \sin 77^\circ + \sin 149^\circ + \dots + \sin 293^\circ$
13.  $\sin \frac{2\pi}{n} + \sin \frac{4\pi}{n} + \sin \frac{6\pi}{n} + \dots + \sin \frac{2n\pi}{n}$
14.  $\sin na + \sin (n-1)a + \sin (n-2)a + \dots$  to  $2n$  terms.
15. Prove that

$$(i) \frac{\sin \theta + \sin 3\theta + \sin 5\theta + \dots \text{ to } n \text{ terms}}{\cos \theta + \cos 3\theta + \cos 5\theta + \dots \text{ to } n \text{ terms}} = \tan n\theta.$$

$$(ii) \sin^2 a + \sin^2 \left( a + \frac{2\pi}{n} \right) + \sin^2 \left( a + \frac{4\pi}{n} \right) + \dots \text{ to } n \text{ terms} \\ = \frac{1}{2}n.$$

Sum to  $n$  terms :

$$16. \sec a \sec 2a + \sec 2a \sec 3a + \sec 3a \sec 4a + \dots$$

$$17. \frac{1}{\sin \theta \sin 2\theta} + \frac{1}{\sin 2\theta \sin 3\theta} + \frac{1}{\sin 3\theta \sin 4\theta} + \dots$$

$$18. \frac{1}{\cos a + \cos 3a} + \frac{1}{\cos a + \cos 5a} + \frac{1}{\cos a + \cos 7a} + \dots$$

$$19. \cot \theta \cot 2\theta + \cot 2\theta \cot 3\theta + \cot 3\theta \cot 4\theta + \dots$$

$$20. \tan a + 2 \tan 2a + 4 \tan 4a + 8 \tan 8a + \dots$$

$$[ \tan a = \cot a - 2 \cot 2a ]$$

$$21. \sin 2\theta \sin^2 \frac{2\theta}{2} + \sin 3\theta \sin^2 \frac{3\theta}{2} + \sin 4\theta \sin^2 \frac{4\theta}{2} + \dots$$

$$22. \frac{\sin x}{\cos 3x} + \frac{\sin 3x}{\cos 3^2 x} + \frac{\sin 3^2 x}{\cos 3^3 x} + \dots$$

$$[ \text{1st term} = \frac{1}{2}(\tan 3x - \tan x) ]$$

$$23. \tan^{-1} \frac{1}{1+1+1^2} + \tan^{-1} \frac{1}{1+2+2^2} + \tan^{-1} \frac{1}{1+3+3^2} + \dots$$

$$24. \tan^{-1} \frac{2}{1+1.3} + \tan^{-1} \frac{2}{1+3.5} + \tan^{-1} \frac{2}{1+5.7} + \dots$$

$$25. \cot^{-1} (2.1^2) + \cot^{-1} (2.2^2) + \cot^{-1} (2.3^2) + \dots$$

$$26. \tan x + \frac{1}{2} \tan \frac{1}{2} x + \frac{1}{2^2} \tan \frac{1}{2^2} x + \dots$$

$$27. \cos x \cos 2x \cos 3x + \cos 2x \cos 3x \cos 4x + \dots$$

$$28. \cos \theta + 2 \cos 2\theta + 3 \cos 3\theta + \dots + n \cos n\theta.$$

$$29. \text{Find the sums of the series}$$

$$(i) \sin a + \sin 2a + \sin 3a + \dots \text{ to } n \text{ terms}$$

$$\text{and } (ii) \sin a + \sin 3a + \sin 5a + \dots \text{ to } n \text{ terms}$$

and hence deduce respectively the sums of the series

(a)  $1 + 2 + 3 + \dots$  to  $n$  terms

and (b)  $1 + 3 + 4 + \dots$  to  $n$  terms.

30. Sum the series

$\tan x \tan 2x + \tan 2x \tan 3x + \dots + \tan nx \tan (n+1)x$   
and hence deduce the sum of the series

$$1.2 + 2.3 + \dots + n(n+1).$$

31. If  $\beta$  be the exterior angle of a regular polygon of  $n$  sides, show that

(i)  $\sin \alpha + \sin (\alpha + \beta) + \sin (\alpha + 2\beta) + \dots$  to  $n$  terms  $= 0$ .

(ii)  $\cos \alpha + \cos (\alpha + \beta) + \cos (\alpha + 2\beta) + \dots$  to  $n$  terms  $= 0$ .

32. A regular polygon of  $n$  sides is inscribed in a circle of radius  $a$ ; prove that

(i) the sum of the lengths of the perpendiculars drawn from the angular points upon any diameter is zero;

(ii) the sum of the lengths of the lines joining any one vertex to each of the other vertices is  $2a \cot \frac{\pi}{2n}$ .

### Sec. C—ELIMINATION

118. The elimination of trigonometrical functions from given equations is a very important and common mathematical problem. There are no set rules to effect the elimination. The form of the equations will often suggest special methods, and in addition to the usual algebraical artifices, we shall always have at our disposal the identical relations subsisting among the trigonometrical functions.

The following examples will illustrate some useful methods of elimination.



**Ex. 1.** *Eliminate  $\theta$  between the equations*

$$a \cos \theta + b \sin \theta + c = 0$$

$$a' \cos \theta + b' \sin \theta + c' = 0.$$

From the given equations, we have, by cross-multiplication,

$$\frac{\cos \theta}{bc' - b'c} = \frac{\sin \theta}{ca' - c'a} = \frac{1}{ab' - a'b}.$$

$$\therefore \cos \theta = \frac{bc' - b'c}{ab' - a'b}, \text{ and } \sin \theta = \frac{ca' - c'a}{ab' - a'b}.$$

Squaring and adding, we get

$$(bc' - b'c)^2 + (ca' - c'a)^2 = (ab' - a'b)^2$$

as the required eliminant.

**Ex. 2.** *Eliminate  $\theta$  from the equations*

$$x \sin \theta + y \cos \theta = 2a \sin 2\theta$$

$$x \cos \theta - y \sin \theta = a \cos 2\theta.$$

Solving as simultaneous equations in  $x$  and  $y$ , we have

$$x = a(\cos 2\theta \cos \theta + 2 \sin 2\theta \sin \theta)$$

$$= a[\cos (2\theta - \theta) + \sin 2\theta \sin \theta]$$

$$= a(\cos \theta + 2 \sin^2 \theta \cos \theta),$$

$$y = a(2 \sin 2\theta \cos \theta - \cos 2\theta \sin \theta)$$

$$= a(\sin \theta + \sin 2\theta \cos \theta) = a(\sin \theta + 2 \sin \theta \cos^2 \theta).$$

$$\therefore x + y = a(\sin \theta + \cos \theta)(1 + 2 \sin \theta \cos \theta)$$

$$= a(\sin \theta + \cos \theta)(\sin \theta + \cos \theta)^2 = a(\cos \theta + \sin \theta)^3;$$

Similarly,

$$x - y = a(\cos \theta - \sin \theta)(1 - 2 \sin \theta \cos \theta)$$

$$= a(\cos \theta - \sin \theta)^3.$$

$$\therefore a^{\frac{1}{3}}(\cos \theta + \sin \theta) = (x + y)^{\frac{1}{3}} \quad \dots (i)$$

$$a^{\frac{1}{3}}(\cos \theta - \sin \theta) = (x - y)^{\frac{1}{3}} \quad \dots (ii)$$

Hence, squaring and adding (i) and (ii), we have,

$$(x + y)^{\frac{2}{3}} + (x - y)^{\frac{2}{3}} = 2a^{\frac{2}{3}}$$

as the required eliminant.

**Ex. 3.** *Eliminate  $x$  and  $y$  from the equations*

$$a \sin^2 x + b \cos^2 x = c, \quad b \sin^2 y + a \cos^2 y = d,$$

$$a \tan x = b \tan y.$$

From the first equation, we have

$$a \sin^2 x + b \cos^2 x = c (\sin^2 x + \cos^2 x).$$

$$\therefore (a - c) \sin^2 x = (c - b) \cos^2 x.$$

$$\therefore \tan^2 x = \frac{c - b}{a - c}.$$

From the second equation, we have similarly

$$b \sin^2 y + a \cos^2 y = d (\sin^2 y + \cos^2 y).$$

$$\therefore \tan^2 y = \frac{d - a}{b - d}.$$

From the third equation,

$$a^2 \tan^2 x = b^2 \tan^2 y.$$

$$\therefore \frac{a^2(c - b)}{a - c} = \frac{b^2(d - a)}{b - d}.$$

This, when simplified, reduces to

$$\frac{1}{a} + \frac{1}{b} = \frac{1}{c} + \frac{1}{d}, \text{ the required eliminant.}$$

## Examples XVII(c)

Eliminate  $\theta$  from the following pair of equations

$$1. \quad \cot \theta (1 + \sin \theta) = 4a.$$

$$\quad \cot \theta (1 - \sin \theta) = 4b.$$

$$2. \quad x = a \cos \theta + b \cos 2\theta$$

$$y = a \sin \theta + b \sin 2\theta.$$

$$3. \quad x = \tan \theta + \tan 2\theta$$

$$y = \cot \theta + \cot 2\theta.$$

$$4. \quad a \sin \theta + b \cos \theta = 1.$$

$$a \operatorname{cosec} \theta - b \sec \theta = 1.$$

$$5. \quad x = \sin \theta + \cos \theta \sin 2\theta$$

$$y = \cos \theta + \sin \theta \sin 2\theta.$$

$$6. \quad x + a = a (2 \cos \theta - \cos 2\theta)$$

$$y = a (2 \sin \theta - \sin 2\theta).$$

$$7. \quad x = 3 \sin \theta - \sin 3\theta$$

$$y = \cos 3\theta + 3 \cos \theta.$$

$$8. \quad x = \cot \theta + \tan \theta$$

$$y = \sec \theta - \cos \theta.$$

$$9. \quad x \sin \theta - y \cos \theta = \sqrt{x^2 + y^2}$$

$$\frac{\cos^2 \theta}{a^2} + \frac{\sin^2 \theta}{b^2} = \frac{1}{x^2 + y^2}.$$

$$10. \quad \frac{x}{a} = \cos \theta + \cos 2\theta$$

$$\frac{y}{b} = \sin \theta + \sin 2\theta.$$

# ELIMINATION

$$11. \frac{ax}{\cos \theta} - \frac{by}{\sin \theta} = a^2 - b^2$$

$$\frac{ax \sin \theta}{\cos^2 \theta} + \frac{by \cos \theta}{\sin^2 \theta} = 0.$$

$$12. \frac{x}{a} \cos \theta - \frac{y}{b} \sin \theta = \cos 2\theta$$

$$\frac{x}{a} \sin \theta + \frac{y}{b} \cos \theta = 2 \sin 2\theta.$$

$$13. x = \operatorname{cosec} \theta - \sin \theta$$

$$y = \sec \theta - \cos \theta.$$

$$14. \sin \theta + \cos \theta = a$$

$$\sin^3 \theta + \cos^3 \theta = b.$$

$$15. \frac{x}{a} \cos \theta + \frac{y}{b} \sin \theta = 1$$

$$x \sin \theta - y \cos \theta = (a^2 \sin^2 \theta + b^2 \cos^2 \theta)^{\frac{1}{2}}$$

Eliminate  $\theta$  and  $\phi$  from the following equations  
(Ex. 16-19) :—

$$16. \sin \theta + \sin \phi = x, \cos \theta + \cos \phi = y, \theta - \phi = a.$$

$$17. \tan \theta + \tan \phi = a, \cot \theta + \cot \phi = b, \theta + \phi = a.$$

$$18. a \sin^2 \theta + b \cos^2 \theta = a \cos^2 \phi + b \sin^2 \phi = 1,$$

$$a \tan \theta = b \tan \phi.$$

$$19. \sin \theta + \sin \phi = a, \cos \theta + \cos \phi = b, \sin 2\theta + \sin 2\phi = 2c.$$

$$20. \text{ If } (a+b) \tan (\theta - \phi) = (a-b) \tan (\theta + \phi) \text{ and}$$

$a \cos 2\phi + b \cos 2\theta = c$ , show that


$$a^2 - b^2 + c^2 = 2ac \cos 2\phi.$$

## APPENDIX

1. To prove that

$$\sin \theta < \theta < \tan \theta$$

where  $\theta$  is the circular measure of any positive acute angle.



Let  $\angle AOP$  be a positive acute angle whose radian measure is  $\theta$ , and let  $\angle QOA$  be an equal angle on the other side of  $OA$ . With centre  $O$  and any radius, a circle is drawn cutting  $OP$ ,  $OA$ ,  $OQ$  at  $P$ ,  $A$ ,  $Q$  respectively.  $PQ$  is joined cutting  $OA$  at  $N$ . The triangles  $OPN$  and  $OQN$  are easily seen to be congruent, so that  $PN = QN$  and  $PNQ$  is perpendicular to  $OA$ . The tangent  $PT$  to the circle at  $P$  cutting  $OA$  at  $T$ ,  $\angle OPT$  is a right angle.  $TQ$  being joined, the triangles  $OPT$  and  $OQT$  are easily proved to be congruent, so that  $TP = TQ$ .

The figure is thus symmetrical about  $OA$ .

Then, from the figure,

$$\sin \theta = \frac{PN}{OP} = \frac{1}{2} \cdot \frac{PQ}{OP}$$

$$\theta = \frac{\text{arc } PAQ}{OP} = \frac{1}{2} \frac{\text{arc } PAQ}{OP}$$

$$\tan \theta = \frac{PT}{OP} = \frac{1}{2} \cdot \frac{PT + QT}{OP}$$

Now we may take it as axiomatic that the straight line  $PQ$  is less than the curved arc  $PAQ$ , and that the curved arc  $PAQ$  which always bends the same way, being within the triangle  $PTQ$ , is less than  $PT + TQ$ .


Hence, since  $PQ < \text{arc } PAQ < PT + TQ$ ,

we have, on dividing throughout by  $2OP$

$$\sin \theta < \theta < \tan \theta.$$

*Alternative method :*

Let  $ABC$  be a circle whose centre is  $O$  and radius  $r$ .



Let  $AOB = \theta$  radians.

Draw  $BT$  the tangent at  $B$  to meet  $OA$  produced at  $T$ ; then  $BT = r \tan \theta$ .

We know that if the angle of a sector of a circle of radius  $r$  is  $\theta$  radians, its area  $= \frac{1}{2}r^2\theta$ .

Now from the figure it is obvious that

$$\triangle OAB < \text{sector } OAB < \triangle OBT.$$

$$\therefore \frac{1}{2}r^2 \sin \theta < \frac{1}{2}r^2 \theta < \frac{1}{2}r \cdot r \tan \theta,$$

$$\text{i.e.,} \quad \sin \theta < \theta < \tan \theta.$$

**Cor.** If now  $\theta$  becomes infinitely small, we can prove

$$\lim_{\theta \rightarrow 0} \frac{\sin \theta}{\theta} = 1,$$

$$\lim_{\theta \rightarrow 0} \cos \theta = 1,$$

$$\text{and } \lim_{\theta \rightarrow 0} \frac{\tan \theta}{\theta} = 1.$$

For, since,  $\sin \theta < \theta < \tan \theta$ ,  
we get, by dividing by  $\sin \theta$ ,

$$1 < \frac{\theta}{\sin \theta} < \frac{1}{\cos \theta}.$$

This is true, however small  $\theta$  may be, provided it is positive. When  $\theta$  becomes infinitely small,  $OP$  and  $ON$  practically come into coincidence, so that

$$\cos \theta = \frac{ON}{OP} \text{ ultimately becomes } 1.$$

$$\text{Hence, } \lim_{\theta \rightarrow 0} \cos \theta = 1.$$

In that case  $\frac{1}{\cos \theta}$  also tends to the value 1. But  $\frac{\theta}{\sin \theta}$  always lies between 1 and  $\frac{1}{\cos \theta}$  which ultimately come into coincidence, and so  $\frac{\theta}{\sin \theta}$  also ultimately coincides with 1.

$$\text{Thus } \frac{\sin \theta}{\theta} = 1 \text{ in the limit.}$$

Again, from

$$\sin \theta < \theta < \tan \theta,$$


We get by dividing by  $\tan \theta$ ,

$$\cos \theta < \frac{\theta}{\tan \theta} < 1$$

and as  $\theta \rightarrow 0$ ,  $\cos \theta \rightarrow 1$  and  $\frac{\theta}{\tan \theta}$  always lying between  $\cos \theta$  and 1 which come into coincidence,  $\frac{\theta}{\tan \theta} = 1$  in the limit, and so  $\lim_{\theta \rightarrow 0} \frac{\tan \theta}{\theta} = 1$ .

Hence, the results.

2. *Formulae for  $\sin(A+B)$  and  $\cos(A+B)$  where  $A$  and  $B$  are of any magnitude. (Generalization of Art. 33.)*



In Article 33, formulæ for  $\sin(A+B)$  and  $\cos(A+B)$  were deduced geometrically with a figure in which  $A$  and  $B$  were acute and  $(A+B)$  less than  $90^\circ$ . We now prove them in a more general case.

Let a revolving line, starting from  $OX$ , trace out an angle  $XOZ = A$  and further trace out an angle  $ZOP = B$ , so that the total angle traced out is  $(A+B)$ . From any point  $P$  on the final position of the revolving line,  $PN$  and  $PT$  are drawn perpendiculars to  $OX$  and  $OZ$  (produced if necessary, as in the above figure), and from  $T$  perpendiculars  $TM$  and  $TR$  are drawn on  $OX$  and  $PN$  (produced if necessary),

In the figure above,  $\angle POT = B - 180^\circ$ , and since  $PN$  and  $PT$  are perpendiculars to  $OX$  and  $OZ$  respectively,  $\angle TPR = \angle TON = 180^\circ - \angle XOZ$  i.e.,  $180^\circ - A$ .

In considering  $\sin(A+B)$  and  $\cos(A+B)$  from the triangle  $NOP$ , it is to be noted that  $PN$  is negative and  $ON$  and  $OP$  are positive.

If we consider only the positive magnitudes of the sides of the acute-angled triangle  $OTM$ ,  $PTR$  and  $OPT$ , then  $PN$  with its proper sign may be written as  $-(TM - PR)$ , and  $ON$  with its proper sign may be written as  $OM + TR$ .

Now, from the figure,

$$\sin(A+B) = \frac{PN}{OP} = \frac{TM - PR}{OP}$$




$$\begin{aligned}
&= -\frac{TM}{OT} \cdot \frac{OT}{OP} + \frac{PR}{PT} \cdot \frac{PT}{OP} \\
&= -\sin TOM \cos POT + \cos TPR \sin POT \\
&= -\sin (180^\circ - A) \cos (B - 180^\circ) \\
&\quad + \cos (180^\circ - A) \sin (B - 180^\circ) \\
&= -\sin A (-\cos B) + (-\cos A) (-\sin B) \\
&= \sin A \cos B + \cos A \sin B.
\end{aligned}$$

Again,

$$\begin{aligned}
\cos (A + B) &= \frac{ON}{OP} = \frac{OM + RT}{OP} \\
&= \frac{OM}{OT} \cdot \frac{OT}{OP} + \frac{RT}{PT} \cdot \frac{PT}{OP} \\
&= \cos TOM \cos POT + \sin TPR \sin POT \\
&= \cos (180^\circ - A) \cos (B - 180^\circ) \\
&\quad + \sin (180^\circ - A) \sin (B - 180^\circ) \\
&= (-\cos A) (-\cos B) + \sin A (-\sin B) \\
&= \cos A \cos B - \sin A \sin B.
\end{aligned}$$

3. *Formulae for  $\sin (A - B)$  and  $\cos (A - B)$  in a more general case. (Generalization of Art. 34.)*



Here  $XOZ$  measured counter-clockwise is  $A$  and  $ZOP$  measured clockwise has magnitude  $B$  so that  $XOP$  measured

clockwise is  $A - B$ . From  $P$ ,  $PN$  and  $PT$  are drawn perpendiculars on  $OX$  and  $OZ$  (produced in this figure), and from  $T$ ,  $TM$  and  $TR$  are drawn perpendiculars on  $OX$  and  $PN$ .

In the present figure, magnitudes of the acute angles  $TOM$  and  $POT$  are  $180^\circ - A$  and  $B - 180^\circ$  respectively, and noting that  $PNOT$  is a cyclic quadrilateral ( $\angle^o N$  and  $T$  being right angles),  $\angle RPT = \angle TOM = 180^\circ - A$  in magnitude.

Now, we see that in considering  $\sin(A - B)$  and  $\cos(A - B)$  from the triangle  $NOP$ ,  $PN$  and  $ON$  are of negative sign.

Hence,

$$\begin{aligned}\sin(A - B) &= \frac{PN}{OP} \\ &= -\frac{MT + PR}{OP},\end{aligned}$$

where the magnitudes of  $MT$ ,  $PR$ , etc. only are considered,

$$\begin{aligned}&= -\frac{MT \cdot OT}{OT \cdot OP} - \frac{PR \cdot PT}{PT \cdot OP} \\ &= -\sin TOM \cos POT - \cos RPT \sin POT \\ &= -\sin(180^\circ - A) \cos(B - 180^\circ) \\ &\quad - \cos(180^\circ - A) \sin(B - 180^\circ) \\ &= -\sin A(-\cos B) - (-\cos A)(-\sin B) \\ &= \sin A \cos B - \cos A \sin B.\end{aligned}$$

Similarly,

$$\begin{aligned}\cos(A - B) &= \frac{ON}{OP} \quad [\text{where } ON \text{ is taken with proper sign}] \\ &= -\frac{RT - OM}{OP} \quad [\text{where magnitudes only of } RT, OM \text{ etc. are considered}] \\ &= -\frac{RT \cdot PT}{PT \cdot OP} + \frac{OM \cdot OT}{OT \cdot OP} \\ &= -\sin RPT \sin POT + \cos TOM \cos POT \\ &= -\sin(180^\circ - A) \sin(B - 180^\circ) \\ &\quad + \cos(180^\circ - A) \cos(B - 180^\circ)\end{aligned}$$

$$\begin{aligned}
 &= -\sin A (-\sin B) + (-\cos A)(-\cos B) \\
 &= \cos A \cos B + \sin A \sin B.
 \end{aligned}$$

4. A few particular cases of  $\sin(A \pm B)$ ,  $\cos(A \pm B)$ .

*Case I.* In the case  $A$  and  $B$  are both acute and  $(A+B) > 90^\circ$ .

Construction same as in Art. 33. Here  $Q$ , the foot of the perpendicular will fall on  $XO$  produced.

$$\angle TPR = 90^\circ - \angle TRP = \angle TRO = \angle ROS = A.$$

$$\sin(A+B) = \sin XOP$$


$$= \frac{PQ}{OP} = \frac{QT+TP}{OP}$$

$$= \frac{RS+PT}{OP} = \frac{RS}{OP} + \frac{PT}{OP}$$

$$= \frac{RS}{OR} \cdot \frac{OR}{OP} + \frac{PT}{PR} \cdot \frac{PR}{OP}$$

$$= \sin A \cos B + \cos TPR \sin B.$$

$$= \sin A \cos B + \cos A \sin B.$$



$$\cos(A+B) = \cos XOP = -\frac{OQ}{OP} \quad [\text{Magnitude of } OQ \text{ being considered}]$$

$$= -\frac{SQ-SO}{OP} = -\frac{OS}{OP} - \frac{SQ}{OP} = -\frac{OS}{OP} - \frac{TR}{OP}$$




# INTERMEDIATE TRIGONOMETRY

*Case III. In the case  $A$  and  $B$  are both obtuse and  $(A - B)$  is acute.*

Construction same as in Art. 34.

Here  $\angle TPR = \angle ROS = 180^\circ - A$ .



$$\begin{aligned}
 \sin (A - B) &= \sin POQ \\
 &= \frac{PQ}{OP} = \frac{PT - RS}{OP} \\
 &= \frac{PT}{OP} - \frac{RS}{OP} = \frac{PT}{PR} \cdot \frac{PR}{OP} - \frac{RS}{OR} \cdot \frac{OR}{OP} \\
 &= \cos TPR \sin POR - \sin ROS \cos POR \\
 &= \cos (180^\circ - A) \sin (180^\circ - B) \\
 &\quad - \sin (180^\circ - A) \cos (180^\circ - B) \\
 &= -\cos A \sin B - \sin A (-\cos B) \\
 &= \sin A \cos B - \cos A \sin B.
 \end{aligned}$$

$$\begin{aligned}
 \cos (A - B) &= \cos POQ \\
 &= \frac{OQ}{OP} = \frac{OS + SQ}{OP} = \frac{OS + RT}{OP} = \frac{OS}{OP} + \frac{RT}{OP} \\
 &= \frac{OS}{OR} \cdot \frac{OR}{OP} + \frac{RT}{PR} \cdot \frac{PR}{OP} \\
 &= \cos ROS \cdot \cos POR + \sin TPR \cdot \sin POR \\
 &= \cos (180^\circ - A) \cos (180^\circ - B) \\
 &\quad + \sin (180^\circ - A) \sin (180^\circ - B)
 \end{aligned}$$

$$\begin{aligned}
 &= (-\cos A)(-\cos B) + \sin A \sin B \\
 &= \cos A \cos B + \sin A \sin B.
 \end{aligned}$$

**Note.** Other particular cases of the above four formulæ can easily be proved exactly in the same way by drawing the corresponding figures in each case and making the same constructions as in Arts. 33 and 34 for  $(A+B)$  and  $(A-B)$  respectively.

**5. An alternative method of proof for  $\sin(A \pm B)$ ,  $\cos(A \pm B)$ .** [ See Arts. 33, 34 ]




Fig. (i)




Fig. (ii)

Let  $\angle XOY = A$ ;  $\angle YOZ = B$ ; in Fig. (i),  $\angle XOZ = A+B (< 90^\circ)$ ; in Fig. (ii),  $\angle XOZ = A-B (A > B)$  [  $A$  and  $B$  being positive and acute ].

Through any point  $P$  on  $OY$ , the common arm of two angles, draw a straight line  $MN$  perpendicular to  $OY$ , meeting  $OX$  in  $M$  and  $OZ$  in  $N$ .

$$\text{Then} \quad \triangle MON = \triangle MOP \pm \triangle NOP$$

$$\therefore \frac{1}{2} OM \cdot ON \sin(A \pm B) = \frac{1}{2} OM \cdot OP \sin A \pm \frac{1}{2} ON \cdot OP \sin B \quad [\text{Art. 88(i)}]$$

$$\therefore \sin(A \pm B) = \sin A \cdot \frac{OP}{ON} \pm \frac{OP}{OM} \sin B$$

$$= \sin A \cos B \pm \cos A \sin B$$

$$\cos(A \pm B) = \cos MON = \frac{OM^2 + ON^2 - MN^2}{2OM \cdot ON} \quad [\text{Art. 83}]$$

$$= \frac{(OP^2 + PM^2) + (OP^2 + PN^2) - (MP \pm PN)^2}{2OM \cdot ON}$$

$$\begin{aligned}
 &= \frac{OP^2 \mp MP \cdot PN}{OM \cdot ON} \\
 &= \frac{OP}{OM} \cdot \frac{OP}{ON} \mp \frac{MP}{OM} \cdot \frac{PN}{ON} \\
 &= \cos A \cos B \mp \sin A \sin B.
 \end{aligned}$$

### 6. Geometrical proof of the expansion of $\tan (A+B)$ .

The figure and the construction are the same as in Art. 33.

$$\begin{aligned}
 \tan (A+B) &= \frac{PQ}{OQ} = \frac{RS+PT}{OS-TR} \\
 &= \frac{\frac{RS}{OS} + \frac{PT}{OS}}{1 - \frac{TR}{OS}} = \frac{\frac{RS}{OS} + \frac{PT}{OS}}{1 - \frac{TR}{TP} \cdot \frac{TP}{OS}}
 \end{aligned}$$

Now,  $\frac{RS}{OS} = \tan A$  and  $\frac{TR}{TP} = \tan TPR = \tan A$ .


The triangles  $ROS$ ,  $TPR$  are similar.

$$\therefore \frac{TP}{OS} = \frac{PR}{OR} = \tan B.$$

$$\therefore \tan (A+B) = \frac{\tan A + \tan B}{1 - \tan A \tan B}.$$

**Note.** Similarly the expansion of  $\tan (A-B)$  can be proved geometrically from the figure and construction of Art. 34.

### 7. Geometrical proof of the formulæ for $2A$ .



Let  $XPY$  be a semi-circle,  $XY$  the diameter and  $O$  the centre.

Draw  $PN$  perpendicular to  $XY$ .



Let  $\angle PXY = A$ ; then  $\angle PCY = 2A$ .  
 $\angle NPY = 90^\circ - \angle PYN = \angle PXY = A$ .

$$\begin{aligned}\sin 2A &= \frac{PN}{CP} = \frac{2PN}{2CP} = \frac{2PN}{XY} = 2 \frac{PN}{XP} \frac{XP}{XY} \\ &= 2 \sin PYN \cdot \cos PXY = 2 \sin A \cos A.\end{aligned}$$

$$\begin{aligned}\cos 2A &= \frac{CN}{CP} = \frac{2CN}{2CP} = \frac{2CN}{XY} = \frac{CN + CN}{XY} \\ &= \frac{(XN - XC) + (CY - YN)}{XY} = \frac{XN - YN}{XY} \\ &= \frac{XN}{XP} \frac{XP}{XY} - \frac{YN}{PY} \frac{PY}{XY} \\ &= \cos A \cdot \cos A - \sin A \cdot \sin A \\ &= \cos^2 A - \sin^2 A.\end{aligned}$$

$$\begin{aligned}\tan 2A &= \frac{PN}{CN} = \frac{2PN}{2CN} = \frac{2PN}{XN - YN} \\ &= \frac{2 \frac{PN}{XN}}{1 - \frac{YN}{XN}} = \frac{2 \frac{PN}{XN}}{1 - \frac{YN}{PN} \frac{PN}{XN}} \\ &= \frac{2 \tan A}{1 - \tan A \cdot \tan A} = \frac{2 \tan A}{1 - \tan^2 A}.\end{aligned}$$

### 8. Trigonometrical Ratios of Generalised angle defined by Projections.





Let  $OXO'$  and  $YOY'$  be a pair of rectangular axes intersecting at the point  $O$  and let an angle  $\theta$ , of any magnitude (positive or negative) be generated by the revolution of  $OP$  from its initial position  $OX$  to its present position. Then the trigonometrical ratios of the generalised angle  $\theta$  are defined as follows

$$\sin \theta = \frac{\text{projection of } OP \text{ on } y\text{-axis}}{OP}$$

$$\cos \theta = \frac{\text{projection of } OP \text{ on } x\text{-axis}}{OP}$$

$$\tan \theta = \frac{\text{projection of } OP \text{ on } y\text{-axis}}{\text{projection of } OP \text{ on } x\text{-axis}}$$

$$\operatorname{cosec} \theta = \frac{OP}{\text{projection of } OP \text{ on } y\text{-axis}}$$

$$\sec \theta = \frac{OP}{\text{projection of } OP \text{ on } x\text{-axis}}$$


$$\cot \theta = \frac{\text{projection of } OP \text{ on } x\text{-axis}}{\text{projection of } OP \text{ on } y\text{-axis}}$$

In the above definitions, projection means algebraic projection; that is, we should consider not only the magnitude but also the sign of the projection; and with the usual convention the projection would be considered *positive* if they are along  $OX$  and  $OY$  and considered *negative* if they are along  $OX'$  and  $OY'$ . By convention,  $OP$  is *always considered positive*. From these definitions, the signs of the trigonometrical ratios can be easily determined according to the position of  $OP$  in one or other of the four quadrants. In the above figures, the positions of  $OP$  in two quadrants only (1st and 3rd) are shown.

**Note 1.** From the above definitions, it is clear that if  $OX$  be a fixed line, and if,  $l$  be the length of any straight line  $OP$  inclined at an angle  $\theta$  to  $OX$ , then the *projection of  $OP$  along  $OX$*  is  $l \cos \theta$ , whatever be the magnitude of the angle  $\theta$ .

**Note 2.** The *Addition and Subtraction Theorems* for generalised angles can also be proved by the method of projection.

## 9. Two important Trigonometrical relations.



If  $D$  be any point in the base  $BC$  of a triangle  $ABC$ , and if  $AD$  divides  $BC$  into two parts  $m$  and  $n$  ( $BD=m$ ,  $CD=n$ ) and the angle  $A$  into two parts  $\alpha$  and  $\beta$  ( $\angle BAD=\alpha$ ,  $\angle CAD=\beta$ ), and if the angle  $ADB$  be  $\theta$ , then

$$(i) (m+n) \cot \theta = n \cot \beta - m \cot \alpha$$

$$(ii) (m+n) \cot \theta = m \cot C - n \cot B.$$

We have

$$\begin{aligned} \frac{m}{n} &= \frac{BD}{DC} = \frac{BD \cdot AD}{AD \cdot DC} = \frac{\sin BAD \cdot \sin ACD}{\sin ABD \cdot \sin DAC} \\ &= \frac{\sin \alpha}{\sin (\theta + \alpha)} \cdot \frac{\sin (\theta - \beta)}{\sin \beta} \left[ \because \begin{array}{l} \angle ABD = \pi - (\alpha + \theta) \\ \angle ACD = \theta - \beta \end{array} \right] \\ &= \frac{\sin \alpha (\sin \theta \cos \beta - \cos \theta \sin \beta)}{\sin \beta (\sin \theta \cos \alpha + \cos \theta \sin \alpha)} \end{aligned}$$

Dividing the numerator and the denominator by

$\sin \alpha \sin \beta \sin \theta$ , we have

$$\frac{m}{n} = \frac{\cot \beta - \cot \theta}{\cot \alpha + \cot \theta}.$$

$$\therefore (m+n) \cot \theta = n \cot \beta - m \cot \alpha.$$

Again,

$$\frac{m}{n} = \frac{\sin BAD \sin ACD}{\sin ABD \sin DAC}$$

$$= \frac{\sin(\theta + B)}{\sin B} \cdot \frac{\sin C}{\sin(\theta - C)} \quad \left[ \begin{array}{l} \angle BAD = \pi - (\theta + B) \\ \angle DAC = \theta - C \end{array} \right]$$

$$= \frac{\sin C (\sin \theta \cos B + \cos \theta \sin B)}{\sin B (\sin \theta \cos C - \cos \theta \sin C)}$$

Dividing the numerator and the denominator by

$\sin B \sin C \sin \theta$ , we have

$$\frac{m}{n} = \frac{\cot B + \cot \theta}{\cot C - \cot \theta}$$

$$\therefore (m+n) \cot \theta = m \cot C - n \cot B.$$

#### 10. Note of Art. 90.

Let us denote the formulæ of Arts. 82, 83, 84 by (I), (II), (III). We have seen in Art. 90, that (II) can be deduced from (III). We shall now show how any one of these can be deduced from any other of the rest.

*To deduce (I) from (III).*

Substituting value of  $b$  from the second relation of Art. 84 in the first,

$$a = (c \cos A + a \cos C) \cos C + c \cos B.$$

$$\therefore a(1 - \cos^2 C) = c(\cos A \cos C + \cos B)$$

$$= c\{\cos A \cos C - \cos(A+C)\}$$

$$[\because A+B+C=\pi]$$

$$\therefore a \sin^2 C = c \sin A \sin C. \quad \therefore a/\sin A = c/\sin C.$$

Similarly substituting the value of  $c$  in the first relation, we get

$$a/\sin A = b/\sin B. \quad \text{Hence etc.}$$

*To deduce (II) and (III) from (I)*

(i) Putting each of the ratios of Art. 82 equal to  $k$ , we get

$$a = k \sin A; \quad b = k \sin B; \quad c = k \sin C.$$

$$\therefore \frac{b^2 + c^2 - a^2}{2bc} = \frac{k^2(\sin^2 B + \sin^2 C - \sin^2 A)}{k^2 \cdot 2 \sin B \sin C}$$

$$= \frac{\sin^2 B + \sin(C+A) \sin(C-A)}{2 \sin B \sin C}$$

$$\begin{aligned}
&= \frac{\sin B \{\sin B + \sin (C - A)\}}{2 \sin B \sin C} \\
&\quad [\because \sin (C + A) = \sin (\pi - B) = \sin B] \\
&= \frac{\sin B \{\sin (C + A) + \sin (C - A)\}}{2 \sin B \sin C} \\
&= \frac{2 \sin B \sin C \cos A}{2 \sin B \sin C} = \cos A.
\end{aligned}$$

$$\begin{aligned}
\text{(ii) } b \cos C + c \cos B &= k (\sin B \cos C + \sin C \cos B) \\
&= k \sin (B + C) = k \sin A \\
&= a.
\end{aligned}$$

$$[\because A + B + C = \pi]$$

To deduce (I) and (III) from (II)

$$\begin{aligned}
\text{(i) } \sin^2 A &= 1 - \cos^2 A \\
&= 1 - \left( \frac{b^2 + c^2 - a^2}{2bc} \right)^2 = \frac{4b^2c^2 - (b^2 + c^2 - a^2)^2}{4b^2c^2} \\
&= \frac{(2bc + b^2 + c^2 - a^2)(2bc - b^2 - c^2 + a^2)}{4b^2c^2} \\
&= \frac{(a + b + c)(b + c - a)(c + a - b)(a + b - c)}{4b^2c^2} \\
&= \frac{K}{4b^2c^2} \text{ say.}
\end{aligned}$$

$$\therefore \frac{\sin^2 A}{a^2} = \frac{K}{4a^2b^2c^2};$$

$$\text{similarly, } \frac{\sin^2 B}{b^2} \text{ and } \frac{\sin^2 C}{c^2} \text{ each} = \frac{K}{4a^2b^2c^2}.$$

$$\therefore \frac{\sin^2 A}{a^2} = \frac{\sin^2 B}{b^2} = \frac{\sin^2 C}{c^2}; \text{ hence etc.}$$

(ii) Adding 2nd and 3rd relations of the formulæ of Art. 83, we get

$$b^2 + c^2 = b^2 + c^2 + 2a^2 - 2ca \cos B - 2ab \cos C.$$

Now transposing and dividing by  $2a$ , we get

$$a = b \cos C + c \cos B; \text{ etc.}$$

### Miscellaneous Examples III

1. The angles of a triangle are as 4 : 5 : 6. Express them in circular measure.

2. The angles of a triangle are in A. P. and the greatest is double the least; express the angles in degrees, and in radians.

3. The number of degrees in one of the acute angles of a right-angled triangle is three-tenths of the number of grades in the other; determine the angles in degrees.

4. Compare the areas of two circles in which the circumference of one is equal to an arc of  $60^\circ$  of the other.

5. A railway train is travelling on a curve of half-a-mile radius at the rate of 20 miles an hour; through what angle has it turned in 10 seconds?

6. An arc of a circle whose radius is 7 inches, subtends an angle of  $15^\circ 39' 7''$ ; what angle will an arc of the same length subtend in a circle whose radius is 2 inches?

Prove the following identities (*Ex. 7 to 22*):—

$$7. \sin^2 \theta \tan \theta + \cos^2 \theta \cot \theta + 2 \sin \theta \cos \theta = \tan \theta + \cot \theta.$$

$$8. \sin^2 \theta (1 + \cot^2 \theta) + \cos^2 \theta (1 + \tan^2 \theta) = 2.$$

$$9. (\tan \theta + \sec \theta)^2 = \frac{1 + \sin \theta}{1 - \sin \theta}. \quad [C. U. 1934.]$$

$$10. 2(\sin^6 \theta + \cos^6 \theta) - 3(\sin^4 \theta + \cos^4 \theta) + 1 = 0.$$

$$11. \frac{\tan x - \cot y}{\tan y - \cot x} = \tan x \cot y.$$

$$12. (\sin x \cos y + \cos x \sin y)^2 + (\cos x \cos y - \sin x \sin y)^2 = 1.$$

13.  $\sin^4 x + \cos^4 x = 1 - 2 \sin^2 x \cos^2 x.$
14.  $\sin^6 x - \cos^6 x = (\sin^2 x - \cos^2 x)(1 - 2 \sin^2 x \cos^2 x).$
15.  $\sqrt{\frac{1 - \sin \theta}{1 + \sin \theta}} = \sec \theta - \tan \theta.$
16.  $(1 + \sec \theta + \tan \theta)(1 - \sec \theta + \tan \theta) = 2 \tan \theta.$
17.  $\frac{\cos \theta}{1 - \tan \theta} + \frac{\sin \theta}{1 - \cot \theta} = \sin \theta + \cos \theta.$
18.  $(\sin x + \cos x)^2 + (\sin x - \cos x)^2 = 2.$
19.  $\cot^2 x \cdot \frac{\sec x - 1}{1 + \sin x} + \sec^2 x \cdot \frac{\sin x - 1}{1 + \sec x} = 0.$
20.  $(\sin x + \cos x)(\tan x + \cot x) = \sec x + \operatorname{cosec} x.$
21.  $(\sin \theta + \sec \theta)^2 + (\cos \theta + \operatorname{cosec} \theta)^2 = (1 + \sec \theta \operatorname{cosec} \theta)^2.$
22.  $\frac{1 - \sin \theta \cos \theta}{\cos \theta (\sec \theta - \operatorname{cosec} \theta)} \cdot \frac{\sin^2 \theta - \cos^2 \theta}{\sin^3 \theta + \cos^3 \theta} = \sin \theta.$
23. If  $a \cos^2 x + b \sin^2 x = c$ , show that  $\tan x = \pm \sqrt{\frac{c-a}{b-a}}.$
24. If  $\operatorname{cosec} A + \operatorname{cosec} B + \operatorname{cosec} C = 0$ , show that  $(\Sigma \sin A)^2 = \Sigma \sin^2 A.$
25. If  $x = a \cos \theta + b \sin \theta$  and  $y = a \sin \theta - b \cos \theta$ , show that  $x^2 + y^2 = a^2 + b^2.$
26. Express  $\frac{\sin x}{\cos^3 x} + \frac{\cos x}{\sin^3 x}$  in terms of  $t$ , where  $t$  stands for  $\tan x.$
27. If  $\sin A = \frac{1}{2}$  and  $\tan B = \sqrt{3}$ , find the value of  $\sin A \cos B + \cos A \sin B.$
28. If  $\cos \theta = \frac{1}{2}$ , find the value of  $\frac{1 - \tan^2 \theta}{1 + \tan^2 \theta}.$
29. If  $5 \tan \theta = 4$ , find the value of  $\frac{5 \sin \theta - 3 \cos \theta}{\sin \theta + 2 \cos \theta}.$

30. If  $\frac{\sin x}{\sin y} = \sqrt{2}$ ,  $\frac{\tan x}{\tan y} = \sqrt{3}$ ,

find  $x$  and  $y$  (given  $x$  and  $y$  acute angles).

31. Which of the statements is possible and which impossible,  $x$ ,  $y$  and  $z$  being unequal real quantities ?

(i)  $\operatorname{cosec} \theta = \frac{x^2 + y^2}{2xy}$ .      (ii)  $\sec \theta = \frac{x^2 - y^2}{x^2 + y^2}$ .

(iii)  $\sin \theta = \frac{x^2 + y^2 + z^2}{yz + zx + xy}$ .

32. Eliminate  $\theta$  from the equations

(i)  $\sin \theta + \cos \theta = a$ ,  $\sin \theta - \cos \theta = b$ .

(ii)  $\frac{x}{a} \cos \theta + \frac{y}{b} \sin \theta = 1$ ,  $\frac{x}{a} \sin \theta - \frac{y}{b} \cos \theta = 1$ .

(iii)  $x = a \cos^3 \theta$ ,  $y = b \sin^3 \theta$ .

33. If  $k \tan \theta = \tan k\theta$ , prove that

$$\frac{\sin^3 k\theta}{\sin^3 \theta} = \frac{k^3}{1 + (k^2 - 1) \sin^2 \theta}.$$

34. If  $\sec x \sec y + \tan x \tan y = \sec z$ ,

then,  $\sec x \tan y + \tan x \sec y = \pm \tan z$ .

35. Show that  $\left( \frac{1 + \cot 60^\circ}{1 - \cot 60^\circ} \right)^2 = \frac{1 + \cos 30^\circ}{1 - \cos 30^\circ}$

36. If  $\tan x = \frac{\sin \theta - \cos \theta}{\sin \theta + \cos \theta}$ , prove that

$$\sin x = \frac{1}{\sqrt{2}} (\sin \theta - \cos \theta)$$

37. Show that the product of  $\sin x(1 + \sin x) + \cos x(1 + \cos x)$  and  $\sin x(1 - \sin x) + \cos x(1 - \cos x)$  is equal to  $2 \sin x \cos x$ .

38. Find the height of a chimney when it is found that on walking towards it 250 feet, in a horizontal line through its base, the angular elevation changes from  $45^\circ$  to  $75^\circ$ .

39. The length of a kite string is 250 yards, and the angle of elevation of the kite is  $30^\circ$ . Find the height of the kite.

40. The angle of elevation of the top of a temple at a distance 300 feet is  $30^\circ$ ; find its height.

41. Find the angle of elevation of the sun when the shadow of a pole 60 feet high, is  $20\sqrt{3}$  yards long.

42. The angles of elevation of a tower at two places due east of it and 200 feet apart are  $45^\circ$  and  $30^\circ$ ; find the height of the tower.

43. An aeroplane leaves the earth at the point  $X$  and rises at a uniform speed. At the end of 15 seconds, an observer stationed at a distance of 660 feet from  $X$ , finds the angular elevation of the aeroplane to be  $45^\circ$ ; at what rate in miles per hour is the aeroplane rising?

44. A ladder 45 feet long just reaches the top of a wall. The ladder makes an angle of  $60^\circ$  with the wall. Find the height of the wall and the distance of the foot of the ladder from the wall.

45. If  $\cos A = \frac{4}{5}$ ,  $\cos B = \frac{3}{5}$ , find the values of  $\sin(A+B)$  and  $\cos(A-B)$ .

46. If  $\tan A = \frac{5}{12}$  and  $\tan B = \frac{9}{16}$ , find the values of  $\sin(A-B)$  and  $\cos(A-B)$ .

47. If  $\tan A = \frac{m+n}{m-n}$ , and  $\tan B = \frac{m-n}{m+n}$ , find  $\tan(A-B)$ .

48. If  $\tan(x+y) = \frac{3}{4}$  and  $\tan x = \frac{5}{6}$ , find  $\tan y$ .

49. If  $\cos \theta = \frac{3}{5}$ , find  $\sin 2\theta$ ,  $\tan 2\theta$ ,  $\cos \frac{\theta}{2}$ .

50. If  $\cos x = \frac{4}{5}$ ,  $\cos y = \frac{3}{5}$  ( $x$  and  $y$  being positive acute angles), find the value of  $\cos \frac{1}{2}(x-y)$ .



51. If  $\sin A = \frac{1}{\sqrt{2}}$ ,  $\sin B = \frac{1}{\sqrt{3}}$ , find the value of

$$\tan \frac{1}{2}(A+B) \cot \frac{1}{2}(A-B).$$

52. If  $\sec x = \frac{1}{\sqrt{2}}$ ,  $\operatorname{cosec} y = \frac{5}{4}$ , find  $\sec(x+y)$ .

53. Prove that

$$\frac{2 \cos 8\theta + 1}{2 \cos \theta + 1} = (2 \cos \theta - 1)(2 \cos 2\theta - 1)(2 \cos 4\theta - 1).$$

54. Show that  $a \cos \theta + b \sin \theta = \sqrt{a^2 + b^2} \cos(\theta - \alpha)$ ,  
where  $\tan \alpha = b/a$ .

55. If  $\sin^4 x + \cos^4 x = 1$ , prove that  $x$  is zero or a multiple of  $\frac{1}{2}\pi$ .

56. If  $\sqrt{2} \cos A = \cos B + \cos^3 B$ ,

$$\text{and } \sqrt{2} \sin A = \sin B - \sin^3 B,$$

$$\text{thnn } \sin(A-B) = \pm \frac{1}{2}.$$

57. Prove that  $\cos^2(\alpha - \beta) - \sin^2(\alpha + \beta) = \cos 2\alpha \cos 2\beta$ .

58. Show that  $\sin 18^\circ + \cos 18^\circ = \sqrt{2} \cos 27^\circ$ .

59. Show that whatever be the value of  $\theta$ ,

$$\sin^2(\theta + \alpha) + \sin^2(\theta + \beta) - 2 \cos(\alpha + \beta) \sin(\theta + \alpha) \sin(\theta + \beta)$$

is independent of  $\theta$ .

60. Show that

$$(i) \frac{\sin \alpha}{\sin(\alpha - \beta) \sin(\alpha - \gamma)} + \frac{\sin \beta}{\sin(\beta - \gamma) \sin(\beta - \alpha)} \\ + \frac{\sin \gamma}{\sin(\gamma - \alpha) \sin(\gamma - \beta)} = 0.$$

$$(ii) \tan(\beta + \gamma - 2\alpha) + \tan(\gamma + \alpha - 2\beta) + \tan(\alpha + \beta - 2\gamma) \\ = \tan(\beta + \gamma - 2\alpha) \tan(\gamma + \alpha - 2\beta) \tan(\alpha + \beta - 2\gamma).$$

61. If  $\tan \frac{1}{2}\theta = \tan^2 \frac{1}{2}\phi$  and  $\tan \phi = 2 \tan \alpha$ ,  
show that  $\theta + \phi = 2\alpha$ .

62. (i) If  $\tan^2 x = 2 \tan^2 y + 1$ , then  $\cos 2x + \sin^2 y = 0$ .

(ii) If  $\cos A = \tan B$ ,  $\cos B = \tan C$ ,  $\cos C = \tan A$ ,  
prove that  $\sin A = \sin B = \sin C$ .

63. Show that  $\tan 20^\circ \tan 40^\circ \tan 80^\circ = \tan 60^\circ$ .

64. If  $\alpha + \beta + \gamma = 0$ , prove that

$$\cos \alpha + \cos \beta + \cos \gamma = 4 \cos \frac{\alpha}{2} \cos \frac{\beta}{2} \cos \frac{\gamma}{2} - 1.$$

65. If in any triangle,  $\tan \phi = \frac{a-b}{a+b} \cot \frac{1}{2}C$ , prove that

$$c = (a+b) \sin \frac{1}{2}C \sec \phi.$$

66. If  $\cos \theta = \frac{a \cos \phi - b}{a - b \cos \phi}$ , then  $\frac{\tan \frac{\theta}{2}}{\sqrt{a+b}} = \frac{\tan \frac{\phi}{2}}{\sqrt{a-b}}$ .

67. If  $\alpha + \beta + \gamma = \frac{1}{2}\pi$ , prove that

$$\sin^2 \alpha + \sin^2 \beta + \sin^2 \gamma + 2 \sin \alpha \sin \beta \sin \gamma = 1.$$

68. If  $\cos^2 A + \cos^2 B + \cos^2 C + 2 \cos A \cos B \cos C = 1$ ,  
show that one of the quantities  $A \pm B \pm C$  is an odd multiple  
of  $\pi$ .

69. Show that  $\sec x = \frac{2}{\sqrt{2} + \sqrt{2+2 \cos 4x}}$ .

70. If  $a \sin (\theta + \alpha) = b \sin (\theta + \beta)$ , prove that

$$\cot \theta = \frac{a \cos \alpha - b \cos \beta}{b \sin \beta - a \sin \alpha}.$$

71. If  $\tan \beta = \frac{n \sin \alpha \cos \alpha}{1 - n \sin^2 \alpha}$ , show that  
 $\tan (\alpha - \beta) = (1 - n) \tan \alpha$ .

In any triangle, prove that (Ex. 72 to 77) :

$$72. \quad \frac{\cos A}{c \cos B + b \cos C} + \frac{\cos B}{a \cos C + c \cos A} + \frac{\cos C}{b \cos A + a \cos B} = \frac{a^2 + b^2 + c^2}{2abc}.$$

$$73. \quad \frac{\tan \frac{A}{2}}{(a-b)(a-c)} + \frac{\tan \frac{B}{2}}{(b-c)(b-a)} + \frac{\tan \frac{C}{2}}{(c-a)(c-b)} = \frac{1}{\Delta}.$$

$$74. \quad \sin 3A \sin (B-C) + \sin 3B \sin (C-A) + \sin 3C \sin (A-B) = 0.$$

$$75. \quad \cot B + \frac{\cos C}{\sin B \cos A} = \cot C + \frac{\cos B}{\sin C \cos A}.$$

$$76. \quad c = (a-b) \sec \theta, \text{ where } \tan \theta = \frac{2\sqrt{ab}}{a-b} \sin \frac{C}{2}.$$

$$77. \quad a(\cos B \cos C + \cos A) = b(\cos C \cos A + \cos B) = c(\cos A \cos B + \cos C).$$

$$78. \quad \text{If in a triangle, } c(a+b) \cos \frac{B}{2} = b(a+c) \cos \frac{C}{2},$$

show that the triangle is isosceles.

79. If in a triangle,  $a, b, c$  be in A. P. and the greatest angle exceeds the least by  $90^\circ$ , prove that

$$a : b : c = \sqrt{7} - 1 : \sqrt{7} : \sqrt{7} + 1.$$

80. In the solution of triangles there can be no ambiguity except when an angle is determined by the sine or cosecant, and in no case whatever, when the triangle has one right angle ; prove this. [ *Cambridge.* ]

81. If  $\sin(\pi \cos \theta) = \cos(\pi \sin \theta)$ , prove that

$$\cos\left(\theta \pm \frac{\pi}{4}\right) = \frac{1}{2\sqrt{2}}.$$

82. If  $\sin(\pi \cot \theta) = \cos(\pi \tan \theta)$ , prove that either  $\operatorname{cosec} 2\theta$  or  $\cot 2\theta$  is equal to  $n + \frac{1}{2}$ ,  $n$  being an integer.

83. If  $\alpha$  and  $\beta$  be the different values of  $\theta$  which satisfy the equation  $a \cos \theta + b \sin \theta = c$ , prove that

$$\sin(\alpha + \beta) = \frac{2ab}{a^2 + b^2}.$$

84. Find all the values of  $\theta$  which satisfy the equation  $\sin \theta + \sin 2\theta + \sin 3\theta = 1 + \cos \theta + \cos 2\theta$ .

85. Prove that in any triangle,

$$\frac{1}{bc} + \frac{1}{ca} + \frac{1}{ab} = \frac{1}{2Rr}.$$

86. If  $r : R : r_1 = 2 : 5 : 12$ , show that the triangle is right-angled.

87. If the angle of elevation of a cloud observed from a point at a height  $h$  above the surface of a lake be  $\phi$  and the angle of depression of its image in the lake be  $\theta$ , prove that the height of the cloud above the lake is  $h \frac{\sin(\theta + \phi)}{\sin(\theta - \phi)}$ , assuming that the image is vertically as much below the surface as the cloud is above it.

[ *A. U. 1942 ; B. H. U. I. 1931.* ]

88. The elevation of a tower due north of a station at  $A$  is  $\alpha$  and at a station  $B$  due west of  $A$  is  $\beta$ . Prove that its altitude is  $\frac{AB \sin \alpha \sin \beta}{\sqrt{\sin^2 \alpha - \sin^2 \beta}}$ . [ *B. H. U. I. 1934.* ]

89. A man walks along a straight road and observes that the greatest angle subtended by two objects is  $\alpha$ ; from the point where this greatest angle is subtended, he walks a distance  $c$  along the road and finds that the two objects are now in a straight line which makes an angle  $\beta$  with the road. Prove that the distance between the objects is  $c \sin \alpha \sin \beta \sec \frac{\alpha + \beta}{2} \sec \frac{\alpha - \beta}{2}$ . [ *B. H. U. I. 1936.* ]

90. On the bank of a river is a column 200 ft. high supporting a statue 30 ft. high. To an observer on the opposite bank with his eye on the level of the ground the statue subtends an angle equal to that subtended by a man 6 ft. high standing at the base of the column; determine the breadth of the river. [ *B. H. U. I. 1941.* ]

## ANSWERS

### Examples I. [ Pages 11-14 ]

1. (i) first quadrant ; (ii) third quadrant ;  
(iii) second quadrant , (iv) fourth quadrant.
2. (i)  $61^{\circ} 34' 44'' 4$  ; (ii)  $175^{\circ} 49' 1'' 776$ .
3. (i)  $253775\pi$  , (ii)  $\frac{3}{10}\pi$ .
4.  $82^{\circ} 30'$  ;  $91^{\circ} 66' 6'' 6$  ,  $1\frac{1}{4}\pi$ . 5.  $\alpha : \beta = 5\pi : 24$ .
6.  $\frac{1}{2} \left( 1 - \frac{\pi}{180} \right)$ . 7.  $6^{\circ}$  and  $9^{\circ}$ . 8.  $\frac{1}{90} \left( D + \frac{M}{60} \right) - \frac{1}{100} \left( G + \frac{m}{100} \right)$ .
9.  $115^{\circ}$  nearly 10.  $20^{\circ}$  and  $30^{\circ}$ . 12.  $20^{\circ}$ ,  $40^{\circ}$ ,  $80^{\circ}$ .
13.  $27^{\circ}$ ,  $9^{\circ}$ ,  $18^{\circ}$ . 14. (i) At  $28\frac{1}{4}$  min. and 48 min. past 7 ;  
(ii) At 7-10. 15.  $20^{\circ}$ ,  $60^{\circ}$ ,  $100^{\circ}$ . 16.  $\frac{\pi}{7}$ ,  $\frac{2\pi}{7}$ ,  $\frac{4\pi}{7}$  ;  $\frac{\pi}{21}$ ,  $\frac{4\pi}{21}$ ,  $\frac{16\pi}{21}$ .
17.  $45^{\circ}$ ,  $60^{\circ}$ ,  $120^{\circ}$ ,  $135^{\circ}$ . 18. 9.
19.  $m\pi$  and  $n\pi$  where  $x = \frac{2(10pm - 9qn)}{mn(10p - 9q)}$ . 20. 2.
21. 3 and 6. 22. 51.41 miles per hour (nearly).
23. 66444 miles per hour (nearly) ; 431445 miles (nearly).
24. 70.8 ft. (nearly). 25. 3959 miles (nearly). 26. 83 ft. 27. 360 yds.

### Examples II. [ Pages 24-26 ]

25.  $(\sin \theta - \cos \theta)^2$ . 26.  $\frac{1}{\tan^2 \theta} - \tan^2 \theta$ . 31.  $\frac{a^2 - b^2}{a^2 + b^2}$ .
33.  $\pm \frac{\sqrt{\sec^2 \alpha - 1}}{\sec \alpha}$  ;  $\pm \frac{\sqrt{1 + \cot^2 \theta}}{\cot \theta}$ . 34.  $\frac{1}{2}$ . 36.  $\frac{1}{2}$ . 37. 1 or  $\frac{1}{2}$ .
39.  $\frac{a^2 - b^2}{2ab}$  ;  $\frac{a^2 + b^2}{a^2 - b^2}$ . 43. (i)  $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$  ; (ii)  $xy = c^2$  ;  
(iii)  $(bc' - b'o)^2 + (ca' - c'a)^2 = (ab' - a'b)^2$ .  
(iv)  $(a'b - b'o)(ab^2 - bc') = (aa' - ac')^2$ .

**Examples III.** [ Pages 85-86 ]

7.  $\frac{\sqrt{3}}{2}$ . 8. (i)  $60^\circ$ ; (ii)  $45^\circ$ ; (iii)  $30^\circ$  (There is another angle which is not one of the standard angles).  
 (iv)  $45^\circ$ ; (v)  $30^\circ$ ; (vi)  $30^\circ$ ; (vii)  $30^\circ$ .  
 9.  $\theta = 52\frac{1}{2}^\circ$ ,  $\phi = 7\frac{1}{2}^\circ$ . 10.  $\alpha = 50^\circ$ ,  $\beta = 10^\circ$ .  
 11.  $A = 22\frac{1}{2}^\circ$ ,  $B = 67\frac{1}{2}^\circ$ ,  $C = 45^\circ$ . 12. (i)  $-\frac{1}{3}$ ; (ii) 1.

**Examples IV.** [ Pages 49-51 ]

1.  $\frac{1}{2}$ ;  $-\frac{1}{\sqrt{3}}$ ;  $\frac{2}{\sqrt{3}}$ ;  $-1$ . 2.  $-\frac{1}{\sqrt{2}}$ ;  $-\frac{2}{\sqrt{3}}$ ;  $-\frac{1}{\sqrt{3}}$ ;  $\frac{\sqrt{3}}{2}$ . 3. 0.  
 4.  $\frac{\sqrt{3}}{2}$ . 5. (i) 1; (ii)  $\pm 2$ ,  $\pm \frac{2}{\sqrt{3}}$ . 10.  $\tan^2 \theta$ ; 1. 12. (i) 2.  
 (ii) 1. (iii)  $\sin x$  or 0 according as  $x$  is odd or even. 13.  $\frac{1}{2}$ .  
 14.  $\frac{\sqrt{40}}{8}$ . 15. (i)  $\cot 26^\circ$ ; (ii)  $\cos 25^\circ$ ; (iii)  $\operatorname{cosec} 39^\circ$ ; (iv)  $\cos \frac{\pi}{9}$ .  
 16. (i)  $300^\circ$ ; (ii)  $480^\circ$ . 17. (i)  $60^\circ$ ; (ii)  $120^\circ$ ,  $240^\circ$ ;  
 (iii)  $30^\circ$ ,  $150^\circ$ ,  $210^\circ$ ,  $330^\circ$ ; (iv)  $30^\circ$ ,  $150^\circ$ ; (v)  $30^\circ$ ,  $135^\circ$ ,  $150^\circ$ ,  $315^\circ$ .

**Examples V.** [ Pages 56-59 ]

1.  $100\sqrt{3}$  ft. 2.  $2'89\ldots$  miles;  $2\frac{1}{2}$  miles. 3.  $20\sqrt{3}$  ft.; 120 ft.  
 4.  $20\sqrt{3}$  ft.; 20 ft. 5.  $30\sqrt{2}$  ft. 6.  $400(\sqrt{3}+1)$  yds.  
 7.  $40\sqrt{3}$  ft. 8.  $\frac{1}{2}(3 \pm \sqrt{3})$  miles. 9.  $22'8$  miles nearly.  
 10.  $94'64$  ft. nearly. 11.  $47'82$  ft. nearly. 12. 60 miles per hour.  
 13.  $50\sqrt{6}$  ft. 14.  $40\sqrt{6}$  ft.;  $40\sqrt{2}(\sqrt{7}+1)$  ft.  
 15.  $\frac{1}{2}(\sqrt{3}+1)$  miles. 16.  $5\sqrt{13}$  miles..  
 17.  $241'6\ldots$  ft.;  $91'6\ldots$  ft. 18.  $5'25\ldots$  miles per hour.  
 19.  $867'38$  ft. 20.  $\frac{1}{2}\sqrt{6}(\sqrt{5}+1)$ .  
 22. 2 miles. 23.  $18'66$  ft.

**Examples VI.** [ Pages 68-70 ]

21.  $\sin A \cos B \cos C - \sin B \cos C \cos A + \sin C \cos A \cos B$   
 $+ \sin A \sin B \sin C$ ;  

$$\frac{\tan A - \tan B - \tan C - \tan A \tan B \tan C}{1 + \tan A \tan B + \tan A \tan C + \tan B \tan C}$$
  
 22. 
$$\frac{\cot A \cot B \cot C - \cot A - \cot B - \cot C}{\cot B \cot C + \cot C \cot A + \cot A \cot B - 1}$$

**Examples VIII.** [ Pages 79-81 ]

27. a.

## Examples IX. [Pages 86-87]

16.  $\frac{b^3 - a^3}{b^3 + a^3}$ . 17. (i)  $2 \sin \frac{1}{2}A = \sqrt{1 + \sin A} + \sqrt{1 - \sin A}$ ;

(ii) No;  $2 \sin \frac{1}{2}\theta = \sqrt{1 + \sin \theta} + \sqrt{1 - \sin \theta}$ .

## Examples XI. [Pages 110-111]

1.  $n\pi \pm \frac{\pi}{4}$ , i.e.  $(2k+1)\frac{\pi}{4}$ . 2. (i)  $n\pi \pm \frac{\pi}{4}$ ; (ii)  $n\pi \pm \frac{\pi}{3}$ .  
 3.  $2n\pi \pm \frac{\pi}{3}$ ,  $(2k+1)\pi$ . 4.  $\frac{n\pi}{2} + (-1)^n \frac{\pi}{12}$ .  
 5.  $n\pi + \frac{\pi}{4}$ , or,  $n\pi + (-1)^n \frac{\pi}{6}$ . 6.  $\frac{n\pi}{3}$ , or,  $n\pi \pm \frac{\pi}{6}$ .  
 7.  $\frac{n\pi}{m+(-1)^n}$ . 8.  $(2n+1)\frac{\pi}{2}$ , or,  $(2n+1)\frac{\pi}{4}$ , or,  $(2n+1)\frac{\pi}{8}$ .  
 9.  $n\pi - \frac{\pi}{4}$ , or,  $\frac{n\pi}{2} + (-1)^n \frac{\pi}{2}$ , where  $\sin \alpha = \frac{\sqrt{5}-1}{2}$ . 10.  $\frac{1}{4}\pi$ .  
 11.  $n\pi + \frac{\pi}{4}$ . 12.  $(4n+1)\frac{\pi}{8}$ . 13.  $2n\pi + \frac{5\pi}{12}$ , or,  $2n\pi - \frac{\pi}{12}$ .  
 14.  $(2n+1)\frac{\pi}{4}$ , or,  $n\pi \pm \frac{\pi}{6}$ . 15.  $2n\pi + \frac{\pi}{2}$ , or,  $2n\pi - \beta$ , where  $\beta$  is a positive acute angle whose sine is  $\frac{1}{2}$ . 16.  $\frac{1}{2}n\pi$ . 17.  $n\pi \pm \frac{1}{6}\pi$ .  
 18.  $(4n+1)\frac{\pi}{12}$ . 19.  $2n\pi + \frac{1}{12}\pi$ , or  $2n\pi + \frac{1}{12}\pi$ . 20.  $-\frac{7}{12}\pi$ ,  $-\frac{1}{6}\pi$ ,  $\frac{1}{3}\pi$ ,  $\frac{11}{12}\pi$ .  
 $\frac{1}{2}(n\pi + \alpha)$ , where  $\tan \alpha = 2$ . 22.  $2n\pi$ . 23.  $2n\pi$ ,  $\frac{1}{6}(4n+1)\pi$ .  
 24.  $90^\circ, 450^\circ, 810^\circ$ . 25.  $\frac{1}{4}\pi, \frac{3}{4}\pi$ . 26. (i)  $\frac{1}{2}n\pi + \frac{1}{4}\pi$ ;  $2n\pi \pm \frac{3}{8}\pi$ .  
 (ii)  $0, \pm \frac{\pi}{12}, \pm \frac{\pi}{6}, \pm \frac{\pi}{4}$ . (iii)  $\frac{n\pi}{3}$ ;  $n\pi \pm \tan^{-1} \frac{1}{\sqrt{2}}$ . (iv)  $2n\pi - \alpha$ ,  $\frac{4n-1}{2}\pi + \alpha$ .  
 (v)  $2n\pi$ , or,  $2n\pi - \frac{1}{2}\pi$ . (vi)  $(2n+1)\frac{\pi}{12}$ ,  $\frac{4n+1}{14}\pi$ ,  $\frac{4n-1}{6}\pi$ .  
 (vii)  $n\pi + \frac{\pi}{2}$ ;  $(2n+1)\frac{\pi}{6} - \frac{\pi}{6}$ . 28.  $n\pi + (-1)^n 21^\circ 48' - 68^\circ 12'$ .  
 29. (i)  $\alpha = \beta = \frac{1}{2}\pi$ ; or,  $\alpha = \frac{3}{2}\pi$ ,  $\beta = -\frac{1}{2}\pi$ .  
 (ii)  $\alpha = \frac{1}{2}\pi$ ,  $\beta = \frac{1}{2}\pi$ ; or,  $\alpha = \frac{3}{2}\pi$ ,  $\beta = \frac{3}{2}\pi$ ;  
 or,  $\alpha = \frac{1}{2}\pi$ ,  $\beta = \frac{3}{2}\pi$ ; or,  $\alpha = \frac{3}{2}\pi$ ,  $\beta = -\frac{1}{2}\pi$ .

## Examples XII. [Pages 119-121]

22. (i) 1; (ii)  $\infty$ ; (iii)  $\frac{x+y}{1-xy}$ . 23.  $y = \frac{4x(1-x^2)}{1-6x^2+x^4}$ .  
 24.  $(x-y)(1+yz) = (y-z)(1+xy)$ . 25. (i)  $\frac{1}{2}$ , or,  $-8$ ; (ii)  $\frac{a-b}{1+ab}$ ;

- (iii)  $\pm \frac{\sqrt{5}}{3}$ ; (iv)  $\pm \frac{1}{\sqrt{3}}$ ; (v)  $\frac{1}{2}$ , or,  $-\frac{1}{2}$ ; (vi)  $\pm \sqrt[4]{21}$ ;  
 (vii) 0, or,  $\frac{1}{2}$ ; (viii) 0,  $\pm \frac{1}{2}$ ; (ix)  $2 - \sqrt{3}$ ; (x)  $\frac{6 + \sqrt{6}}{8}$ .

### Miscellaneous Examples I. [ Pages 122-123 ]

2.  $\pm \sqrt{\frac{b^2 - c^2}{a^2 - c^2}}$ . 19.  $a^2 + b^2 = 2(1 + c)$ .

### Examples XIII(a). [ Pages 135-137 ]

1. (i) 6; (ii) -3. 2. -2. 5.  $\frac{n}{n-1}$ . 9. (i) 1; (ii)  $1\frac{1}{2}$ .  
 10. I'1173942, '3861209. 13. 2'425805. 14. '41369.  
 15. (i) I'8969092; (ii) '898665. 16. 39'879.  
 17. (i) 13; (ii) 6; (iii) 25. 18. (i) 24; (ii) 4; (iii) 70.  
 19. (i)  $\frac{\log 2}{\log 3}$  i.e. '63.....; (ii)  $4 + \frac{\log 7}{\log 3}$  i.e. 5'77...  
 (iii)  $\frac{2 \log 7 - 3 \log 3}{6 \log 5 - \log 7 - 2 \log 3}$ ; i.e. '108...  
 (iv)  $x = \frac{\log 3}{\log 3 - \log 2} = 2'71$  nearly,  $y = \frac{\log 2}{\log 3 - \log 2} = 1'71$  nearly;  
 (v)  $\frac{2b(2a-b)}{5ab+3ac-2b^2-bc}$  and  $\frac{2ab}{5ab+3ac-2b^2-bc}$ ,  
 where  $a = \log 2$ ,  $b = \log 3$ ,  $c = \log 7$ .  
 20. (i)  $\log x = \frac{a+3b}{5}$ ,  $\log y = \frac{a-2b}{5}$ .

### Examples XIII(b). [ Pages 142-144 ]

1. 3'2766077. 2. I'3686646. 3. 37'6018. 4. '7400827.  
 5. '8455104;  $32^\circ 16' 21''$ . 6. '7928863.  
 7. 9'9440554, 10'1559446. 8.  $36^\circ 24' 36''$ .  
 9.  $58^\circ 13' 55''$ . 10. 9'6198509;  $22^\circ 36' 28''$ .  
 12. 10'0957589. 13. 9'9147334. 14. 9'8718486.  
 16.  $\theta = 50^\circ 7' 48''$  nearly. 17. '2394.

### Examples XIV(a). [ Pages 157-160 ]

23.  $120^\circ$ . 24.  $A = 60^\circ$ . 29.  $A = 90^\circ$ ,  $B = 30^\circ$ ,  $C = 60^\circ$ .  
 39.  $\sqrt{\frac{y}{s} + \frac{s}{x} + \frac{x}{y}}$ . 40. 84.

### Examples XIV(b). [ Pages 166-168 ]

15.  $r = 4$ ;  $R = 8\frac{1}{2}$ .



**Examples XV(a).** [ Pages 172-173 ]

1.  $35^\circ 5' 49''$ .      2.  $102^\circ 1' 28''$ .      3.  $58^\circ 59' 38''$ .
4.  $104^\circ 30'$ ;  $46^\circ 36'$ ;  $28^\circ 54'$ .      5. (i)  $88^\circ 59' 40.9''$ .  
(ii)  $78^\circ 27' 46.86''$ .      6. (i)  $48^\circ 11' 28''$ ;  $58^\circ 24' 43''$ ;  $73^\circ 23' 54''$ .  
(ii)  $132^\circ 34' 24''$ .      7.  $A=120^\circ$ ,  $B=45^\circ$ ,  $C=15^\circ$ .
8.  $A=45^\circ$ ,  $B=30^\circ$ ,  $C=105^\circ$ .      9.  $A=60^\circ$ ,  $B=38^\circ 11'$ ,  $C=81^\circ 49'$ .
10.  $A=105^\circ$ ,  $B=45^\circ$ ,  $C=30^\circ$ .      11.  $(\sqrt{3}+1) : \sqrt{6} : (\sqrt{3}-1)$ .
12.  $\sqrt{5}+1 : \sqrt{5}-1$ .      13.  $3 : 4 : 5$ .

**Examples XV(b).** [ Pages 176-178 ]

1.  $B=88^\circ 12' 48''$ ,  $C=21^\circ 47' 12''$ .
2.  $B=56^\circ 19' 46.3''$ ,  $C=63^\circ 40' 13.7''$ .
3.  $A=117^\circ 38' 45''$ ,  $B=27^\circ 38' 45''$ .
4.  $A=94^\circ 42' 54''$ ,  $B=25^\circ 17' 6''$ .
5.  $B=71^\circ 44' 29.5''$ ,  $C=48^\circ 15' 30.5''$ .
6. (i)  $70^\circ 53' 36''$ ;  $49^\circ 6' 14''$ .      (ii)  $74^\circ 13' 50''$ ,  $35^\circ 16' 10''$ .  
(iii)  $A=64^\circ 21'$ ,  $B=77^\circ 25'$ ,  $c=27.39$ .
7. (i)  $B=78^\circ 17' 39.6''$ ,  $C=49^\circ 36' 20.4''$ .  
(ii)  $116^\circ 33' 54''$ ;  $26^\circ 33' 54''$ .
8.  $A=B=75^\circ$ ,  $C=30^\circ$ ,  $b=2\sqrt{6}$ .      9.  $\sqrt{6}$ ,  $15^\circ$ ,  $105^\circ$ .
10. (i)  $A=45^\circ$ ,  $B=75^\circ$ ,  $c=\sqrt{6}$ .      (ii)  $A=80^\circ$ ,  $B=90^\circ$ .
11.  $27.0875$ .      12.  $172.6436$  ft.      13.  $79.063$ .
14. (i)  $A=31^\circ 20'$ ,  $b=185$ ,  $c=192$ .  
(ii)  $b=18.46$ ,  $c=37.16$ ,  $C=70^\circ 30'$ .      (iii)  $b=118.9$ ,  $c=117.2$ .
15.  $C=75^\circ$ ,  $a=c=2\sqrt{3}+2$ .      16.  $C=105^\circ$ ,  $a=\sqrt{2}$ ,  $c=\sqrt{3}+1$ .
17.  $72^\circ$ ,  $72^\circ$ ,  $36^\circ$ ; each side  $=\sqrt{5}+1$ .      18.  $8, 1$ .

**Examples XV(c).** [ Pages 184-185 ]

1. (i) One solution;      (ii) Ambiguous; two solutions;  
(iii) No solution;      (iv) One solution (right-angled triangle).
2. (i)  $C=75^\circ$ ,  $A=60^\circ$ ,  $a=\sqrt{6}$  } (ii)  $60^\circ$ , or,  $120^\circ$ .  
or  $C=105^\circ$ ,  $A=30^\circ$ ,  $a=\sqrt{2}$  }
3.  $A=45^\circ$ ,  $C=75^\circ$ ,  $c=\sqrt{3}+1$ . (no ambiguity).      4. Impossible.
5.  $C=58^\circ 56' 56.3''$  } or,  $C=121^\circ 3' 3.7''$  }  
 $A=87^\circ 48' 3.7''$  }  $A=25^\circ 41' 56.3''$  }
6.  $B=34^\circ 27'$ ,  $C=100^\circ 33'$ .
7.  $A=5^\circ 44' 21''$ .      8.  $A=33^\circ 39' 84''$ ,  $B=86^\circ 30' 26''$ .
9.  $A=80^\circ 36'$ ,  $C=64^\circ 14'$ ;      or,  $A=29^\circ 4'$ ,  $C=115^\circ 46'$ .

## Miscellaneous Examples II. [ Pages 186-188 ]

11. 4, 5, 6.                      14.  $B = 44^\circ 25' 39''$ , or,  $135^\circ 34' 4''$ .  
 21.  $\frac{1}{3}(n\pi + \frac{1}{2}\pi - (a+b+c))$ .      24.  $\frac{1}{3}(n\pi + \frac{1}{2}\pi)$ .

## Examples XVI. [ Pages 213-214 ]

4.  $\theta = \frac{1}{2}\pi$ .      5.  $x = 38^\circ 10'$  nearly.      6.  $\frac{1}{2}\pi$ .      7.  $-37$  nearly.  
 8. (i)  $x = 0$ ; (ii)  $46^\circ 25'$  (nearly) and  $90^\circ$ ; (iii)  $22\frac{1}{2}^\circ$  and  $112\frac{1}{2}^\circ$ ;  
 (iv)  $\frac{3}{2}\pi$ ; (v)  $14^\circ$  nearly; (vi)  $1'19$ ,  $2'72$ ,  $4'92$ .  
 (vii)  $1'16$ ,  $3'28$ ,  $4'95$ . (viii)  $\pm 82$ . (ix)  $64$ .

## Examples XVII(a). Pages 221-224 ]

4.  $\sqrt{2135}$  ft.      14.  $17^\circ 27' 30''$ .      18.  $18^\circ 26' 58''$  nearly.

## Examples XVII(b). [ Pages 231-233 ]

1.  $-\cos\left(a + \frac{\pi}{2n}\right) / \sin \frac{\pi}{2n}$ .      2. 0.  
 3.  $\frac{\sin\{a + \frac{1}{2}(n-1)(a+\pi)\} \sin \frac{1}{2}n(a+\pi)}{\sin \frac{1}{2}(a+\pi)}$ .  
 4.  $\frac{n}{2} + \frac{\sin n\theta}{2 \sin \theta} \cos(n+1)\theta$ .      5.  $\frac{1}{4} \left( \frac{3 \sin^2 na}{\sin a} - \frac{\sin^2 3na}{\sin 3a} \right)$ .  
 6.  $(-1)^{n-1} \frac{\sin n\theta \sin(n+1)\theta}{2 \cos \theta}$ .  
 7.  $\frac{3}{8}n - \frac{1}{2} \frac{\sin na}{\sin a} \cos(n+1)a + \frac{1}{8} \frac{\sin 2na}{\sin 2a} \cos 2(n+1)a$ .  
 8.  $\cos\{\theta + \frac{1}{2}(n-1)(\theta + \frac{1}{2}\pi)\} \frac{\sin \frac{1}{2}n(\theta + \frac{1}{2}\pi)}{\sin \frac{1}{2}(\theta + \frac{1}{2}\pi)}$ .  
 9.  $\frac{1}{4 \sin a} \left\{ (n+1) \sin 2a - \sin 2(n+1)a \right\}$ .  
 10.  $\frac{n}{2} \cos 2a + \frac{\cos 2(n+1)a \sin 2na}{2 \sin 2a}$ .      11.  $\frac{1}{2}$ .      12. 0.  
 13. 0.      14.  $\sin na$ .      16.  $\operatorname{cosec} a \{\tan(n+1)a - \tan a\}$ .  
 17.  $\operatorname{cosec} \theta \{\cot \theta - \cot(n+1)\theta\}$ .      18.  $\frac{1}{2} \operatorname{cosec} a \{\tan(n+1)a - \tan a\}$ .  
 19.  $\cot \theta \{\cot \theta - \cot(n+1)\theta\} - n$ .      20.  $\cot a - 2^n \cot 2^n a$ .  
 21.  $\frac{1}{2} \frac{\sin \frac{1}{2}n\theta}{\sin \frac{1}{2}\theta} \sin \frac{1}{2}(n+3)\theta - \frac{1}{4} \frac{\sin n\theta}{\sin \theta} \sin(n+3)\theta$ .  
 22.  $\frac{1}{2}(\tan 3^n x - \tan x)$ .      23.  $\tan^{-1} \frac{n}{2+n}$ .      24.  $\tan^{-1} \frac{n}{n+1}$ .  
 25.  $\tan^{-1} \frac{n}{n+1}$ .      26.  $\frac{1}{2^{n-1}} \cot \frac{\pi}{2^{n-1}} - 2 \cot 2\pi$ .

27.  $\frac{1}{4} \left[ \frac{\sin \frac{1}{2}nx}{\sin \frac{1}{2}x} \cos \frac{1}{2}(n+3)x(1+2 \cos 2x) + \frac{\sin \frac{1}{2}nx}{\sin \frac{1}{2}x} \cos \frac{1}{2}(n+3)x \right]$ .
28.  $\frac{(n+1) \cos n\theta - n \cos (n+1)\theta - 1}{2(1 - \cos \theta)}$ . 29. (a)  $\frac{1}{2}n(n+1)$ ; (b)  $n^2$ .
30.  $\cot x \tan (n+1)x - (n+1)$ ;  $\frac{1}{3}n(n+1)(n+2)$ .

### Examples XVII (c). [Pages 236-237]

1.  $(a^2 - b^2)^2 = ab$ . 2.  $a^2\{(x+b)^2 + y^2\} = (x^2 + y^2 - b^2)^2$ .
3.  $(x+3y)^2 = xy^2(x+2y)$ . 4.  $a^2 + b^2 = 1 + b^{\frac{2}{3}} - b^{\frac{4}{3}}$ .
5.  $(x+y)^{\frac{2}{3}} + (x-y)^{\frac{2}{3}} = 2$ . 6.  $(x^2 + y^2 + 2ax)^2 = 4a^2(x^2 + y^2)$ .
7.  $x^{\frac{2}{3}} + y^{\frac{2}{3}} = 4^{\frac{2}{3}}$ . 8.  $x^{\frac{4}{3}}y^{\frac{2}{3}} - x^{\frac{2}{3}}y^{\frac{4}{3}} = 1$ . 9.  $\frac{x^2}{b^2} + \frac{y^2}{a^2} = 1$ .
10.  $\frac{2x}{a} = \left( \frac{x^2}{a^2} + \frac{y^2}{b^2} \right) \left( \frac{x^2}{a^2} + \frac{y^2}{b^2} - 3 \right)$ . 11.  $(ax)^{\frac{2}{3}} + (by)^{\frac{2}{3}} = (a^2 - b^2)^{\frac{2}{3}}$ .
12.  $\left( \frac{x}{a} + \frac{y}{b} \right)^{\frac{2}{3}} + \left( \frac{x}{a} - \frac{y}{b} \right)^{\frac{2}{3}} = 2$ . 13.  $x^{\frac{2}{3}}y^{\frac{2}{3}}(x^{\frac{2}{3}} + y^{\frac{2}{3}}) = 1$ .
14.  $3a - 2b = a^2$ . 15.  $\frac{x^2}{a} + \frac{y^2}{b} = a + b$ . 16.  $x^2 + y^2 - 2 \cos a = 2$ .
17.  $ab = (b-a) \tan a$ . 18.  $a + b = 2ab$ . 19.  $(ab - c)(a^2 + b^2) = 2ab$ .

### Miscellaneous Examples III. [Pages 253-260]

1.  $\frac{1}{15}\pi, \frac{1}{3}\pi, \frac{2}{3}\pi$ . 2.  $40^\circ, 60^\circ, 80^\circ, \frac{1}{3}\pi, \frac{1}{3}\pi, \frac{1}{3}\pi$ . 3.  $90^\circ, 22\frac{1}{2}^\circ, 67\frac{1}{2}^\circ$ .
4.  $1:36$ . 5.  $6\frac{1}{4}$  degrees. 6.  $54^\circ 46' 54''$ .
26.  $\frac{(t^2+1)(t^4+1)}{t^2}$ . 27. 1. 28.  $\frac{7}{15}$ . 29.  $\frac{1}{15}$ .
30.  $x = \frac{1}{3}\pi, y = \frac{1}{3}\pi$ . 31. (i) Possible; (ii) Impossible; (iii) Impossible.
32. (i)  $a^2 + b^2 = 2$ ; (ii)  $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 2$ ; (iii)  $\left( \frac{x}{a} \right)^{\frac{2}{3}} + \left( \frac{y}{b} \right)^{\frac{2}{3}} = 1$ .
33. 341.5 ft. approximately. 39. 125 yds. 40. 173.2 ft.
41.  $30^\circ$ . 42. 273.2 ft. 43. 30 miles per hour.
44. height = 22.5 ft., distance = 38.97 ft. 45.  $1, \frac{1}{3}, \frac{1}{3}$ .
46.  $\frac{2}{3}, \frac{1}{3}, \frac{1}{3}$ . 47.  $\frac{2mn}{m^2 - n^2}$ . 48.  $-\frac{1}{3}$ . 49.  $\frac{1}{3}, -\frac{1}{3}, \frac{1}{3}\sqrt{5}$ .
50.  $\frac{1}{15}\sqrt{2}$ . 51.  $5 + 2\sqrt{6}$ . 52.  $-\frac{1}{3}$ .
54.  $(2n+1)\frac{\pi}{2}$ , or,  $(2n+1)\pi \pm \frac{\pi}{2}$ , or,  $n\pi + (-1)^n\frac{\pi}{6}$ . 90. 107.2 ft.

# CALCUTTA UNIVERSITY QUESTIONS

1. (a) Show that  $\cot 2A + \tan A = \operatorname{cosec} 2A$ . [ See Ex. 3, Art. 38. ]

(b) If  $\alpha + \beta + \gamma = \pi$ , prove that

$$\cos^2 \alpha + \cos^2 \beta + \cos^2 \gamma + 2 \cos \alpha \cos \beta \cos \gamma = 1.$$

[ See Ex. 8, Art. 56. ]

2. (a)  $\sin^{-1} \frac{2a}{1+a^2} + \sin^{-1} \frac{2b}{1+b^2} = 2 \tan^{-1} x,$

show that  $x = \frac{a+b}{1-ab}.$  [ See Ex. 7, Art. 71. ]

(b) Solve  $\sin \theta + \sqrt{3} \cos \theta = \sqrt{2}$ . [ See Ex. 13, Examples XI. ]

3. (a) In a plane triangle, establish geometrically

$$\cos A = \frac{b^2 + c^2 - a^2}{2bc}$$

discussing separately the cases (i) when  $A$  is acute and (ii) when  $A$  is obtuse. [ See Art. 83. ]

(b) If  $B = 45^\circ$ ,  $C = 10^\circ$  and  $a = 200$  ft., find  $b$ , having given that  $\log 2 = .30103$ ,  $L \sin 55^\circ = 9.9139645$

$$\log 1726.4 = 3.2371414, \log 1726.5 = 3.2371666.$$

[ See Ex. 12, Examples XV(b). ]

4. Solve graphically the equation  $5 \sin \theta + 2 \cos \theta = 5$  between  $\theta = 0^\circ$  to  $\theta = 270^\circ$ . [ See Ex. 8(ii), Examples XVI. ]

1. (a) Prove geometrically that

$$\cos(A+B) = \cos A \cos B - \sin A \sin B. \quad [ \text{See Art. 33.} ]$$

(b) If  $A+B+C=\pi$ , prove that

$$\cos^2 \frac{A}{2} + \cos^2 \frac{B}{2} + \cos^2 \frac{C}{2} = 2 \left( 1 + \sin \frac{A}{2} \sin \frac{B}{2} \sin \frac{C}{2} \right).$$

2. (a) Show that  $\tan^{-1}(\frac{1}{2} \tan 2A) + \tan^{-1}(\cot A) + \tan^{-1}(\cot^3 A) = 0$ .

(b) Solve  $\tan x + \tan 2x + \tan x \tan 2x = 1$ . [ See Ex. 18, p. 110. ]

3. (a) In a plane triangle, prove that

$$\tan \frac{A}{2} = \sqrt{\frac{(s-b)(s-c)}{s(s-a)}}. \quad [ \text{See Art. 86.} ]$$

(b) The sides of a triangle are 7, 8, 9; determine the greatest angle, [ See Ex. 6(i), Examples XV(a). ]

4. Solve graphically the equation  $2 \sin^2 x = \cos 2x$  between

$$x = -\frac{\pi}{2} \text{ and } x = \frac{3\pi}{2}. \quad [\text{See Ex. 1, Art. 111.}]$$

1. (a) Prove geometrically that

$$\sin(A-B) = \sin A \cos B - \cos A \sin B,$$

when  $A$  and  $B$  are positive and acute and  $A > B$ . [See Art. 34.]

- (b) If  $A+B+C=\pi$ , prove that

$$\frac{\cos A}{\sin B \sin C} + \frac{\cos B}{\sin C \sin A} + \frac{\cos C}{\sin A \sin B} = 2. \quad [\text{See Ex. 11, p. 95.}]$$

$$\sqrt{2}. (a) \text{ Show that } \tan^{-1} \frac{1}{a+b} + \tan^{-1} \frac{b}{a^2+ab+1} = \tan^{-1} \frac{1}{a}.$$

[See Ex. 4(ii), p. 119.]

$$(b) \text{ Solve } \tan\left(\frac{\pi}{4} + \theta\right) + \tan\left(\frac{\pi}{4} - \theta\right) = 4. \quad [\text{See Ex. 17, p. 110.}]$$

3. (a) In a plane triangle, prove that

$$\tan \frac{B-C}{2} = \frac{b-c}{b+c} \cot \frac{A}{2}. \quad [\text{See Art. 89.}]$$

(b) Two sides of a triangle are 3 and 5 feet, and the included angle is  $120^\circ$ ; find the other angles. [See Ex. 1, p. 176.]

4. Solve graphically the equation  $\cot \theta - \tan \theta = 2$  between  $\theta = 0$  to  $\theta = \pi$ . [See Ex. 8(iii), p. 214.]

1. Prove geometrically  $\cos(A-B) = \cos A \cos B + \sin A \sin B$ , where the angles  $A$ ,  $B$ ,  $A-B$  are all positive and lie in the first quadrant. [See Art. 34.]

$$\text{Show that } \cos \frac{\phi-\theta}{2} - \sin \theta \sin \frac{\phi+\theta}{2} = \cos \theta \cos \frac{\phi+\theta}{2}.$$

Find  $\sin 18^\circ$ . [See Art. 53.]

2. For a triangle  $ABC$ , establish the formula

$$\sin A = \frac{2}{bc} \{s(s-a)(s-b)(s-c)\}^{\frac{1}{2}},$$

where  $s$  is the semi-perimeter of the triangle. [See Art. 87.]

Corresponding to the inequality  $a+b > c$  concerning the sides of a triangle, can you prove  $\sin A + \sin B > \sin C$ ?

3. A person walks one mile bearing an angle  $\theta_1$  with a fixed direction, and then another mile bearing  $\theta_2$  with the same direction. Find (a) final distance from the starting point and (b) final bearing.

Show that

$$\tan^{-1}x + \tan^{-1}y + \tan^{-1}z = \tan^{-1} \frac{x+y+z-xyz}{1-ys-zx-xy}. \quad [\text{See Art. 70.}]$$

4. Solve  $4 \sin \theta \cos \theta = 1 - 2 \sin \theta + 2 \cos \theta$  in the interval  $0 < \theta < \pi$ .

Draw the graph of  $3 \sin x + 4 \cos x$ . What is its maximum value?

1. Establish geometrically the formula

$$\sin(A-B) = \sin A \cos B - \cos A \sin B,$$

where  $A, B, A-B$  are positive and lie in the first quadrant.

[See Art. 34]

$PQR$  is a triangle and  $S$  is the projection of  $P$  on  $QR$  produced. If  $\angle PQS = 30^\circ$ ,  $\angle PRS = 45^\circ$ , and  $QR = 2$  ft, find  $RS$ .

2. Given  $A+B+C = \pi$ , show that

$$\checkmark (i) \tan A + \tan B + \tan C = \tan A \tan B \tan C;$$

[See Art. 56 Ex. 5]

$$\checkmark (ii) \text{ if } A = \tan^{-1} 2, B = \tan^{-1} 3, \text{ then } C = \pi/4.$$

3. In any triangle  $ABC$ , prove that

$$\tan \frac{B-C}{2} = \frac{b-c}{b+c} \cot \frac{A}{2}.$$

[See Art. 89]

If  $b = \sqrt{3}$ ,  $c = 1$  and  $A = 30^\circ$ , solve the triangle.

4. (i) What is meant by saying that the trigonometrical ratios are *periodic functions*? What values of  $x$  would indicate the end of the period beginning from  $x=0$  of  $\sin x$  and  $\sin \pi x$ ?

[See Art. 102 Note]

Sketch a period of the tangent-graph,  $y = \tan x$ , including  $x = \pi/2$ , and discuss the behaviour of the graph near  $x = \pi/2$ .

[See Art. 106 and Note]

- (ii) Solve the equation  $\sin 4\theta = \cos 3\theta + \sin 2\theta$  in  $0 < \theta < \pi$ .

1. (a) Define a *radian*. What is the length of an arc of a circle of radius  $r$  which subtends an angle of  $\theta$  radians at the centre? In a diagram with acute angle  $\theta$  justify that  $\theta > \sin \theta$ .

[See Art. 5 and Appendix Art. 21]

- (b) If  $A, B$  are positive and  $A+B$  is acute, establish geometrically

$$\cos(A+B) = \cos A \cos B - \sin A \sin B. \quad [\text{See Art. 33}]$$

2. (a) If  $A+B+C = 180^\circ$ , show that

$$1 + 4 \sin \frac{A}{2} \sin \frac{B}{2} \sin \frac{C}{2} = \cos A + \cos B + \cos C.$$

[See Art. 56 Ex. 4]

- (b) Prove that

$$\tan^{-1} \frac{1}{3} + \tan^{-1} \frac{1}{5} + \tan^{-1} \frac{1}{7} + \tan^{-1} \frac{1}{8} = \frac{\pi}{4}.$$

[ See Ex. 4 of Examples XII ]

3. (a) Assuming the formula of the type

$$c^2 = a^2 + b^2 - 2ab \cos C,$$

deduce that the area of the triangle  $ABC$  is given by

$$\{s(s-a)(s-b)(s-c)\}^{\frac{1}{2}}.$$

(b) From an aeroplane vertically over a straight horizontal road, the angles of depression of two consecutive milestones on opposite sides of the aeroplane are observed to be  $\alpha$  and  $\beta$ . Show that the height in miles of the aeroplane above the road is given by

$$\frac{\tan \alpha \tan \beta}{\tan \alpha + \tan \beta}.$$

4. (a) Sketch the graphs of  $y=x$ ,  $y=\sin x$  and  $y=\tan x$  in the range between  $-\pi/2$  and  $+\pi/2$  with reference to the same axes in  $x$  and  $y$ . From the nature of the graphs near the origin can you suggest any relation among them at the origin?

- (b) Solve
- $\cos \theta - \sin \theta = 1/\sqrt{2}$
- in
- $-\pi < \theta < +\pi$
- .

1. (a) Find the relation between a
- degree*
- and a
- radian*
- .

- (b) Assuming
- $A$
- and
- $B$
- to be positive and
- $A+B$
- to be acute,

$$\text{Prove that } \sin(A+B) = \sin A \cos B + \cos A \sin B.$$

[ See Art. 33 ]

2. (a) Show that
- $\cos A + \cos(120^\circ + A) + \cos(120^\circ - A) = 0$
- .

[ See Ex. 23(i) of Examples VIII ]

- (b) If
- $A+B+C=180^\circ$
- , show that

$$\sin 2A + \sin 2B + \sin 2C = 4 \sin A \sin B \sin C.$$

[ See Art. 56 Ex. 1 ]

3. (a) Prove that, in a triangle,

$$\frac{\sin A}{a} = \frac{\sin B}{b} = \frac{\sin C}{c}. \quad [\text{See Art. 82}]$$

(b) A spherical balloon whose radius is  $r$  ft. subtends at an observer's eye an angle  $\alpha$ , when the angular elevation of its centre is  $\beta$ . Determine the height of the centre of the balloon.

4. (a) Draw the graphs of
- $y=\cos x$
- and
- $y=\sec x$
- , from
- $x=0$
- to
- $x=2\pi$
- .

[ See Art. 105 and 109 ]

- (b) Solve the equation
- $\sin 2\theta = \cos \theta$
- .

## PATNA UNIVERSITY QUESTIONS

1. (a) Obtain  $\tan (A+B+C)$  in terms of  $\tan A$ ,  $\tan B$ , and  $\tan C$ .

(b) Show that if an angle  $\alpha$  be divided into two parts, such that the ratio of the tangents of the parts is  $\lambda$ , then the difference  $x$  between the parts is given by

$$\sin x = \frac{\lambda - 1}{\lambda + 1} \sin \alpha.$$

2. (a) If  $A+B+C=180^\circ$ , prove that

$$\sin^2 \frac{A}{2} + \sin^2 \frac{B}{2} + \sin^2 \frac{C}{2} = 1 - 2 \sin \frac{A}{2} \sin \frac{B}{2} \sin \frac{C}{2}.$$

- (b) Show that  $\cos 7^\circ 30' = \frac{1}{4}(-1 + \sqrt{2} + \sqrt{3})$ .

3. (a) Prove that

$$\left( \frac{\cos A + \cos B}{\sin A - \sin B} \right)^n + \left( \frac{\sin A + \sin B}{\cos A - \cos B} \right)^n = 2 \cot^n \frac{A-B}{2},$$

when  $n$  is even, and is zero when  $n$  is odd.

- (b) Draw the graph of  $y = \sec x$  from 0 to  $2\pi$ .

4. Prove that in a triangle,

$$\cos \frac{A}{2} = \sqrt{\frac{s(s-a)}{bc}}.$$

Find the greatest angle in a triangle whose sides are 5, 6, 7 ft. respectively, having given

$$\log 6 = .7781513$$

$$L \cos 39^\circ 14' = 9.8890644, \text{ diff. for } 60'' = .0001032.$$

5. A railway curve in the shape of a quadrant of a circle, has  $n$  telegraph posts at its ends and at equal distances along the curve. A man stationed at a point  $P$  on one of the extreme radii produced sees the  $p$ th and  $q$ th posts from the end nearest him in a straight line. Show that the radius of the curve is

$$\frac{a}{2} \cos (p+q)\phi . \operatorname{cosec} p\phi . \operatorname{cosec} q\phi ,$$

where  $\phi = \frac{\pi}{4(n-1)}$ , and  $a$  is the distance of  $P$  from the nearest end of the curve.



1. (a) Find  $\sin 3A$  in terms of  $\sin A$ .

(b) Show that the equation  $\sec^2 \theta = \frac{4xy}{(x+y)^2}$  is only possible when  $x=y$ .

2. (a) If  $A+B=90^\circ$ , find the greatest value of  $\cos A \cos B$ .

(b) Prove that

$$\cos^4 \frac{\pi}{8} + \cos^4 \frac{3\pi}{8} + \cos^4 \frac{5\pi}{8} + \cos^4 \frac{7\pi}{8} = \frac{3}{2}.$$

3. (b) If  $A+B+C=180^\circ$ , prove that

$$\sin 2A + \sin 2B + \sin 2C = 4 \sin A \sin B \sin C.$$

(b) In a triangle  $ABC$ , if  $\tan \frac{A}{2}, \tan \frac{B}{2}, \tan \frac{C}{2}$  be in A.P., then show that  $\cos A, \cos B, \cos C$  are also in A.P.

4. (a) "If  $b, c$  and  $B$  be given, then solution of the triangle may be ambiguous." Discuss this statement in detail.

(b) Draw the graph of  $y = \tan x$  between  $-\pi$  and  $\pi$ .

5. The elevation of a tower due North at a station  $A$  is  $\alpha$ , and at a station  $B$  due West of  $A$  is  $\beta$ .

Prove that the altitude is

$$\frac{AB \cdot \sin \alpha \sin \beta}{\sqrt{\sin^2 \alpha - \sin^2 \beta}}.$$

1. (a) The angles of a triangle are in A.P., and the number of degrees in the least is to the number of radians in the greatest as 60 to  $\pi$ ; find the angles in degrees.

(b) Trace the graph of  $\sin x$  from  $x = -\pi$  to  $x = \pi$ .

2. (a) Prove that  $\cos^2 A \cdot \cos 3A + \sin^2 A \cdot \sin 3A = \cos^2 2A$ .

(b) If  $a$  and  $\beta$  are two distinct angles satisfying the equation  $a \cos \theta + b \sin \theta = c$ , show that  $\sin(a+\beta) = \frac{2ab}{a^2+b^2}$ .

3. (a) Prove that in any triangle,  $2bc \cos A = b^2 + c^2 - a^2$ .

(b) In a triangle, prove that

$$a^2 \cos(B-C) + b^2 \cos(C-A) + c^2 \cos(A-B) = 3abc.$$

4. If the angles of a triangle be in A.P. and the lengths of the greatest and least sides be 24 and 16 feet respectively, find the length of the third side and the angles, given  $\log 2 = .30103$ ,  $\log 3 = .4771213$  and  $L \tan 19^\circ 6' = 9.5894287$ , diff. for  $1' = 4084$ .

5. (a) If  $\alpha + \beta + \gamma = \frac{1}{2}\pi$ , prove that

$$\tan \beta \tan \gamma + \tan \gamma \tan \alpha + \tan \alpha \tan \beta = 1.$$

(b) A tower stands in a field whose shape is that of an equilateral triangle and whose side is 80 feet. It subtends angles at the three corners whose tangents are respectively  $\sqrt{3}+1$ ,  $\sqrt{2}$ ,  $\sqrt{3}$ . Find its height.

1. (a) Prove that  $\sin(A+B) = \sin A \cos B + \cos A \sin B$ , when  $A$  and  $A+B$  are both obtuse.

(b) Prove that  $\sin^2 A + \sin^2(120^\circ + A) + \sin^2(240^\circ + A) = -\frac{1}{4} \sin 3A$ .

2. (a) If  $A+B+C=180^\circ$ , prove that

$$\sin \frac{A}{2} + \sin \frac{B}{2} + \sin \frac{C}{2} = 1 + 4 \sin \frac{\pi-A}{4} \sin \frac{\pi-B}{4} \sin \frac{\pi-C}{4}.$$

(b) Prove that in a triangle,  $\tan \frac{A-B}{2} = \frac{a-b}{a+b} \cot \frac{C}{2}$ .

3. (a) Solve  $\sin 7x - \sin x = \sin 3x$ .

(b) The angles of a polygon (which has no re-entrant angle) are in A.P. The least angle is  $\frac{2\pi}{3}$  radians and the common difference is  $5^\circ$ . Find the number of sides.

4. (a) If  $a, b, A$  are given and if  $c_1, c_2$  are the two values of the third side in the ambiguous case, prove that if  $c_1 > c_2$ ,  $c_1 - c_2 = 2a \cos B$ .

(b) Draw the graph of  $y = \cos x$  between  $x = -\pi$  and  $x = \pi$ .

5. Two towers stand on a horizontal plane and their distance apart is 120 ft. A person standing successively at the bases observes that the angular elevation of one is double that of the other, but when half way between them, their elevations appear to be complementary. Show that the heights are 90 ft. and 40 ft. respectively.

1. (a) Find the value of  $\sin 18^\circ$ .

(b) If  $\tan \frac{\theta}{2} = \left(\frac{1+c}{1-c}\right)^{\frac{1}{2}} \tan \frac{\phi}{2}$ , prove that  $\cos \theta = \frac{\cos \phi - c}{1 - c \cos \phi}$ .

2. Prove that in a triangle  $ABC$ ,  $\frac{a}{\sin A} = \frac{b}{\sin B} = \frac{c}{\sin C}$ ,

and deduce that  $a \sin\left(\frac{A}{2} + B\right) = (b+c) \sin \frac{A}{2}$ .

3. (a) If  $A+B+C=\pi$ , prove that

$$\sin^2 A + \sin^2 B + \sin^2 C = 2 \sin A \cos B \sin C.$$

(b) Trace the graph of  $\cos x$  between  $x=90^\circ$  and  $x=360^\circ$ , and find from the graph the value of  $\cos 160^\circ$  approximately.

4. At each end of a horizontal base of length  $2a$  it is found that the angular height of a certain peak is  $\theta$  and that at the middle point it is  $\phi$ . Prove that the vertical height of the peak is

$$\frac{a \sin \theta \sin \phi}{\sqrt{\sin(\phi+\theta) \sin(\phi-\theta)}}$$

5. (a) Prove that  $\log_b n = \log_a n \times \log_b a$ .

(b) In a triangle  $ABC$ ,  $a:b=7:3$ , and  $C=60^\circ$ ; find  $A$  and  $B$ , having given  $\log 2 = .3010300$ ,  $\log 3 = .4771213$ ;  $L \tan 34^\circ 42' = 9.8403776$ , difference for  $1' = 2699$ .

1. (a) Prove that  $\cos(A-B) = \cos A \cos B + \sin A \sin B$ , where  $A$  is obtuse and  $A-B$  is acute.

(b) If  $\tan \theta = \sec 2a$ , prove that  $\sin 2\theta = \frac{1 - \tan^4 a}{1 + \tan^4 a}$ .

2. In any triangle  $ABC$ , prove that (i)  $\tan \frac{B-C}{2} = \frac{b-c}{b+c} \cot \frac{A}{2}$ .

(ii)  $\sin(B+2C) + \sin(C+2A) + \sin(A+2B)$   
 $= 4 \sin \frac{1}{2}(B-C) \sin \frac{1}{2}(C-A) \sin \frac{1}{2}(A-B).$

3. (a) Express the cosine of an angle of a triangle in terms of its sides.

(b) In a triangle  $ABC$ , prove that

$$(b+c-a) \left( \cot \frac{B}{2} + \cot \frac{C}{2} \right) = 2a \cot \frac{A}{2}.$$

4. (a) Trace the graph of  $\sin x$  between  $x=0$  and  $x=2\pi$  and find from the graph the angles whose sine is .7.

(b) In a triangle  $ABC$ , if  $B=45^\circ$ ,  $C=10^\circ$  and  $a=200$  ft., find  $b$ , having given  $\log 2 = .30103$ ,  $L \sin 55^\circ = 9.9133645$ ,  $\log 172.64 = 2.2371414$ ,  $\log 172.65 = 2.2371666$ .

5. An object is observed from three points  $A, B, C$ , lying in a horizontal straight line which passes directly underneath the object; the angular elevation at  $B$  is twice that at  $A$ , and at  $C$  is three times that at  $A$ ; if  $AB=a$ ,  $BC=b$ , show that the height of the object is

$$\frac{a}{2b} \sqrt{\{(a+b)(3b-a)\}}.$$

1. (a) Draw the graph of  $y = \cos x$  for values of  $x$  varying from  $-\pi$  to  $\pi$ .

(b) If  $A + B + C = 180^\circ$ , show that

$$\sin \frac{A}{2} + \sin \frac{B}{2} + \sin \frac{C}{2} - 1 = 4 \sin \frac{\pi - A}{4} \sin \frac{\pi - B}{4} \sin \frac{\pi - C}{4}.$$

2. (a) If  $\tan \beta = \frac{n \sin a \cos a}{1 - n \sin^2 a}$ , show that  $\tan(a - \beta) = (1 - n) \tan a$ .

(b) Find a value of  $\theta$  satisfying the equation

$$\cos 3\theta + \cos 2\theta + \cos \theta = 0.$$

3. (a) In any triangle, prove that  $\tan \frac{A - B}{2} = \frac{a - b}{a + b} \cot \frac{C}{2}$ .

(b) If in any triangle  $a \cos^2 \frac{C}{2} + c \cos^2 \frac{A}{2} = \frac{3b}{2}$ , show that the sides of the triangle are in Arithmetical Progression.

4. The sides  $b$  and  $c$  of a triangle and the angle  $B$  are given. If  $b > c \sin B$  but  $< c$ , discuss the solution of the triangle.

If  $a_1$  and  $a_2$  are the values of the third side in the two solutions, prove that

$$a_1^2 + a_2^2 - 2a_1 a_2 \cos 2B = 4b^2 \cos^2 B.$$

5. The angles of elevation of a bird flying in a horizontal straight line from a fixed point at four successive observations are  $\alpha, \beta, \gamma, \delta$ , the observations being taken at equal intervals of time. Assuming the speed of the bird to be uniform, show that

$$\cot^2 \alpha - \cot^2 \delta = 3(\cot^2 \beta - \cot^2 \gamma).$$

1. (a) Draw the graph of  $y = \tan x$  from  $-\frac{\pi}{2}$  to  $\frac{\pi}{2}$ .

(b) Is it possible to find a value of  $\theta$  if  $\sec \theta = \frac{x^2 - y^2}{x^2 + y^2}$ ,  $x$  and  $y$  being two real and unequal numbers? Justify your answer.

2. (a) Find the value of  $\sin 18^\circ$ .

(b) If  $\frac{\tan(A - B)}{\tan A} + \frac{\sin^2 C}{\sin^2 A} = 1$ , prove that  $\tan A \tan B = \tan^2 C$ .

3. In any triangle, prove that

(i)  $c^2 = a^2 + b^2 - 2ab \cos C$ .

(ii)  $a \sin(B - C) + b \sin(C - A) + c \sin(A - B) = 0$ .

4. In a triangle,  $b=2.25$ ,  $c=1.75$ ,  $A=54^\circ$ . Find  $B$  and  $C$ , having given  $\log 2=.301030$ ,  $L \tan 63^\circ=10.292834$ ,  $L \tan 13^\circ 47'=9.889724$  and  $L \tan 13^\circ 48'=9.890270$ .

5. A man walking towards a building on which a flagstaff is fixed, observes the angle subtended by the flagstaff to be greatest when he is at a distance  $d$  from the building. If  $\theta$  be the observed greatest angle, find the length of the flagstaff and the height of the building.

1. (a) Given  $\sec \theta + \tan \theta = u$ , express  $\tan \theta$  in terms of  $u$ .

(b) Find all the values of  $\theta$  lying between  $0^\circ$  and  $360^\circ$  which satisfy the equation

$$\sec^2 \frac{\theta}{2} = 2 \sqrt{2} \tan \frac{\theta}{2}.$$

2. (a) Draw the graph of  $y = \tan x$  from  $-\pi$  to  $\pi$ .

(b) If  $A+B+C=\pi$ , prove that

$$\cos \frac{A}{2} + \cos \frac{B}{2} - \cos \frac{C}{2} = 4 \cos \frac{\pi+A}{4} \cos \frac{\pi+B}{4} \cos \frac{\pi-C}{4}.$$

3. (a) Prove that the value of  $\frac{\cot A}{1+\cot A} \times \frac{\cot (45^\circ-A)}{1+\cot (45^\circ-A)}$  is the same for all values of  $A$ .

(b) Show that  $\sin 20^\circ \sin 40^\circ \sin 60^\circ \sin 80^\circ = \frac{1}{16}$ .

4. (a) Discuss the ambiguous case in the solution of triangles.

(b) If a triangle is such that  $2b=a+c$ , prove that

$$2 \cot \frac{B}{2} = \cot \frac{A}{2} + \cot \frac{C}{2}.$$

5. Two points  $A$  and  $B$  of a straight horizontal road are at a distance of 400 feet apart. A vertical pole 100 feet high is at equal distances from  $A$  and  $B$ , and the angle subtended by  $AB$  at the foot of the pole (which is in the same horizontal plane as the road) is  $60^\circ$ . Find (a) the distance from the road to the foot of the pole, and (b) the cosine of the angle subtended by  $AB$  at the top of the pole.

1. (a) If  $\sin \alpha = -\frac{1}{2}$ , and  $\alpha$  lies between  $180^\circ$  and  $270^\circ$ , find the values of  $\sin \frac{\alpha}{2}$  and  $\cos \frac{\alpha}{2}$ .

(b) If  $\cos \theta = \frac{\cos \alpha - \cos \beta}{1 - \cos \alpha \cos \beta}$ , prove that one value of  $\tan \frac{\theta}{2}$  is  $\tan \frac{\alpha}{2} \cot \frac{\beta}{2}$ .

2. (a) Find the values of  $\theta$  lying between  $0^\circ$  and  $360^\circ$  satisfying the equation  $\tan^2 \theta + \cot^2 \theta = 2$ .

(b) If  $A+B+C=180^\circ$ , prove that

$$\sin^2 \frac{A}{2} + \sin^2 \frac{B}{2} + \sin^2 \frac{C}{2} = 1 - 2 \sin \frac{A}{2} \sin \frac{B}{2} \sin \frac{C}{2}.$$

3. A person walking along a straight road observes that the greatest angle which two objects subtends is  $\alpha$ . From the spot where this is the case he walks a distance  $c$  and the objects now appear as one, their direction making an angle  $\beta$  with the road. Shew that the distance between the objects is  $\frac{2c \sin \alpha \sin \beta}{\cos \alpha + \cos \beta}$ .

4. (a) Find the value of  $\tan \frac{A}{2}$  in terms of the sides of the triangle, adopting the usual notation.

(b) In any triangle, prove that

$$(b-c) \cos \frac{A}{2} = a \sin \frac{B-C}{2}.$$

5. Pick out, giving reasons, the ambiguous case out of the following and solve it.

(i)  $A=30^\circ$ ,  $c=250$  ft.,  $a=125$  ft.

(ii)  $A=30^\circ$ ,  $c=250$  ft.,  $a=200$  ft.,

given  $\log 2 = .30103$ ,  $\log 6.03893 = .7809601$ .

$L \sin 38^\circ 41' = 9.7958800$  and  $L \sin 8^\circ 41' = 9.1789001$ .

1. (a) Show that  $\cos^2 A \cos 3A + \sin^2 A \sin 3A = \cos^2 2A$ .

(b) If  $x \sin^2 \theta + y \cos^2 \theta = \sin \theta \cos \theta$ , and  $x \sin \theta - y \cos \theta = 0$ , show that  $x^2 + y^2 = 1$ .

2. (a) Establish the formula

$$\cos B - \cos A = 2 \sin \frac{A+B}{2} \sin \frac{A-B}{2}.$$

(b) Prove that  $\cos^2 A + \cos^2 B + \cos^2 C - 2 \cos A \cos B \cos C = 1$ , if  $A+B+C = \pi$ .

3. (a) Prove that in a triangle,  $\cos \frac{A}{2} = \sqrt{\frac{s(s-a)}{bc}}$ .

$$(b) \frac{a^2 \sin(B-C)}{\sin B + \sin C} + \frac{b^2 \sin(C-A)}{\sin C + \sin A} + \frac{c^2 \sin(A-B)}{\sin A + \sin B} = 0.$$

4. (a) Draw the graph of  $y = \sin x + \cos x$  as  $x$  ranges from 0 to  $\pi$ .

(b) Prove that  $\cot A + \cot B + \cot C = \cot A \cot B \cot C$ , if  $A+B+C = \frac{3}{2}\pi$ .

5. (a) Prove that  $\log_b n = \log_a n \times \log_a b$ .

(b) To determine the breadth  $AB$  of a canal an observer places himself at  $C$  in the straight line  $AB$  produced through  $B$ , and then walks 100 yards at right angles to this line. He then finds that  $AB$  and  $BC$  subtend angles  $15^\circ$  and  $25^\circ$  at his eyes. Find the breadth of the canal, given  $L \cos 25^\circ = 9.9572757$ ;  $L \cos 40^\circ = 9.8842540$ ;  $L \cos 75^\circ = 9.4129962$ ;  $\log 37279 = 4.5714643$ ;  $\log 3728 = 3.5714759$ .

1. (a) Evaluate  $\sin 18^\circ$ .

(b) If  $\sec(\phi + \alpha) + \sec(\phi - \alpha) = 2 \sec \phi$ , prove that

$$\cos \phi = \sqrt{2} \cos \frac{\alpha}{2}.$$

2. (a) If  $A + B + C = \pi$ , prove that

$$\cos \frac{A}{2} + \cos \frac{B}{2} + \cos \frac{C}{2} = 4 \cos \frac{\pi - A}{4} \cos \frac{\pi - B}{4} \cos \frac{\pi - C}{4}.$$

(b) Draw the graph of  $y = \tan x$  from  $x = 0$  to  $x = 2\pi$ .

3. In a triangle  $ABC$ , prove that

$$(i) \cos \frac{A}{2} = \sqrt{\frac{s(s-a)}{bc}}.$$

(ii)  $\cot A, \cot B, \cot C$  are in A. P., if  $a^2, b^2, c^2$  are in A. P.

4. Two sides of a triangle are in the ratio of 9 to 7, and the included angle is  $64^\circ 12'$ ; find the other angles, having given  $\log 2 = .3010300$ ,  $L \tan 57^\circ 54' = 10.2025255$ ,  $L \tan 11^\circ 16' = 9.2998216$ ,  $L \tan 11^\circ 17' = 9.2999804$ .

5. A flagstaff  $PN$  stands vertically on level ground. A base  $XY$  is measured at right angles to  $XN$ , the points  $X, Y, N$  being in the same horizontal plane, and the angles  $PXN$  and  $PYN$  are found to be  $\alpha$  and  $\beta$  respectively. Prove that the height of the flagstaff is

$$\frac{\sin \alpha \sin \beta}{\sqrt{\sin(\alpha - \beta) \sin(\alpha + \beta)}} \cdot XY.$$

1. (a) Obtain  $\cot 3A$  in terms of  $\cot A$ .

(b) Prove that

$$\cot A + \cot(60^\circ + A) + \cot(120^\circ + A) = 3 \cot 3A.$$

2. (a) Prove geometrically that

$$\cos(180^\circ + A) = -\cos A, \text{ for all values of } A.$$

(b) Solve

$$\sin 3A + \sin 2A + \sin A = 0, \text{ where } A \text{ lies between } 0 \text{ and } 2\pi.$$

3. In a triangle  $ABC$ , prove that

$$(i) \sin \frac{C}{2} = \sqrt{\frac{(s-a)(s-b)}{ab}},$$

$$(ii) (b-c) \cot \frac{A}{2} + (c-a) \cot \frac{B}{2} + (a-b) \cot \frac{C}{2} = 0.$$

4. The sides of a triangle are 4, 5, 6; find  $B$  having given  $\log 2 = .30103$ ,  $L \cot 27^\circ 53' = 9.9464040$ , diff. for  $1' = .0000669$ .

5. (a) The elevation of a steeple at a place due south of it is  $45^\circ$ , and at another place due west of the former place the elevation is  $30^\circ$ . If the distance between the two places be  $a$ , find the height of the steeple.

(b) Draw the graph of  $y = \cot x$  from  $x=0$  to  $x=\pi$ .

1. (a) If  $\tan \frac{x}{\theta+\alpha} = \tan \frac{y}{\theta+\beta} = \tan \frac{z}{\theta+\gamma}$ , prove that

$$\frac{x+y}{x-y} \sin^2(\alpha-\beta) + \frac{y+z}{y-z} \sin^2(\beta-\gamma) + \frac{z+x}{z-x} \sin^2(\gamma-\alpha) = 0.$$

(b) Prove that  $\cot A + \cot (60^\circ + A) + \cot (120^\circ + A) = 3 \cot 3A$ .

2. (a) If  $A+B+C=180^\circ$  and  $\sin \left(A + \frac{C}{2}\right) = n \sin \frac{C}{2}$ , show that

$$\tan \frac{A}{2} \tan \frac{B}{2} = \frac{n-1}{n+1}.$$

(b) If  $A+B+C=180^\circ$ , prove that

$$\sin^2 A + \sin^2 B + \sin^2 C - 2 \cos A \cos B \cos C = 2.$$

3. In a triangle, prove that

$$(i) \sin \frac{A}{2} = \sqrt{\frac{(s-b)(s-c)}{bc}},$$

$$(ii) (b^2 - c^2) \cot A + (c^2 - a^2) \cot B + (a^2 - b^2) \cot C = 0.$$

4. (a) Draw the graph of  $y = \sin x$  from  $x=0$  to  $x=\pi$ , and from the graph find the angles whose sine is .7.

(b) If  $a=70$ ,  $b=35$ ,  $C=36^\circ 52' 12''$ , find the other angles having given  $\log 3 = .4771213$ ,  $L \cot 18^\circ 26' 6'' = 10.4771213$ .

5. A flagstaff is on the top of a tower which stands on a level plane. At a certain point in the plane the tower subtends an angle  $\alpha$ , and the flagstaff an angle  $\beta$ . At another point ' $a$ ' ft. nearer the base of the tower, the flagstaff again subtends the angle  $\beta$ . Show that the height of the tower is  $\frac{a \tan \alpha}{1 - \tan \alpha \tan (\alpha + \beta)}$ .



# INTERMEDIATE EXAMINATION PAPERS

U. P.

ALLAHABAD

1. Show that  $\sin \theta = \frac{2 \tan \frac{\theta}{2}}{1 + \tan^2 \frac{\theta}{2}}$  and  $\cos \theta = \frac{1 - \tan^2 \frac{\theta}{2}}{1 + \tan^2 \frac{\theta}{2}}$ .

If  $\tan \frac{\theta}{2} = \sqrt{\frac{1-e}{1+e}} \tan \frac{\phi}{2}$ , show that  $\cos \phi = \frac{\cos \theta - e}{1 - e \cos \theta}$ .

2. (a) Solve the equation  $\tan \theta + \tan 2\theta + \tan 3\theta = 0$ .

(b) Show that  $\frac{\cos 6\theta + 6 \cos 4\theta + 15 \cos 2\theta + 10}{\cos 5\theta + 5 \cos 3\theta + 10 \cos \theta}$  can be reduced to the simple form  $2 \cos \theta$ .

3. Show that, in a triangle  $ABC$ , the distance of the ortho-centre from the side  $BC$  is  $2R \cos B \cos C$ ,  $R$  being the radius of the circum-circle.

Establish  $4R \sin A \sin B \sin C = a \cos A + b \cos B + c \cos C$ .

4. Prove that in a triangle,  $\cos \frac{C}{2} = \sqrt{\frac{s(s-c)}{ab}}$ .

Find the greatest angle of the triangle whose sides are 5, 6 and 7, having given

$$\log_{10} 6 = .7781518$$

$$L \cos 39^\circ 14' = 9.8890644$$

$$\text{diff. for } 60'' = .0001032.$$

5. The elevation of a tower due north of a point  $A$  is  $\theta$  and at a point  $B$  due west of  $A$  is  $\phi$ . Show that its altitude is

$$\frac{AB \sin \theta \sin \phi}{\sqrt{\sin^2 \theta + \sin^2 \phi}}.$$

1. Prove that  $\sin(\theta + \phi) = \sin \theta \cos \phi + \cos \theta \sin \phi$ , when  $\theta$  and  $\phi$  are both acute angles.

Show that if an angle  $\alpha$  be divided into two parts so that the ratio of the tangents of the parts is  $\lambda$ , the difference  $x$  between the parts is given by

$$\sin x = \frac{\lambda - 1}{\lambda + 1} \sin \alpha.$$

2. Determine the height of a mountain if the elevation of its top at unknown distance from the base is  $28^\circ$ ; and at a distance 8 miles 77 yards further off from the mountain along the same line, the angle of elevation is  $16^\circ$ . Given  $1.6071 = .2060$ ,  $L \sin 16^\circ = 9.4403$ ,  $L \sin 28^\circ = 9.6716$ ,  $L \sin 12^\circ = 9.3179$ .

3. Solve completely the following equations :—

(i)  $\cot \frac{1}{2}x - \cot x = \operatorname{cosec} \frac{1}{2}x$ . (ii)  $2 \sin^2 x + \sin^2 2x = 2$ .

4. Show that

$$(i) \cos \alpha + \cos \beta + \cos \gamma + \cos (\alpha + \beta + \gamma) \\ = 4 \cos \frac{\alpha + \beta}{2} \cos \frac{\beta + \gamma}{2} \cos \frac{\gamma + \alpha}{2}.$$

(ii)  $\tan A + \tan B + \tan C = \tan A \tan B \tan C$ , when  $A + B + C = \pi$ .

5. Establish the following relations in a triangle :—

(i)  $a \cot A + b \cot B + c \cot C = 2R + 2r$ . (ii)  $r_1 + r_2 + r_3 = r + 4R$ .

1. Prove that in any triangle  $ABC$ ,

$$\tan \frac{B-C}{2} = \frac{b-c}{b+c} \cot \frac{A}{2}.$$

2. (a) Show that in a triangle  $ABC$ ,

$$\sin A + \sin B + \sin C = 4 \cos \frac{A}{2} \cos \frac{B}{2} \cos \frac{C}{2}.$$

(b) Prove that in a triangle  $ABC$ ,

$$r_1 + r_2 = c \cot \frac{C}{2},$$

where  $r_1$  and  $r_2$  are the radii of the two escribed circles opposite to  $A$  and  $B$  respectively.

3. (a) Find the value of  $\sin 15^\circ$ .

(b) Find the number of digits in  $13^{12} \times 2^8$ , given

$$\log 2 = .30103 \text{ and } \log 3 = .4771213.$$

4. (a) What is the angle of elevation of the sun when the length of the shadow of a pole is  $\sqrt{3}$  times the height of the pole?

(b) Prove that the radian is a constant angle.

5. If  $(a-b)(s-c) = (b-c)(s-a)$ , show that the radii of the escribed circles of the triangle are in A.P., where  $a, b, c$  are the lengths of the sides of the triangle and  $s$  is semi-perimeter.

6. If  $K$  is a point in the side  $AB$  of a triangle  $ABC$  such that  $AK : KB = m : n$ , and if  $\theta$  be the  $\angle CKB$ , show that

$$(m+n) \cot \theta = n \cot A - m \cot B.$$

7. (a) If  $\tan \theta = k$ , find  $\tan 5\theta$ .

(b) Solve  $\sin 9\theta = \sin \theta$ .

8. In a triangle,  $a = 10$ ,  $A = 51^\circ 31' 40''$ ,  $B = 76^\circ$ ; find  $b$ ,

given  $\log 12396 = 4.0932816$ ,

$\log 12397 = 4.0933166$ ,

$L \sin 76^\circ = 9.9869041$ ,

$L \sin 51^\circ 30' = 9.8935444$ ,

$L \sin 51^\circ 31' = 9.8936448$ .

1. Prove that

$$(a) \frac{\sin(A+3B) \sin(3A+B)}{\sin 2A + \sin 2B} = 2 \cos A$$

$$(b) \tan^{-1} \frac{1}{4} + \tan^{-1} \frac{2}{9} = \frac{1}{2} \cos^{-1} \frac{3}{5}.$$

2. Solve the following equation, giving the general solution

$$\cos \theta - \sin 3\theta = \cos 2\theta.$$

3. A hillside is a plane sloping at  $30^\circ$  to the horizontal. For climbing the hill there is a straight road inclined at an angle of  $45^\circ$  to a line of the greatest slope. Find the tangent of the angle which the road makes with the horizontal.

4. (a) If  $\tan A + \sin A = m$  and  $\tan A - \sin A = n$ , prove that

$$m^2 - n^2 = 4 \sqrt{mn}.$$

(b) Prove that

$$\cos \tan^{-1} \sin \cot^{-1} x = \left( \frac{x^2 + 1}{x^2 + 2} \right)^{\frac{1}{2}}.$$

5. (a) State, without proof, the formulæ which you would use in solving a triangle when two sides and the included angle are known.

(b) A landmark  $A$  is observed from two points  $B, C$ , 4400 yds. apart. The angle  $ABC$  is found to be  $68^\circ$  and the angle  $ACB$   $72^\circ$ . Find the distance of the landmark from  $B$ .

[ Given  $\log 4.4 = 0.6435$ ,  $\log 6.51 = 0.8136$ ,

$L \sin 72^\circ = 9.9782$ ,  $\log \operatorname{cosec} 40^\circ = 0.1919$ . ]

6. (a) Find an expression for the radius of the circumscribed circle of a triangle.

(b) If  $r_1, r_2, r_3$  are the radii of the escribed circles of the triangle  $ABC$  opposite  $A, B, C$  respectively, and  $r$  is the radius of the inscribed circle, prove that

$$rr_1/r_2r_3 = \tan^2 \frac{1}{2} A.$$

1. State the value of  $\cos(90^\circ + \theta)$ , and prove the truth of your statement. ( $\theta < 90^\circ$ ).

What sign has  $\sin \theta + \cos \theta$  when  $\theta = 100^\circ$ ? Give reasons.

2. Prove that  $\sin A + \sin B = 2 \sin \frac{1}{2}(A+B) \cos \frac{1}{2}(A-B)$ .

Simplify

$$\frac{\sin A + \sin 3A + \sin 5A + \sin 7A}{\cos A + \cos 3A + \cos 5A + \cos 7A}$$

3. (a) An angle  $x$  is divided into two parts  $\alpha, \beta$  such that

$$\frac{\tan \alpha}{\tan \beta} = \frac{a}{b}.$$

Prove that

$$\sin(\alpha - \beta) = \frac{a-b}{a+b} \sin x.$$

(b) Eliminate  $\theta$  between the pair of equations :

$$x = 2 \cos \theta, y = 3 \cos(\theta - 30^\circ).$$

4. (a) Find the general solution of the equation

$$\cos 2\theta + \sin 2\theta = 1.$$

(b) If  $A+B+C=180^\circ$ , prove that

$$\sin^2 A + \sin^2 B + \sin^2 C = 2 + 2 \cos A \cos B \cos C.$$

5. At each end of a base of length  $2a$  the elevation of the top of a mountain is  $B$ , and at the middle point of the base the elevation is  $A$ . Prove that the height of the mountain is

$$\frac{a \sin A \sin B}{\sqrt{\{\sin(A+B) \sin(A-B)\}}}.$$

6. If  $AG$  bisects the angle  $A$  of a triangle  $ABC$ , and  $G$  lies on  $BC$ , find the length of  $AG$ .

In the triangle  $ABC$ ,  $D$  is the foot of the perpendicular from  $A$  on  $BC$  and  $A'$  is the middle point of  $BC$ . If  $AD=h$ ,  $AA'=m$ , and  $BC=a$ , prove that

$$\cot A = (4m^2 - a^2)/4ah.$$

1. Prove that in a triangle  $ABC$ ,

$$\tan \frac{B-C}{2} = \frac{b-c}{b+c} \cot \frac{A}{2}.$$

2. (a) Show that

$$\cos 2x = \frac{1 - \tan^2 x}{1 + \tan^2 x} \text{ and } \sin 2x = \frac{2 \tan x}{1 + \tan^2 x}.$$

- (b) If
- $2 \tan A = 3 \tan B$
- , show that

$$\tan (A - B) = \frac{\sin 2B}{5 - \cos 2B}.$$

3. (a) Prove that
- $\sin 15^\circ = (\sqrt{3} - 1)/2 \sqrt{2}$
- .

- (b) Solve the equation

$$\sin \theta + \cos \theta = \sqrt{2}.$$

4. Prove that
- $\log_a a = \log_a a / \log_a b$
- .

If  $x = \log_a a$ ,  $y = \log_a 2a$  and  $z = \log_a 3a$ , prove that  $xyz + 1 = 2yz$ .

5. Prove that in any triangle,

$$(i) \frac{b^2 - c^2}{a^2} \sin 2A + \frac{c^2 - a^2}{b^2} \sin 2B + \frac{a^2 - b^2}{c^2} \sin 2C = 0.$$

$$(ii) (r_1 - r)(r_2 - r)(r_3 - r) = 4r^2 R,$$

where the letters have their usual meanings.

6. (a) If
- $\tan \frac{\theta}{2} = \sqrt{\frac{1 - e}{1 + e}} \tan \frac{\phi}{2}$
- , prove that

$$\cos \phi = \frac{\cos \theta - e}{1 - e \cos \theta}.$$

- (b) Eliminate
- $\theta$
- between the equations

$$x = \sin (\theta + \alpha), y = \cos (\theta - \beta).$$

7. (a) Solve
- $\tan^{-1} 2x + \tan^{-1} 3x = \frac{1}{2}\pi$
- .

(b) Determine the height of a mountain if the elevation of its top at an unknown distance from the base is  $28^\circ$ , and at a distance 8 miles 77 yards further off from the mountain along the same line, the elevation is  $16^\circ$ , given

$$\log 1.6071 = .2060, \text{ L } \sin 12^\circ = 9.3179.$$

$$\text{L } \sin 16^\circ = 9.4403 \text{ and } \text{L } \sin 28^\circ = 9.6716.$$

8. Prove that the square of the distance between the circum-centre and the in-centre of any triangle is
- $R^2 - 2Rr$
- , in the usual notation.

If the circum-centre lies on the in-circle, prove that

$$\cos A + \cos B + \cos C = \sqrt{2}.$$

1. Show that
- $\cos 36^\circ = \frac{\sqrt{5} + 1}{4}$
- .

2. In any triangle, prove that

(i)  $2bc \cos A = b^2 + c^2 - a^2$ .

(ii)  $\sin 2A + \sin 2B + \sin 2C = 4 \sin A \sin B \sin C$ .

3. (a) Prove the identity

$$\tan\left(\frac{\pi}{4} + \frac{\theta}{2}\right) = \sqrt{\frac{1+\sin\theta}{1-\sin\theta}} = \tan\theta + \sec\theta.$$

(b) If  $\tan A + \sin A = m$  and  $\tan A - \sin A = n$ , show that  $m^2 - n^2 = 4\sqrt{mn}$ .

4. (a) Solve the equation  $\tan\theta + \tan 2\theta + \tan 3\theta = 0$ .

(b) If  $\tan\theta = \frac{1}{7}$  and  $\tan\phi = \frac{1}{3}$ , show that

$$\cos 2\theta = \sin 4\phi$$

5. (a) Eliminate  $\theta$  between the pair of equations

$$x = 2 \cos \theta, y = 3 \cos(\theta - 30^\circ).$$

(b) If  $K$  is a point in the side  $AB$  of a triangle  $ABC$  such that  $AK : KB = m : n$ , and  $\theta$  be the angle  $CKB$ , show that

$$(m+n) \cot \theta = n \cot A - m \cot B.$$

6. (a) The angle of elevation of the top of a pole is  $15^\circ$  from a point on the ground. On walking 100 feet towards the pole, the angle is found to be  $30^\circ$ . Find the height of the pole.

(b) Prove that  $\cos \tan^{-1} \sin \cot^{-1} x = \left(\frac{x^2+1}{x^2+2}\right)^{\frac{1}{2}}$ .

7. (a) Show that the radius of the circle circumscribing a regular polygon of  $n$  sides, equal to  $a$ , is given by

$$R = \frac{1}{2} a \operatorname{cosec} \frac{\pi}{n}.$$

(b) If the area of a triangle is 96 and the radii of its escribed circles are 8, 12, 24 respectively, calculate the length of the sides.

1. (a) If  $A$  is an angle between  $90^\circ$  and  $180^\circ$  and if  $\sin A = \frac{4}{5}$ , find  $\tan \frac{A}{2}$ .

(b) If  $A+B+C=\pi$ , prove that

$$\sin A - \sin B + \sin C = 4 \sin \frac{A}{2} \cos \frac{B}{2} \sin \frac{C}{2}.$$

2. (a) Prove geometrically the formula

$$\cos(A+B) = \cos A \cos B - \sin A \sin B.$$

(b) Show that  $\sin^2 24^\circ - \sin^2 6^\circ = \frac{\sqrt{5}-1}{8}$ .

3. (a) How would you solve a triangle having given two sides and the included angle?

(b) In a triangle  $ABC$ ,  $b=14$ ,  $c=11$ ,  $A=60^\circ$ ; find  $B$ , having given

$$L \tan 11^\circ 44' 29'' = 9.31774$$

$$\log 2 = .30103, \log 3 = .47712.$$

4. Show that

$$(a) \sqrt{r_1 r_2 r_3} = S = r^2 \cot \frac{A}{2} \cot \frac{B}{2} \cot \frac{C}{2},$$

$$(b) \cos A + \cos B + \cos C = 1 + \frac{r}{R},$$

where the symbols have their usual meanings.

5. (a) A spherical balloon of radius  $r$  feet subtends at an observer's eye an angle  $\alpha$ , when the angular elevation of its centre is  $\beta$ . Find the height of the centre of the balloon.

(b) Trace the changes in the sign and magnitude of  $\cos A - \sin A$  as  $A$  changes from 0 to  $2\pi$ .

6. (a) Prove that  $\cos A - \sqrt{3} \sin A = 2 \cos \left( A + \frac{\pi}{3} \right)$ ;

hence find the maximum value of

$$\cos A - \sqrt{3} \sin A.$$

(b) Solve the equation

$$7 \sin^2 x + 3 \cos^2 x = 4.$$

(c) If  $\cos^2 A + \cos^2 B + \cos^2 C + 2 \cos A \cos B \cos C = 1$ , shew that  $A \pm B \pm C$  is an odd multiple of  $\pi$ .

7. (a) Prove geometrically that

$$\tan A > A > \sin A.$$

(b) Shew by means of trigonometrical formulæ that if  $x+y+s=xyz$ , then

$$\frac{2x}{1-x^2} + \frac{2y}{1-y^2} + \frac{2s}{1-s^2} = \frac{2x}{1-x^2} \cdot \frac{2y}{1-y^2} \cdot \frac{2s}{1-s^2}.$$

1. (a) Prove geometrically the formula

$$\sin(A+B) = \sin A \cos B + \cos A \sin B.$$

(b) Solve the equation  $\sqrt{3} \sin \theta - \cos \theta = \sqrt{2}$ .

2. (a) In a triangle  $ABC$ , prove that

$$a \sin\left(\frac{A}{2} + B\right) = (b+c) \sin \frac{A}{2}.$$

- (b) Find the value of

$$2 \cos \frac{\pi}{13} \cos \frac{9\pi}{13} + \cos \frac{3\pi}{13} + \cos \frac{5\pi}{13}.$$

3. (a) Obtain an expression for the radius of an inscribed circle of a triangle.

- (b) If  $r_1, r_2, r_3$  be the radii of escribed circles and  $r$  the radius of inscribed circle of a triangle  $ABC$ , prove that

$$\frac{1}{r_1} + \frac{1}{r_2} + \frac{1}{r_3} + \frac{1}{r} = \frac{a^2 + b^2 + c^2}{S^2},$$

where  $S$  stands for the area of the triangle  $ABC$ .

4. (a) If  $\sin \theta = \sin \alpha$ , shew that all the values of  $\theta$  are included in the expression

$$n\pi + (-1)^n \alpha.$$

- (b) If  $\cos(A-B) = \frac{1}{2}$ , and  $\sin(A+B) = \frac{1}{2}$ , find the smallest positive values of  $A$  and  $B$ , and also their most general values.

5. (a) In an ambiguous case of the solution of a triangle, if  $a, b$ , and  $A$  are given, prove that the difference between the two values of  $c$  is  $2\sqrt{a^2 - b^2 \sin^2 A}$ .

- (b) Solve the triangle  $ABC$ , having given  $a=100$ ,  $c=100\sqrt{2}$ , and  $A=30^\circ$ .

6. (a) In a triangle  $ABC$ , if  $a^2, b^2, c^2$  are in A.P., show that  $\tan A, \tan B, \tan C$  are in H.P.

$$(b) \text{ If } \frac{a \cos A \sec B - x}{a \sin(A+B)} = \frac{y - b \sin A \sec B}{b \cos(A+B)} = \tan B,$$

$$\text{prove that } \frac{x^2}{a^2} + \frac{y^2}{b^2} = 1.$$

7. (a) Prove the rule which determines the characteristic of the logarithm (to base 10) of a number less than unity.

- (b) Find the smallest integral power  $a$ , of 7, which makes  $7^a$  greater than  $10^{50}$ .

$$(\log 7 = .8451)$$

- (c) Find the number of digits in  $3^{66}$ , having given  $\log 3 = .4771213$ .

1. (a) Deduce the signs of

- (i)  $\sin A$ , (ii)  $\cos A$  for all values of  $A$  between 0 and  $2\pi$ .



(b) Prove that

$$\frac{\sin(A-C) + 2 \sin A + \sin(A+C)}{\sin(B-C) + 2 \sin B + \sin(B+C)} = \frac{\sin A}{\sin B}.$$

2. (a) Prove geometrically the formula

$$\cos(A-B) = \cos A \cos B + \sin A \sin B.$$

(b) Solve the equation

$$\sin \frac{n+1}{2} \theta = \sin \frac{n-1}{2} \theta + \sin \theta.$$

3. In a triangle  $ABC$ , prove that

$$(i) \cot B \cot C + \cot C \cot A + \cot A \cot B = 1.$$

$$(ii) \sin(B+C-A) + \sin(C+A-B) + \sin(A+B-C) \\ = 4 \sin A \sin B \sin C.$$

4. In a triangle  $ABC$ , prove that

$$(i) (r_1 + r_2) \tan \frac{C}{2} = (r_2 - r) \cot \frac{C}{2} = c.$$

(ii) if  $8R^2 = a^2 + b^2 + c^2$ , then the triangle is a right-angled triangle.

5. The sides of a triangle are 32, 40 and 66 feet; find the angle opposite the greatest side, having given that

$$\log 207 = 2.3159703,$$

$$\log 1073 = 3.0305997,$$

$$L \cot 66^\circ 18' = 9.6424341,$$

$$\text{difference for } 1' = 3431.$$

6. (a) In a triangle  $ABC$ , prove that if

$$\cos A = \frac{\sin B}{2 \sin C},$$

then the triangle is isosceles.

(b) If  $\theta = \frac{\pi}{2^n + 1}$ , prove that

$$2^n \cos \theta \cos 2\theta \cos 2^2\theta \dots \cos 2^{n-1}\theta = 1.$$

7. A man notices two objects in a straight line due west. After walking a distance  $c$  due north, he observes that the objects subtend an angle  $\alpha$  at his eye; and, after walking a further distance  $c$  due north, an angle  $\beta$ . Show that the distance between the objects is

$$\frac{8c}{2 \cot \beta - \cot \alpha}.$$

## BENARES HINDU UNIVERSITY QUESTIONS

1. Define a radian, and show that if  $\theta$  be the circular measure on an angle subtended at the centre of a circle of radius  $r$  by an arc whose length is  $l$ ,  $\theta = \frac{l}{r}$ .

Two circles, the sum of whose radii is ' $a$ ', are placed in the same plane, with their centres at a distance ' $2a$ ', and an endless string, quite stretched, partly surrounds the circles and crosses itself between them. Show that the length of the string is  $(\frac{1}{2}\pi + 2\sqrt{3})a$ .

2. Prove that

$$(i) \ 1 - \frac{\sin^2 a}{1 + \cot a} - \frac{\cos^2 a}{1 + \tan a} = \sin a \cos a.$$

$$(ii) \ \tan 70^\circ = 2 \tan 50^\circ + \tan 20^\circ.$$

$$(iii) \ \sin^2 A + \sin^2 B - \sin^2 C = 2 \sin A \sin B \cos C, \\ \text{where } A + B + C = 180^\circ.$$

3. (a) Solve completely  $\cos \theta - \sin 3\theta = \cos 2\theta$ .

(b) If  $\sin^{-1} a + \sin^{-1} \beta + \sin^{-1} \gamma = \pi$ ,

$$\text{show that } a\sqrt{1-a^2} + \beta\sqrt{1-\beta^2} + \gamma\sqrt{1-\gamma^2} = 2a\beta\gamma.$$

4. In any triangle, prove that

$$(i) \ \frac{a}{\sin A} = \frac{b}{\sin B} = \frac{c}{\sin C}; \quad (ii) \ \cos A = \frac{b^2 + c^2 - a^2}{2bc}.$$

If  $\cos B = \frac{\sin A}{2 \sin C}$ , show that the triangle is isosceles.

5. The perpendiculars from the angles of a triangle on the opposite sides meet at  $O$ , and  $OA = x$ ,  $OB = y$ ,  $OC = z$ . Show that

$$\frac{a}{x} + \frac{b}{y} + \frac{c}{z} = \frac{abc}{xyz}.$$

6. A tower stood at the foot of an inclined plane whose inclination to the horizon was  $9^\circ$ . A line 100 ft. in length was measured straight up the incline from the foot of the tower, and at the end of this line the tower subtended an angle of  $54^\circ$ . Find the height of the tower, having given

$$\log 2 = .30103, \log 114.4123 = 2.0584726,$$

$$\text{and } L \sin 54^\circ = 9.9079576.$$

1. (a) Prove the following identities :—

$$(i) \ \cos^6 A + \sin^6 A = 1 - 3 \sin^2 A \cos^2 A.$$

$$(ii) \frac{\cos \theta}{1 - \tan \theta} + \frac{\sin \theta}{1 - \cot \theta} = \sin \theta + \cos \theta.$$

$$(b) \text{ If } 2 \tan \alpha = 3 \tan \beta, \text{ prove that } \tan(\alpha - \beta) = \frac{\sin \beta}{5 - \cos 2\beta}.$$

$$2. \text{ In any triangle, prove that } \tan \frac{B-C}{2} = \frac{b-c}{b+c} \cot \frac{A}{2}.$$

Two sides of a triangle are 540 and 420 yards long respectively and include an angle of  $52^\circ 6'$ . Find the remaining angles, given that  $\log 2 = .30103$ ,  $L \tan 26^\circ 3' = 9.6891480$ ,  $L \tan 14^\circ 20' = 9.4074189$ ,  $L \tan 14^\circ 21' = 9.4079458$ .

$$3. (a) \text{ Solve the equation } \sin(\theta - \phi) = \frac{1}{2}, \text{ and } \cos(\theta + \phi) = \frac{1}{2}.$$

$$(b) \text{ Prove that (i) } \sin^{-1} x + \cos^{-1} x = \frac{\pi}{2}.$$

$$(ii) \sin \cot^{-1} \cos \tan^{-1} x = \sqrt{\frac{x^2 + 1}{x^2 + 2}}.$$

$$4. (a) \text{ Prove geometrically the identity}$$

$$\cos(A - B) = \cos A \cos B + \sin A \sin B.$$

$$(b) \text{ In any triangle } ABC, \text{ prove that}$$

$$\frac{a^2 \sin(B - C)}{\sin B + \sin C} + \frac{b^2 \sin(C - A)}{\sin C + \sin A} + \frac{c^2 \sin(A - B)}{\sin A + \sin B} = 0.$$

5. An object is observed at three points  $A, B, C$  lying in a horizontal straight line which passes directly underneath the object. The angular elevation at  $B$  is twice that at  $A$ , and at  $C$  three times that at  $A$ ;  $AB = a$ ,  $BC = b$ . Show that the height of the object is  $\frac{a}{2b} \sqrt{(a+b)(3b-a)}$ .

6. If  $r_1, r_2, r_3$  be the radii of the three escribed circles, and  $\left(1 - \frac{r_1}{r_2}\right)\left(1 - \frac{r_2}{r_3}\right) = 2$ , show that the triangle must be right-angled.

$$1. (i) \text{ Show that } \frac{1}{\sec A - \tan A} = \sec A + \tan A.$$

$$(ii) \text{ If } \cos \theta - \sin \theta = \sqrt{2} \sin \theta, \text{ prove that } \cos \theta + \sin \theta = \sqrt{2} \cos \theta.$$

$$(iii) \text{ Prove that } \sin(A + B) \sin(A - B) = \sin^2 A - \sin^2 B.$$

$$2. \text{ Prove that}$$

$$(i) \tan 3A \tan 2A \tan A = \tan 3A - \tan 2A - \tan A.$$

$$(ii) \cos^4 \frac{\pi}{8} + \cos^4 \frac{3\pi}{8} + \cos^4 \frac{5\pi}{8} + \cos^4 \frac{7\pi}{8} = \frac{3}{2}.$$

$$3. (i) \text{ Find the value of } \sin 18^\circ.$$

$$(ii) \text{ If } A + B + C = 180^\circ, \text{ prove that}$$

$$\tan A + \tan B + \tan C = \tan A \tan B \tan C.$$

4. Solve any of the following equations :—

(i)  $\sin 5\theta \cos 3\theta = \sin 9\theta \cos 7\theta$ .

(ii)  $\tan \theta + \tan 2\theta + \tan 3\theta = 0$ .

(iii)  $\sin \theta + \cos \theta = \sqrt{2}$ .

5. (i) Discuss the ambiguous case in the solution of triangles.

(ii) In a  $\triangle ABC$ , if  $a=5$ ,  $b=4$ , and  $A=45^\circ$ , find the other angles, having given

$$\log 2 = .30103$$

$$L \sin 34^\circ 26' = 9.7523919,$$

and

$$L \sin 34^\circ 27' = 9.7525761.$$

6. (i) Find the radius of the circumscribed circle about a given triangle in terms of its sides and its area.

(ii) Prove that  $\frac{1}{bc} + \frac{1}{ca} + \frac{1}{ab} = \frac{1}{2\Delta r}$ ,

where  $a, b, c$  are the sides of a  $\triangle ABC$ ,  $R$  and  $r$  are respectively the radii of its circumscribed and inscribed circles.

1. (a) Prove that  $\sqrt{\frac{1 - \sin A}{1 + \sin A}} = \sec A - \tan A$ .

(b) What is the angle of elevation of the sun when the length of the shadow of a pole is  $\sqrt{3}$  times the height of the pole?

2. (a) Prove that  $\cos(A-B) = \cos A \cos B + \sin A \sin B$ .

(b) Prove that  $\frac{\sin(A-B)}{\cos A \cos B} + \frac{\sin(B-C)}{\cos B \cos C} + \frac{\sin(C-A)}{\cos C \cos A} = 0$ .

3. (a) Prove that  $\frac{\sec 8A - 1}{\sec 4A - 1} = \tan 8A$ .

(b) Prove that  $16 \cos \frac{2\pi}{15} \cos \frac{4\pi}{15} \cos \frac{8\pi}{15} \cos \frac{14\pi}{15} = 1$ .

4. Solve :—

(i)  $\tan^2 \theta - (1 + \sqrt{3}) \tan \theta + \sqrt{3} = 0$ .

(ii)  $\sqrt{3} \cos \theta + \sin \theta = \sqrt{2}$ .

5. (a) Prove that  $\log_a m = \log_a m \times \log_a b$ .

(b) In a triangle  $ABC$ , if  $a=21$ ,  $b=11$  and  $C=34^\circ 42' 30''$ , find  $A$  and  $B$ , given

$$\log 2 = .30103,$$

$$\text{and } L \tan 72^\circ 38' 45'' = 10.50515.$$

6. (a) Prove that  $\frac{1}{x} + \frac{1}{y} + \frac{1}{z} - \frac{1}{x+y+z} = 0$ ,

where  $r_1, r_2, r_3$  are the radii of the escribed circles and  $r$  the radius of the incircle of a triangle.

$$(b) \text{ Prove that } \sin^{-1} \frac{9}{5} - \cos^{-1} \frac{12}{13} = \sin^{-1} \frac{16}{65}.$$

$$1. (a) \text{ Prove that } \sin 2A = \frac{2 \tan A}{1 + \tan^2 A}.$$

$$(b) 2 \operatorname{cosec} 2A = \tan A + \cot A.$$

$$2. (a) \text{ Prove that } \sin(A-B) = \sin A \cos B - \cos A \sin B.$$

$$(b) \sin 10^\circ + \sin 20^\circ + \sin 40^\circ + \sin 50^\circ = \sin 70^\circ + \sin 80^\circ.$$

$$3. (a) \text{ Prove that } \tan 3A \tan 2A \tan A = \tan 3A - \tan 2A - \tan A.$$

$$(b) \text{ If } A+B+C=180^\circ, \text{ prove that}$$

$$\tan A + \tan B + \tan C = \tan A \tan B \tan C.$$

4. Solve :—

$$(a) \cos 3x + \cos 2x + \cos x = 0.$$

$$(b) \sin \theta + \cos \theta = \frac{1}{\sqrt{2}}.$$

5. (a) In any triangle, prove that

$$\tan \frac{B-C}{2} = \frac{b-c}{b+c} \cot \frac{A}{2}.$$

(b) If in a triangle,  $b=14$ ,  $c=11$ , and  $A=60^\circ$ , find  $B$  and  $C$  given that

$$\log 2 = .30103.$$

$$\log 3 = .4771218.$$

$$L \tan 11^\circ 44' = 9.3174299.$$

$$L \tan 11^\circ 45' = 9.3180640.$$

$$6. (a) \text{ Prove that } \frac{1}{bc} + \frac{1}{ca} + \frac{1}{ab} = \frac{1}{2Rr},$$

where  $R$  and  $r$  are the radii of the circum-circle and the in-circle respectively of the  $\triangle ABC$ .

$$(b) \text{ Prove that } \sin^{-1} \frac{8}{5} + \sin^{-1} \frac{8}{17} = \sin^{-1} \frac{77}{85}.$$

1. Prove that the circumference of a circle is equal to  $\pi$  times the diameter.

Examine the following statement :—

' $\pi=180^\circ$ ; therefore the circumference of a circle is 180 times the diameter'.

The angles of a triangle are in A. P., and the number of degrees in the least is to the circular measure of the greatest as  $60 : \pi$ ; find the angles.

2. (i) Prove that  $\frac{1}{\sec \theta - \tan \theta} - \frac{1}{\cos \theta} = \frac{1}{\cos \theta} - \sec \theta + \tan \theta$ .

(ii)  $\tan 7A - \tan 4A - \tan 3A = \tan 7A \tan 4A \tan 3A$ .

(iii)  $\sin^2 A + \sin^2 B + \sin^2 C = 2 \sin A \sin B \sin C$ ,  
given  $A + B + C = \pi$ .

3. Solve any two of the following :—

(i)  $\sin 4\theta = \sin \theta$ .

(ii)  $3 \cos \theta + \sqrt{3} \sin \theta = \sqrt{6}$ .

(iii)  $\cos^2 \theta - \sin \theta - \frac{1}{4} = 0$ .

4. In any triangle, prove that

(i)  $a = b \cos C + c \cos B$ .

(ii)  $\frac{a}{\sin A} = \frac{b}{\sin B}$ .

Prove that  $2a - b = 2c \cos B$ , given angle  $C = 60^\circ$ .

5. Prove  $r = \frac{S}{s}$  and state the corresponding results for the three escribed circles of the triangle, in the usual notation.

Prove that  $\frac{1}{r_1} + \frac{1}{r_2} + \frac{1}{r_3} = \frac{1}{r}$  and  $\frac{1}{ab} + \frac{1}{bc} + \frac{1}{ca} = \frac{1}{2Rr}$ .

6. Explain clearly what you mean by inverse trigonometrical functions and their principal values.

Prove that  $\tan^{-1} \frac{1}{2} + \tan^{-1} \frac{1}{3} = 1$  and  $\sin \operatorname{cosec}^{-1} \cot \tan^{-1} x = x$ .

7. A man on a boat moving direct towards a tower observes that at a certain point, the angular height of the tower is  $10^\circ$ . After advancing 50 yards nearer the tower, the elevation is observed to be  $15^\circ$ . Find the height of the tower above the water level; given

$$L \sin 15^\circ = 9.4129962,$$

$$L \cos 5^\circ = 9.9983442$$

$$\log 25.783 = 1.4113834$$

$$\text{and } \log 25.784 = 1.4118503.$$

1. (a) If  $\tan \theta + \sin \theta = a$  and  $\tan \theta - \sin \theta = b$ , then prove that  $a^2 - b^2 = 4\sqrt{ab}$ .

(b) Show that  $1 - \frac{\sin^2 \theta}{1 + \cot \theta} - \frac{\cos^2 \theta}{1 + \tan \theta} = \sin \theta \cos \theta$ .

2. (a) In a triangle  $ABC$ , prove that

$$\tan \frac{A}{2} = \sqrt{\frac{(s-b)(s-c)}{s(s-a)}}, \text{ where } 2s = a + b + c.$$

(b) The sides of a triangle are 130 ft., 123 ft. and 77 ft. Find the greatest angle, having given

$$\log 2 = .3010300,$$

$$L \tan 38^\circ 39' = 9.9029376, \quad L \tan 38^\circ 40' = 9.9031966.$$

3. (a) In any triangle  $ABC$ , prove that

$$\frac{a}{\sin A} = \frac{b}{\sin B} = \frac{c}{\sin C}.$$

(b) If in a triangle  $ABC$ ,

$$\cos A = \frac{\sin B}{2 \sin C},$$

prove that the triangle is isosceles.

4. (a) Solve the equation  $\sqrt{3} \cos \theta + \sin \theta = \sqrt{2}$ .

(b) Prove that  $\sin^{-1} \frac{1}{\sqrt{5}} + \cot^{-1} 3 = \frac{\pi}{4}$ .

5. (a) In any triangle  $ABC$ , prove that

$$\begin{aligned} & (\text{i}) \sin(B+2C) + \sin(C+2A) + \sin(A+2B) \\ & \quad = 4 \sin \frac{B-C}{2} \sin \frac{C-A}{2} \sin \frac{A-B}{2}. \end{aligned}$$

$$(\text{ii}) \frac{1}{r_1} + \frac{1}{r_2} + \frac{1}{r_3} = \frac{1}{r}.$$

where the symbols have their usual meanings.

6. In the same horizontal plane there are two inaccessible points  $P$  and  $Q$  and two stations  $S$  and  $T$  at each of which  $PQ$  is observed to subtend an angle  $\alpha$ .  $PT$  subtends at  $S$  an angle  $\beta$  and  $QS$  subtends at  $T$  an angle  $\gamma$ . Prove that

$$PQ = ST \cdot \frac{\sin \alpha}{\sin(\beta + \gamma - \alpha)}.$$

1. (a) If  $\sin^{-1} x + \sin^{-1} y = \frac{\pi}{2}$ , prove that

$$x \sqrt{1-y^2} + y \sqrt{1-x^2} = 1.$$

(b) Prove that the radian is a constant angle, and find its value in degrees, minutes and seconds.

2. (a) If  $A+B+C=180^\circ$ , prove that

$$\cos \frac{A}{2} + \cos \frac{B}{2} + \cos \frac{C}{2} = 4 \cos \frac{\pi-A}{4} \cos \frac{\pi-B}{4} \cos \frac{\pi-C}{4}.$$

(b) In any triangle, prove that

$$\frac{(r_1 + r_2)(r_2 + r_3)(r_3 + r_1)}{r_1 r_2 + r_2 r_1 + r_1 r_3} = 4R,$$

where the symbols have their usual meaning.

3. (a) Discuss the ambiguous case in the solution of triangles.

(b) In a triangle  $ABC$ , if  $a=5$ ,  $b=4$  and  $A=45^\circ$ , find the other angles having given

$$\begin{aligned}\log 2 &= .30103 ; \\ L \sin 34^\circ 26' &= 9.7523919 ; \\ L \sin 34^\circ 27' &= 9.7525761.\end{aligned}$$

4. Prove that

$$\begin{aligned}(i) \tan 70^\circ &= 2 \tan 50^\circ + \tan 20^\circ. \\ (ii) \cos \alpha \cos (60^\circ - \alpha) \cos (60^\circ + \alpha) &= \frac{1}{4} \cos 3\alpha. \\ (iii) \sin (\alpha + \beta + \gamma) + \sin (\alpha + \beta - \gamma) + \sin (\alpha - \beta + \gamma) \\ &= 4 \sin \alpha \cos \beta \cos \gamma, \\ &\text{provided } \alpha = \beta + \gamma.\end{aligned}$$

5. Solve the following equations :—

$$\begin{aligned}(a) \tan (\pi \cot \theta) &= \cot (\pi \tan \theta). \\ (b) \tan \theta + \tan 2\theta + \sqrt{3} \tan \theta \tan 2\theta &= \sqrt{3}.\end{aligned}$$

6. An object is observed at three points  $A, B, C$  lying in a horizontal straight line which passes directly underneath the object. The angular elevation at  $B$  is twice that at  $A$  and at  $C$  it is three times that at  $A$ ;  $AB=a$ ,  $BC=b$ . Show that the height of the object is

$$\frac{a}{2b} \sqrt{(a+b)(3b-a)}.$$

1. (a) Establish  $\cos (A+B) = \cos A \cos B - \sin A \sin B$ .

$$(b) \text{ Prove that } \cos 20^\circ \cdot \cos 40^\circ \cdot \cos 60^\circ \cdot \cos 80^\circ = \frac{1}{16}.$$

$$2. (a) \text{ Show that } \tan \frac{A}{2} = \sqrt{\frac{(s-b)(s-c)}{s(s-a)}}.$$

for any triangle  $ABC$ , each letter being used in its usual significance.

(b) The sides of a triangle are 2, 3 and 4; find the greatest angle, having given  $\log 2 = .30103$ ,  $\log 3 = .4771213$ ,  $L \tan 52^\circ 14' = 10.1103395$ ,  $L \tan 52^\circ 15' = 10.1111004$ .

3. (a) Prove the following :—

$$\frac{\tan A + \sec A - 1}{\tan A - \sec A + 1} = \frac{1 + \sin A}{\cos A}.$$

(b)  $\tan A + \tan B + \tan C = \tan A \tan B \tan C$ , if  $A+B+C=\pi$ .

$$4. (a) \text{ Prove that } \tan^{-1} \frac{1}{2} + \tan^{-1} \frac{1}{3} = \sin^{-1} \frac{1}{\sqrt{5}} + \cot^{-1} 3 = 45^\circ.$$

(b) The angles of a triangle are in A.P. and the number of radians in the least angle is to the number of degrees in the mean angle as 1 : 120. Find the angles in radians.

5. (a) Solve  $\sqrt{3} \cos \theta + \sin \theta = \sqrt{2}$ .

(b) A square tower stands upon a horizontal plane. From a point in this plane, from which three of its upper corners are visible,



their angular elevations are respectively  $45^\circ$ ,  $60^\circ$  and  $45^\circ$ . Show that the height of the tower is to the breadth of one of its sides as  $\sqrt{6}(\sqrt{5}+1)$  to 4.

1. Establish the following identities :—

$$(i) \frac{1}{\operatorname{cosec} A - \cot A} - \frac{1}{\sin A} = \frac{1}{\sin A} - \frac{1}{\operatorname{cosec} A + \cot A}.$$

$$(ii) \sin 65^\circ + \cos 65^\circ = \sqrt{2} \cos 20^\circ.$$

$$(iii) \tan \frac{A+B}{2} + \tan \frac{A-B}{2} = \frac{2 \sin A}{\cos A + \cos B}.$$

2. If  $R$  be the radius of the circle circumscribing the triangle  $ABC$  and  $p_1, p_2, p_3$  be the perpendiculars from the centre on the sides of the triangle, prove that

$$(i) \frac{a}{\sin A} = \frac{b}{\sin B} = \frac{c}{\sin C} = 2R,$$

$$(ii) \frac{a}{p_1} + \frac{b}{p_2} + \frac{c}{p_3} = \frac{abc}{p_1 p_2 p_3}.$$

3. In a triangle  $ABC$  calculate the sine of  $A/2$  in terms of the lengths of the sides and hence find the greatest angle of the triangle in which  $a=74$ ,  $b=26$  and  $c=60$ , given that

$$\log 3 = .4771213$$

$$\log 13 = 1.1139434$$

$$L \sin 56^\circ 19' = 9.9201836$$

$$\text{Diff. for } 1' = .0000342.$$

4. Solve the equations

$$(i) \cos \theta - \sin 3\theta = \cos 2\theta.$$

$$(ii) \tan \theta + \tan 2\theta + \tan 3\theta = 0.$$

5. The elevation of a tower at a place  $P$  due east of it is  $\theta$ ; and at  $Q$  due north of  $P$ , the elevation is  $\phi$ . Show that the height of the tower is

$$\frac{PQ \cdot \sin \theta \sin \phi}{\sqrt{\sin(\theta + \phi) \sin(\theta - \phi)}}.$$

6.-(a) Prove that

$$\tan^{-1} a + \tan^{-1} b = \tan^{-1} \frac{a+b}{1-ab}.$$

(b) Expand the determinant

$$\begin{vmatrix} 1 & & 1 \\ \sin A & \sin B & \sin C \\ \cos A & \cos B & \cos C \end{vmatrix}$$

and hence or otherwise prove that in a triangle  $ABC$

$$a \sin(B-C) + b \sin(C-A) + c \sin(A-B) = 0.$$

**TABLES OF LOGARITHMS, NATURAL SINES,  
NATURAL TANGENTS, LOGARITHMIC SINES,  
LOGARITHMIC TANGENTS ETC.**

TABLE I  
LOGARITHMS OF NUMBERS

	0	1	2	3	4	5	6	7	8	9		1	2	3	4	5	6	7	8	9
10	00000	00432	00860	01284	01703	02119	02531	02938	03342	03743		42	83	125	166	208	248	290	331	373
11	04189	04582	04922	05308	05690	06070	06446	06819	07188	07555		38	76	114	152	190	227	265	303	340
12	07918	08279	08636	08991	09343	09691	10037	10380	10721	11059		35	70	105	140	175	209	243	278	313
13	11394	11727	12057	12385	12710	13038	13364	13672	13988	14301		32	65	97	129	162	198	226	258	290
14	14613	14922	15229	15534	15836	16137	16435	16732	17026	17319		30	60	90	120	150	180	210	240	270
15	17609	17898	18184	18469	18752	19033	19312	19590	19866	20140		28	56	84	112	140	168	196	224	252
16	20412	20683	20952	21219	21484	21748	22011	22272	22531	22789		26	53	79	105	132	158	184	210	237
17	23045	23300	23553	23805	24055	24304	24551	24797	25042	25285		25	50	74	99	124	149	174	199	223
18	25527	25768	26007	26245	26482	26717	26951	27184	27416	27646		23	47	70	94	117	141	164	188	211
19	27875	28108	28330	28556	28780	28993	29226	29447	29667	29885		22	45	67	89	111	134	156	178	201
20	30108	30320	30535	30750	30963	31175	31387	31597	31806	32015		21	42	64	85	106	127	148	170	191
21	32222	32428	32634	32838	33041	33244	33445	33646	33846	34044		20	40	61	81	101	121	141	162	182
22	34243	34439	34635	34830	35025	35218	35411	35603	35793	35984		19	39	58	77	97	116	135	154	174
23	36173	36361	36549	36736	36922	37107	37291	37475	37658	37840		19	37	56	74	93	111	130	148	167
24	38021	38202	38382	38561	38739	38917	39094	39270	39445	39620		18	36	53	71	89	107	124	142	160
25	39794	39967	40140	40312	40483	40654	40824	40993	41162	41330		17	34	51	68	85	102	119	136	153
26	41497	41664	41830	41996	42160	42325	42488	42651	42813	42975		16	33	49	66	82	98	115	131	148
27	43186	43357	43527	43696	43861	44028	44191	44354	44516	44678		16	32	47	63	79	95	111	126	142
28	44716	44871	45025	45179	45332	45484	45637	45788	45939	46090		15	30	46	61	76	91	106	122	137
29	46240	46389	46538	46687	46835	46982	47129	47276	47422	47567		15	29	44	59	74	88	103	118	133



## LOGARITHMS OF NUMBERS

	Mean Differences									
	1	2	3	4	5	6	7	8	9	
55	74086	74115	74194	74373	74351	74429	74507	74586	74663	74741
56	74819	74896	74974	75051	75138	75205	75282	75358	75435	75511
57	75587	75664	75740	75815	75891	75967	76042	76118	76193	76268
58	76343	76418	76493	76567	76641	76716	76790	76864	76938	77012
59	77085	77159	77232	77305	77379	77452	77525	77597	77670	77743
60	77815	77887	77960	78032	78104	78176	78247	78319	78390	78462
61	78538	78604	78675	78746	78817	78888	78958	79029	79099	79169
62	79239	79309	79379	79449	79518	79588	79657	79727	79796	79865
63	79934	80003	80073	80140	80209	80277	80346	80414	80483	80550
64	80618	80686	80754	80821	80889	80956	81023	81090	81158	81224
65	81291	81358	81425	81491	81558	81624	81690	81757	81823	81889
66	81954	82020	82086	82151	82217	82282	82347	82413	82478	82543
67	82607	82672	82737	82802	82866	82930	82995	83059	83123	83187
68	83251	83315	83378	83442	83506	83569	83632	83696	83759	83822
69	83885	83948	84011	84073	84136	84198	84261	84323	84386	84448
70	84510	84572	84634	84696	84757	84819	84880	84942	85003	85065
71	85126	85187	85248	85309	85370	85431	85491	85552	85612	85673
72	85738	85794	85854	85914	85974	86034	86094	86153	86213	86273
73	86332	86392	86451	86510	86570	86629	86688	86747	86806	86864
74	86923	86982	87040	87099	87157	87216	87274	87332	87390	87448



TABLE II  
NATURAL SINES

	0'	10'	20'	30'	40'	50'	60'		Mean Differences									1'	2'	3'	4'	5'	6'	7'	8'	9'
0°	0.00000	0.00091	0.00362	0.00673	0.01164	0.01454	0.01745	89°										29	58	87	116	145	175	204	233	261
1°	.01745	.02086	.02327	.02619	.02908	.03190	.03490	88°										29	58	87	116	145	175	204	233	261
2°	.03490	.03781	.04071	.04363	.04653	.04943	.05234	87°										29	58	87	116	145	175	204	233	261
3°	.05234	.05524	.05814	.06105	.06395	.06685	.06976	86°										29	58	87	116	145	174	203	232	261
4°	.06976	.07266	.07556	.07846	.08136	.08426	.08716	85°										29	58	87	116	145	174	203	232	261
5°	.08716	.09005	.09295	.09585	.09874	.10164	.10453	84°										29	58	87	116	145	174	203	232	261
6°	.10453	.10742	.11031	.11320	.11609	.11898	.12187	83°										29	58	87	116	145	174	203	232	261
7°	.12187	.12476	.12764	.13053	.13341	.13629	.13917	82°										29	58	87	116	145	173	202	231	261
8°	.13917	.14205	.14493	.14781	.15069	.15356	.15643	81°										29	58	86	115	144	173	202	230	258
9°	.15643	.15931	.16218	.16505	.16792	.17078	.17365	80°										29	57	86	115	144	172	201	230	258
10°	.17365	.17651	.17937	.18224	.18509	.18795	.19061	79°										29	57	86	115	144	172	201	229	258
11°	.19061	.19366	.19652	.19937	.20222	.20507	.20791	78°										29	57	86	114	143	171	200	228	257
12°	.20791	.21076	.21360	.21644	.21927	.22212	.22495	77°										28	57	85	114	142	170	199	227	256
13°	.22495	.22778	.23062	.23345	.23627	.23910	.24192	76°										28	57	85	113	141	170	198	226	255
14°	.24192	.24474	.24756	.25038	.25320	.25601	.25882	75°										28	56	85	113	141	169	197	226	254
15°	.25882	.26163	.26443	.26724	.27004	.27284	.27564	74°										28	56	84	112	140	168	196	224	253
16°	.27564	.27843	.28123	.28402	.28680	.28959	.29237	73°										28	56	84	112	140	167	195	223	251
17°	.29237	.29515	.29793	.30071	.30348	.30625	.30902	72°										28	56	83	111	139	166	194	222	250
18°	.30902	.31178	.31454	.31730	.32006	.32282	.32557	71°										28	55	83	110	138	166	193	221	248
19°	.32557	.32833	.33106	.33381	.33655	.33929	.34202	70°										27	55	83	110	137	164	192	219	247

TABLE II.]

## NATURAL SINES AND COSINES

VII

	60'	50'	40'	30'	20'	10'	0'		1'	2'	3'	4'	5'	6'	7'	8'	9'
20°	0.84202	0.84475	0.84748	0.85021	0.85293	0.85565	0.85837	69°	27	55	83	109	137	164	191	218	246
21°	0.86887	0.86108	0.86379	0.86650	0.86921	0.87191	0.87461	68°	27	54	81	108	136	163	190	217	244
22°	0.87461	0.87780	0.88098	0.88367	0.88637	0.88905	0.89173	67°	27	54	81	108	135	161	188	215	242
23°	0.88073	0.88411	0.88748	0.89085	0.89422	0.89759	0.90096	66°	27	53	80	107	134	160	187	214	240
24°	0.88674	0.89039	0.89404	0.89769	0.90134	0.90499	0.90864	65°	27	53	80	106	133	159	186	212	238
25°	0.89263	0.89655	0.89788	0.89951	0.90113	0.90275	0.90437	64°	26	52	79	105	131	157	184	210	236
26°	0.89837	0.90259	0.90488	0.90680	0.90869	0.91058	0.91246	63°	26	52	78	104	130	156	182	208	234
27°	0.90399	0.90851	0.91103	0.91303	0.91488	0.91670	0.91851	62°	26	52	77	103	129	155	181	206	232
28°	0.90947	0.91439	0.91716	0.91916	0.92116	0.92316	0.92516	61°	26	51	77	102	128	154	179	204	230
29°	0.91481	0.91973	0.92248	0.92448	0.92648	0.92848	0.93048	60°	25	51	76	101	127	153	177	202	228
30°	0.92000	0.92552	0.93103	0.93655	0.94207	0.94759	0.95311	59°	25	50	75	100	125	150	175	200	225
31°	0.92504	0.93075	0.93646	0.94217	0.94788	0.95359	0.95930	58°	25	50	74	99	124	149	174	198	223
32°	0.92993	0.93584	0.94175	0.94766	0.95357	0.95948	0.96539	57°	25	49	74	98	123	147	172	196	221
33°	0.93464	0.94085	0.94706	0.95327	0.95948	0.96569	0.97190	56°	24	49	73	97	122	146	170	194	219
34°	0.93919	0.94570	0.95221	0.95872	0.96523	0.97174	0.97825	55°	24	48	72	96	120	144	168	192	216
35°	0.94358	0.95039	0.95720	0.96401	0.97082	0.97763	0.98444	54°	24	47	71	95	119	142	166	190	213
36°	0.94779	0.95490	0.96201	0.96912	0.97623	0.98334	0.99045	53°	23	47	70	94	117	140	164	187	211
37°	0.95183	0.95924	0.96665	0.97406	0.98147	0.98888	0.99629	52°	23	46	70	92	116	139	162	185	208
38°	0.95565	0.96336	0.97107	0.97878	0.98649	0.99420	1.00191	51°	23	46	68	91	114	137	159	182	205
39°	0.95932	0.96733	0.97534	0.98335	0.99136	0.99937	1.00738	50°	22	45	67	90	112	135	157	179	202
40°	0.96279	0.97110	0.97941	0.98772	0.99603	1.00434	1.01265	49°	22	44	66	88	111	133	155	177	199
41°	0.96606	0.97467	0.98328	0.99189	1.00050	1.00911	1.01772	48°	22	43	65	87	109	131	153	174	196
42°	0.96913	0.97804	0.98695	0.99586	1.00477	1.01368	1.02259	47°	21	43	64	86	107	129	150	172	193
43°	0.97200	0.98121	0.99042	0.99963	1.00884	1.01805	1.02726	46°	21	42	63	84	106	127	148	169	190
44°	0.97466	0.98417	0.99368	1.00319	1.01270	1.02221	1.03172	45°	21	42	62	83	104	124	145	166	187

NATURAL COSINES



## NATURAL SINES

	0'	10'	20'	30'	40'	50'	60'		Mean Differences									1'	2'	3'	4'	5'	6'	7'	8'	9'
45°	0.70711	0.70916	0.71121	0.71325	0.71529	0.71732	0.71934	44°	20	41	61	82	102	122	143	163	184									
46°	.71934	.72136	.72337	.72537	.72737	.72937	.73135	43°	20	40	60	80	100	120	140	160	180									
47°	.73135	.73338	.73531	.73728	.73924	.74120	.74314	42°	20	39	59	78	98	118	138	157	177									
48°	.74314	.74509	.74703	.74896	.75088	.75280	.75471	41°	19	39	58	77	96	116	135	154	173									
49°	.75471	.75661	.75851	.76041	.76229	.76417	.76604	40°	19	38	57	76	95	113	132	151	170									
50°	0.76604	0.76791	0.76977	0.77162	0.77347	0.77531	0.77715	39°	19	37	56	74	93	111	130	148	167									
51°	.77715	.77897	.78079	.78261	.78442	.78622	.78801	38°	18	36	54	72	91	109	127	145	163									
52°	.78801	.78980	.79158	.79335	.79512	.79688	.79864	37°	18	35	53	71	89	106	124	142	159									
53°	.79864	.80038	.80212	.80386	.80558	.80730	.80902	36°	17	35	52	69	87	104	121	138	156									
54°	.80902	.81072	.81242	.81412	.81580	.81748	.81915	35°	17	34	51	68	85	101	118	135	152									
55°	0.81915	0.82082	0.82248	0.82413	0.82577	0.82741	0.82904	34°	16	33	49	66	82	99	115	132	148									
56°	.82904	.83066	.83228	.83389	.83549	.83708	.83867	33°	16	32	48	64	80	96	112	128	144									
57°	.83867	.84025	.84183	.84339	.84495	.84650	.84805	32°	16	31	47	63	78	94	110	125	141									
58°	.84805	.84959	.85112	.85264	.85416	.85567	.85717	31°	15	30	46	61	76	91	106	122	137									
59°	.85717	.85866	.86015	.86163	.86310	.86457	.86603	30°	15	30	44	59	74	89	103	118	133									
60°	0.86603	0.86748	0.86892	0.87036	0.87178	0.87321	0.87462	29°	14	29	43	57	72	86	100	114	129									
61°	.87462	.87603	.87743	.87882	.88020	.88158	.88295	28°	14	28	42	55	69	83	97	111	125									
62°	.88295	.88431	.88566	.88701	.88835	.88968	.89101	27°	13	27	40	54	67	81	94	108	121									
63°	.89101	.89232	.89363	.89493	.89628	.89752	.89879	26°	13	26	39	52	65	78	91	104	117									
64°	.89879	.90007	.90133	.90259	.90383	.90507	.90631	25°	13	25	38	50	63	75	88	100	113									

TABLE II ]

## NATURAL SINES AND COSINES

IX

	66°	67°	68°	69°	70°	71°	72°	73°	74°	75°	76°	77°	78°	79°	80°	81°	82°	83°	84°	85°	86°	87°	88°	89°	90°
	0.90831	0.90753	0.90675	0.90596	0.90516	0.90437	0.90357	0.90276	0.90194	0.90111	0.90028	0.89944	0.89859	0.89773	0.89687	0.89599	0.89511	0.89421	0.89330	0.89238	0.89145	0.89051	0.88956	0.88860	0.88763
	0.91855	0.91772	0.91688	0.91603	0.91518	0.91432	0.91346	0.91259	0.91172	0.91084	0.90996	0.90907	0.90818	0.90728	0.90637	0.90545	0.90452	0.90359	0.90265	0.90171	0.90076	0.89981	0.89885	0.89789	0.89692
	0.92878	0.92792	0.92705	0.92617	0.92529	0.92440	0.92351	0.92261	0.92171	0.92080	0.91989	0.91897	0.91805	0.91712	0.91619	0.91525	0.91431	0.91337	0.91242	0.91147	0.91051	0.90955	0.90859	0.90762	0.90665
	0.93896	0.93808	0.93719	0.93629	0.93538	0.93446	0.93353	0.93260	0.93166	0.93072	0.92977	0.92882	0.92786	0.92690	0.92593	0.92496	0.92398	0.92300	0.92202	0.92103	0.92004	0.91905	0.91805	0.91705	0.91604
	0.94909	0.94817	0.94724	0.94630	0.94535	0.94439	0.94343	0.94246	0.94148	0.94050	0.93951	0.93852	0.93752	0.93651	0.93550	0.93448	0.93345	0.93242	0.93138	0.93033	0.92928	0.92822	0.92715	0.92608	0.92500
	0.95918	0.95823	0.95727	0.95630	0.95532	0.95434	0.95335	0.95236	0.95136	0.95035	0.94934	0.94832	0.94729	0.94625	0.94520	0.94415	0.94309	0.94202	0.94094	0.93985	0.93876	0.93766	0.93655	0.93543	0.93430
	0.96924	0.96827	0.96729	0.96630	0.96530	0.96429	0.96327	0.96224	0.96120	0.96015	0.95909	0.95802	0.95694	0.95585	0.95475	0.95364	0.95252	0.95139	0.95025	0.94910	0.94794	0.94677	0.94559	0.94440	0.94320
	0.97928	0.97829	0.97729	0.97628	0.97525	0.97421	0.97316	0.97210	0.97103	0.96995	0.96886	0.96776	0.96664	0.96551	0.96437	0.96322	0.96206	0.96089	0.95971	0.95852	0.95732	0.95611	0.95489	0.95365	0.95240
	0.98929	0.98828	0.98726	0.98623	0.98518	0.98412	0.98305	0.98197	0.98088	0.97978	0.97867	0.97754	0.97640	0.97525	0.97409	0.97291	0.97172	0.97052	0.96931	0.96809	0.96686	0.96562	0.96437	0.96311	0.96184
	0.99928	0.99825	0.99721	0.99616	0.99510	0.99402	0.99293	0.99183	0.99071	0.98958	0.98844	0.98728	0.98611	0.98493	0.98374	0.98254	0.98132	0.98009	0.97885	0.97760	0.97634	0.97507	0.97379	0.97250	0.97120
	1.00000	0.99999	0.99997	0.99994	0.99990	0.99985	0.99979	0.99972	0.99964	0.99955	0.99945	0.99934	0.99922	0.99909	0.99895	0.99880	0.99864	0.99847	0.99829	0.99811	0.99792	0.99773	0.99753	0.99732	0.99710
	1' 2'	1' 1'	1' 0'	0' 59"	0' 58"	0' 57"	0' 56"	0' 55"	0' 54"	0' 53"	0' 52"	0' 51"	0' 50"	0' 49"	0' 48"	0' 47"	0' 46"	0' 45"	0' 44"	0' 43"	0' 42"	0' 41"	0' 40"	0' 39"	0' 38"
	1' 2'	1' 1'	1' 0'	0' 59"	0' 58"	0' 57"	0' 56"	0' 55"	0' 54"	0' 53"	0' 52"	0' 51"	0' 50"	0' 49"	0' 48"	0' 47"	0' 46"	0' 45"	0' 44"	0' 43"	0' 42"	0' 41"	0' 40"	0' 39"	0' 38"

## NATURAL COSINES

TABLE III  
NATURAL TANGENTS

	0'	10'	20'	30'	40'	50'	60'		1'	2'	3'	4'	5'	6'	7'	8'	9'
0°	0.00000	0.00391	0.00782	0.00873	0.01164	0.01455	0.01746	89°	29	58	87	116	146	175	204	233	262
1°	0.01746	0.02087	0.02328	0.02619	0.02910	0.03201	0.03492	88°	29	58	87	116	146	175	204	233	262
2°	0.03492	0.03783	0.04075	0.04366	0.04658	0.04949	0.05241	87°	29	58	87	116	146	175	204	233	262
3°	0.05241	0.05533	0.05824	0.06116	0.06408	0.06700	0.06993	86°	29	58	88	117	146	175	204	234	263
4°	0.06993	0.07285	0.07578	0.07870	0.08163	0.08456	0.08749	85°	29	58	88	117	146	175	204	234	263
5°	0.08749	0.09042	0.09335	0.09629	0.09923	0.10216	0.10510	84°	29	59	88	118	147	176	206	235	265
6°	0.10510	0.10805	0.11099	0.11394	0.11688	0.11983	0.12278	83°	29	59	88	118	147	176	206	235	265
7°	0.12278	0.12574	0.12869	0.13165	0.13461	0.13758	0.14054	82°	30	59	89	118	148	178	207	237	266
8°	0.14054	0.14351	0.14648	0.14945	0.15243	0.15540	0.15838	81°	30	59	89	119	149	178	208	238	267
9°	0.15838	0.16137	0.16435	0.16734	0.17033	0.17333	0.17633	80°	30	60	90	120	150	179	209	239	269
10°	0.17633	0.17933	0.18233	0.18534	0.18835	0.19136	0.19438	79°	30	60	90	120	151	181	211	241	271
11°	0.19438	0.19740	0.20042	0.20345	0.20648	0.20952	0.21256	78°	30	61	91	121	152	182	212	242	272
12°	0.21256	0.21560	0.21864	0.22169	0.22475	0.22781	0.23087	77°	31	61	92	123	153	183	214	244	275
13°	0.23087	0.23393	0.23700	0.24008	0.24316	0.24624	0.24933	76°	31	62	92	123	154	185	216	246	277
14°	0.24933	0.25242	0.25552	0.25862	0.26173	0.26483	0.26795	75°	31	62	93	124	155	186	217	248	279
15°	0.26795	0.27107	0.27419	0.27732	0.28046	0.28360	0.28675	74°	31	63	94	125	157	188	219	250	283
16°	0.28675	0.28990	0.29305	0.29621	0.29938	0.30255	0.30573	73°	32	63	95	126	158	190	221	252	285
17°	0.30573	0.30891	0.31210	0.31530	0.31850	0.32171	0.32492	72°	32	64	96	128	160	192	224	256	288
18°	0.32492	0.32814	0.33136	0.33460	0.33788	0.34108	0.34433	71°	32	65	97	129	162	194	226	259	291
19°	0.34433	0.34758	0.35085	0.35412	0.35740	0.36068	0.36397	70°	33	65	98	131	164	196	229	262	294

TABLE III.]

## NATURAL TANGENTS

XI

	60'	50'	40'	30'	20'	10'	0'		1'	2'	3'	4'	5'	6'	7'	8'	9'
30°	0.58997	0.56797	0.54597	0.52397	0.50197	0.47997	0.45797	0.43597	0.41397	0.39197	0.36997	0.34797	0.32597	0.30397	0.28197	0.25997	0.23797
31°	0.59886	0.57686	0.55486	0.53286	0.51086	0.48886	0.46686	0.44486	0.42286	0.40086	0.37886	0.35686	0.33486	0.31286	0.29086	0.26886	0.24686
32°	0.60775	0.58575	0.56375	0.54175	0.51975	0.49775	0.47575	0.45375	0.43175	0.40975	0.38775	0.36575	0.34375	0.32175	0.29975	0.27775	0.25575
33°	0.61664	0.59464	0.57264	0.55064	0.52864	0.50664	0.48464	0.46264	0.44064	0.41864	0.39664	0.37464	0.35264	0.33064	0.30864	0.28664	0.26464
34°	0.62553	0.60353	0.58153	0.55953	0.53753	0.51553	0.49353	0.47153	0.44953	0.42753	0.40553	0.38353	0.36153	0.33953	0.31753	0.29553	0.27353
35°	0.63442	0.61242	0.59042	0.56842	0.54642	0.52442	0.50242	0.48042	0.45842	0.43642	0.41442	0.39242	0.37042	0.34842	0.32642	0.30442	0.28242
36°	0.64331	0.62131	0.59931	0.57731	0.55531	0.53331	0.51131	0.48931	0.46731	0.44531	0.42331	0.40131	0.37931	0.35731	0.33531	0.31331	0.29131
37°	0.65220	0.63020	0.60820	0.58620	0.56420	0.54220	0.52020	0.49820	0.47620	0.45420	0.43220	0.41020	0.38820	0.36620	0.34420	0.32220	0.30020
38°	0.66109	0.63909	0.61709	0.59509	0.57309	0.55109	0.52909	0.50709	0.48509	0.46309	0.44109	0.41909	0.39709	0.37509	0.35309	0.33109	0.30909
39°	0.66998	0.64798	0.62598	0.60398	0.58198	0.55998	0.53798	0.51598	0.49398	0.47198	0.44998	0.42798	0.40598	0.38398	0.36198	0.33998	0.31798
40°	0.67887	0.65687	0.63487	0.61287	0.59087	0.56887	0.54687	0.52487	0.50287	0.48087	0.45887	0.43687	0.41487	0.39287	0.37087	0.34887	0.32687
41°	0.68776	0.66576	0.64376	0.62176	0.59976	0.57776	0.55576	0.53376	0.51176	0.48976	0.46776	0.44576	0.42376	0.40176	0.37976	0.35776	0.33576
42°	0.69665	0.67465	0.65265	0.63065	0.60865	0.58665	0.56465	0.54265	0.52065	0.49865	0.47665	0.45465	0.43265	0.41065	0.38865	0.36665	0.34465
43°	0.70554	0.68354	0.66154	0.63954	0.61754	0.59554	0.57354	0.55154	0.52954	0.50754	0.48554	0.46354	0.44154	0.41954	0.39754	0.37554	0.35354
44°	0.71443	0.69243	0.67043	0.64843	0.62643	0.60443	0.58243	0.56043	0.53843	0.51643	0.49443	0.47243	0.45043	0.42843	0.40643	0.38443	0.36243

## NATURAL COTANGENTS

## NATURAL TANGENTS

	0'	10'	20'	30'	40'	50'	60'		Mean Differences									1'	2'	3'	4'	5'	6'	7'	8'	9'
45°	1.00000	1.00583	1.01170	1.01761	1.02355	1.02952	1.03553	44°										59	118	178	237	296	355	414	474	533
46°	.08553	.04158	.04768	.05378	.05994	.06618	.07237	43°										61	123	184	246	307	368	430	491	553
47°	.07287	.07864	.08496	.09131	.09770	.10414	.11061	42°										64	127	191	255	319	382	446	510	573
48°	.11081	.11713	.12369	.13029	.13694	.14363	.15037	41°										66	132	199	265	332	397	463	530	596
49°	.15037	.15715	.16398	.17085	.17777	.18474	.19175	40°										69	138	207	276	345	413	482	552	620
50°	1.19175	1.19893	1.20693	1.21310	1.22031	1.22758	1.23490	39°										72	144	216	288	360	431	503	575	647
51°	.28490	.24227	.24969	.25717	.26471	.27230	.27994	38°										75	150	225	300	376	451	526	601	676
52°	.27994	.28764	.29541	.30323	.31110	.31904	.32704	37°										78	157	235	314	392	471	549	628	707
53°	.32704	.33511	.34323	.35142	.35968	.36800	.37638	36°										82	164	247	329	411	493	576	658	740
54°	.37638	.38484	.39336	.40195	.41061	.41934	.42815	35°										86	172	259	345	431	517	603	690	776
55°	1.42815	1.43703	1.44598	1.45501	1.46411	1.47330	1.48256	34°										91	181	272	363	453	544	634	725	816
56°	.48356	.49190	.50133	.51084	.52043	.53010	.53987	33°										96	191	287	382	478	573	669	764	860
57°	.53987	.54972	.55966	.56969	.57981	.59002	.60038	32°										101	201	302	403	504	604	706	806	907
58°	.60038	.61074	.62125	.63185	.64256	.65337	.66428	31°										107	213	320	426	533	639	746	852	959
59°	.66428	.67530	.68643	.69766	.70901	.72047	.73205	30°										113	226	339	451	565	677	790	903	1016
60°	1.7391	1.7437	1.7556	1.7675	1.7796	1.7917	1.8040	29°										13	24	36	48	60	73	84	96	108
61°	1.8040	1.8165	1.8291	1.8418	1.8546	1.8676	1.8807	28°										13	25	38	51	64	77	89	102	115
62°	1.8907	1.9041	1.9174	1.9310	1.9447	1.9586	1.9726	27°										14	27	41	54	68	82	96	109	123
63°	1.9636	1.9768	1.9912	2.0057	2.0204	2.0353	2.0503	26°										15	29	44	58	73	88	102	117	131
64°	2.0503	2.0655	2.0809	2.0965	2.1123	2.1283	2.1445	25°										16	31	47	63	79	94	110	126	141

TABLE III.]

## NATURAL TANGENTS

XIII

	65°	66°	67°	68°	69°	70°	71°	72°	73°	74°	75°	76°	77°	78°	79°	80°	81°	82°	83°	84°	85°	86°	87°	88°	89°	90°
	2.1445	2.1609	2.1775	2.1943	2.2113	2.2286	2.2460	2.2635	2.2811	2.2988	2.3166	2.3345	2.3525	2.3705	2.3886	2.4068	2.4251	2.4435	2.4620	2.4806	2.4993	2.5181	2.5370	2.5560	2.5751	2.5943
	2.3460	2.3637	2.3817	2.3998	2.4183	2.4369	2.4559	2.4751	2.4945	2.5140	2.5336	2.5534	2.5733	2.5933	2.6134	2.6336	2.6539	2.6743	2.6948	2.7154	2.7361	2.7569	2.7778	2.7988	2.8198	2.8410
	2.8559	2.8750	2.8945	2.9142	2.9342	2.9545	2.9751	2.9958	3.0166	3.0376	3.0587	3.0799	3.1012	3.1226	3.1441	3.1657	3.1874	3.2092	3.2311	3.2531	3.2752	3.2974	3.3197	3.3421	3.3646	3.3872
	2.4751	2.4960	2.5172	2.5386	2.5605	2.5826	2.6051	2.6275	2.6501	2.6728	2.6958	2.7188	2.7419	2.7651	2.7884	2.8118	2.8353	2.8589	2.8826	2.9064	2.9303	2.9543	2.9784	3.0026	3.0269	3.0513
	2.6051	2.6379	2.6611	2.6846	2.7085	2.7328	2.7575	2.7825	2.8077	2.8331	2.8588	2.8847	2.9108	2.9371	2.9636	2.9903	3.0171	3.0441	3.0712	3.0985	3.1259	3.1535	3.1812	3.2091	3.2371	3.2652
	2.7475	2.7725	2.7980	2.8239	2.8502	2.8770	2.9042	2.9317	2.9595	2.9877	3.0161	3.0447	3.0735	3.1025	3.1317	3.1611	3.1907	3.2205	3.2505	3.2807	3.3111	3.3417	3.3725	3.4035	3.4347	3.4660
	2.9042	2.9319	2.9600	2.9887	3.0178	3.0475	3.0777	3.1081	3.1387	3.1696	3.2007	3.2320	3.2635	3.2952	3.3271	3.3592	3.3915	3.4240	3.4567	3.4896	3.5227	3.5559	3.5893	3.6229	3.6567	3.6906
	3.0777	3.1084	3.1397	3.1716	3.2041	3.2371	3.2705	3.3044	3.3387	3.3734	3.4084	3.4437	3.4793	3.5152	3.5513	3.5877	3.6243	3.6611	3.6982	3.7355	3.7730	3.8107	3.8486	3.8867	3.9250	3.9635
	3.3709	3.4052	3.4402	3.4759	3.5124	3.5495	3.5871	3.6251	3.6634	3.7020	3.7409	3.7800	3.8194	3.8591	3.8991	3.9393	3.9797	4.0203	4.0611	4.1022	4.1435	4.1850	4.2267	4.2686	4.3107	4.3530
	3.4874	3.5261	3.5656	3.6059	3.6470	3.6891	3.7321	3.7761	3.8201	3.8649	3.9106	3.9573	4.0049	4.0534	4.1019	4.1504	4.1990	4.2477	4.2965	4.3454	4.3944	4.4435	4.4927	4.5420	4.5914	4.6409
	3.7321	3.7760	3.8208	3.8667	3.9136	3.9617	4.0108	4.0611	4.1126	4.1653	4.2193	4.2747	4.3315	4.3898	4.4486	4.5079	4.5677	4.6279	4.6885	4.7496	4.8111	4.8731	4.9355	4.9984	5.0617	5.1254
	4.0108	4.0611	4.1126	4.1653	4.2193	4.2747	4.3315	4.3898	4.4486	4.5079	4.5677	4.6279	4.6885	4.7496	4.8111	4.8731	4.9355	4.9984	5.0617	5.1254	5.1895	5.2540	5.3189	5.3842	5.4499	5.5160
	4.3315	4.3898	4.4486	4.5079	4.5677	4.6279	4.6885	4.7496	4.8111	4.8731	4.9355	4.9984	5.0617	5.1254	5.1895	5.2540	5.3189	5.3842	5.4499	5.5160	5.5825	5.6494	5.7167	5.7844	5.8525	5.9210
	4.7046	4.7729	4.8430	4.9152	4.9894	5.0658	5.1445	5.2246	5.3051	5.3861	5.4676	5.5496	5.6321	5.7151	5.7986	5.8826	5.9671	6.0521	6.1376	6.2236	6.3101	6.3971	6.4846	6.5726	6.6611	6.7501
	5.1446	5.2257	5.3093	5.3955	5.4845	5.5764	5.6713	5.7692	5.8691	5.9710	6.0749	6.1808	6.2877	6.3956	6.5045	6.6154	6.7273	6.8402	6.9541	7.0690	7.1849	7.3018	7.4197	7.5386	7.6585	7.7794
	5.6713	5.7694	5.8708	5.9758	6.0844	6.1970	6.3138	6.4349	6.5606	6.6912	6.8269	6.9682	7.1154	7.2687	7.4287	7.5958	7.7704	7.9530	8.1443	8.3445	8.5545	8.7750	8.9969	9.2202	9.4450	9.6713
	6.3139	6.4349	6.5606	6.6912	6.8269	6.9682	7.1154	7.2687	7.4287	7.5958	7.7704	7.9530	8.1443	8.3445	8.5545	8.7750	8.9969	9.2202	9.4450	9.6713	9.9000	10.1311	10.3646	10.6005	10.8389	11.0798
	7.1164	7.2687	7.4287	7.5958	7.7704	7.9530	8.1443	8.3445	8.5545	8.7750	8.9969	9.2202	9.4450	9.6713	9.9000	10.1311	10.3646	10.6005	10.8389	11.0798	11.3231	11.5689	11.8172	12.0681	12.3215	12.5774
	8.3450	8.5545	8.7750	8.9969	9.2202	9.4450	9.6713	9.9000	10.1311	10.3646	10.6005	10.8389	11.0798	11.3231	11.5689	11.8172	12.0681	12.3215	12.5774	12.8343	13.0932	13.3551	13.6200	13.8879	14.1588	14.4327
	9.5144	9.7892	10.078	10.385	10.712	11.059	11.4301	11.8172	12.2202	12.6391	13.0640	13.4959	13.9348	14.3807	14.8336	15.2935	15.7604	16.2343	16.7152	17.2031	17.6980	18.2000	18.7091	19.2253	19.7486	20.2790
	11.4301	11.8326	12.251	12.705	13.197	13.737	14.301	14.889	15.502	16.140	16.803	17.491	18.204	18.942	19.705	20.494	21.308	22.148	23.014	23.906	24.824	25.768	26.739	27.737	28.762	29.814
	14.301	14.924	15.605	16.350	17.169	18.075	19.081	20.186	21.391	22.606	23.941	25.396	26.881	28.396	29.941	31.516	33.121	34.756	36.421	38.126	39.871	41.656	43.481	45.346	47.251	49.196
	19.081	20.206	21.470	22.904	24.542	26.432	28.636	31.184	34.126	37.501	41.359	45.751	50.726	56.344	62.656	69.721	77.596	86.341	96.026	106.721	118.506	131.371	145.406	160.721	177.426	195.521
	26.881	28.396	31.242	34.368	38.185	42.961	49.104	57.290	67.721	80.721	96.721	116.221	140.721	171.721	211.721	263.721	330.721	417.721	531.721	680.721	873.721	1119.721	1437.721	1847.721	2377.721	3057.721
	57.290	68.750	85.940	114.59	171.89	343.77	+ ∞																			
	60'	50'	40'	30'	20'	10'	0'																			
	1'	2'	3'	4'	5'	6'	7'	8'	9'																	

The differences change very rapidly here so that they cannot be tabulated.

The cotangent of a small angle of  $x'$  or the tangent of  $90^\circ - x'$  is very nearly equal to  $3437.7$  divided by  $x'$ .

## NATURAL COTANGENTS



TABLE IV ]

## LOGARITHMIC SINES

XV

	20°	21°	22°	23°	24°	25°	26°	27°	28°	29°	30°	31°	32°	33°	34°	35°	36°	37°	38°	39°	40°	41°	42°	43°	44°	
	9.53405	9.53751	9.54093	9.54433	9.54769	9.55102	9.55433	9.55758	9.56085	9.56408	9.56727	9.57044	9.57358	9.57668	9.57978	9.58284	9.58588	9.58889	9.59188	9.59484	9.59778	9.60070	9.60359	9.60646	9.60931	9.61214
	9.61494	9.61773	9.62049	9.62323	9.62595	9.62865	9.63133	9.63398	9.63662	9.63924	9.64184	9.64442	9.64698	9.64953	9.65205	9.65456	9.65705	9.65952	9.66197	9.66441	9.66683	9.66923	9.67161	9.67398	9.67633	9.67866
	9.68100	9.68334	9.68566	9.68794	9.69019	9.69242	9.69463	9.69682	9.69898	9.70112	9.70323	9.70532	9.70739	9.70944	9.71148	9.71350	9.71550	9.71748	9.71944	9.72138	9.72329	9.72518	9.72705	9.72890	9.73073	9.73254
	9.73433	9.73611	9.73787	9.73962	9.74135	9.74306	9.74475	9.74642	9.74807	9.74971	9.75133	9.75293	9.75451	9.75607	9.75762	9.75915	9.76067	9.76218	9.76368	9.76516	9.76662	9.76807	9.76951	9.77093	9.77234	9.77373
	9.77511	9.77648	9.77783	9.77916	9.78048	9.78179	9.78308	9.78435	9.78561	9.78686	9.78809	9.78931	9.79052	9.79172	9.79290	9.79407	9.79523	9.79638	9.79751	9.79863	9.79974	9.80084	9.80192	9.80299	9.80404	9.80508
	9.80611	9.80713	9.80814	9.80914	9.81012	9.81109	9.81205	9.81299	9.81392	9.81483	9.81572	9.81660	9.81746	9.81831	9.81915	9.82000	9.82082	9.82163	9.82242	9.82320	9.82397	9.82472	9.82546	9.82619	9.82691	9.82762
	9.82832	9.82902	9.82971	9.83039	9.83106	9.83172	9.83237	9.83301	9.83364	9.83426	9.83487	9.83547	9.83606	9.83664	9.83721	9.83777	9.83832	9.83886	9.83939	9.83991	9.84042	9.84092	9.84141	9.84189	9.84236	9.84282
	9.84327	9.84372	9.84417	9.84461	9.84504	9.84546	9.84588	9.84629	9.84669	9.84708	9.84746	9.84783	9.84820	9.84856	9.84891	9.84926	9.84960	9.84994	9.85027	9.85060	9.85092	9.85124	9.85155	9.85186	9.85216	9.85246
	9.85275	9.85304	9.85333	9.85361	9.85389	9.85416	9.85443	9.85469	9.85495	9.85520	9.85545	9.85570	9.85594	9.85618	9.85642	9.85665	9.85688	9.85711	9.85734	9.85756	9.85778	9.85800	9.85822	9.85843	9.85864	9.85885
	9.85905	9.85926	9.85946	9.85966	9.85986	9.86005	9.86024	9.86043	9.86062	9.86080	9.86098	9.86116	9.86134	9.86152	9.86170	9.86187	9.86204	9.86221	9.86238	9.86255	9.86271	9.86288	9.86304	9.86320	9.86336	9.86352
	9.86367	9.86383	9.86398	9.86413	9.86428	9.86443	9.86457	9.86472	9.86486	9.86500	9.86514	9.86528	9.86542	9.86556	9.86569	9.86583	9.86597	9.86610	9.86624	9.86637	9.86650	9.86663	9.86676	9.86689	9.86702	9.86715
	9.86728	9.86741	9.86754	9.86767	9.86780	9.86792	9.86805	9.86818	9.86831	9.86843	9.86856	9.86868	9.86881	9.86893	9.86905	9.86917	9.86929	9.86941	9.86953	9.86965	9.86976	9.86988	9.86999	9.87010	9.87021	9.87032
	9.87043	9.87054	9.87065	9.87076	9.87086	9.87097	9.87107	9.87117	9.87127	9.87137	9.87147	9.87156	9.87166	9.87175	9.87185	9.87194	9.87203	9.87212	9.87221	9.87230	9.87239	9.87248	9.87257	9.87266	9.87275	9.87284
	9.87293	9.87302	9.87311	9.87320	9.87329	9.87338	9.87347	9.87355	9.87364	9.87373	9.87381	9.87390	9.87398	9.87406	9.87415	9.87423	9.87431	9.87439	9.87447	9.87455	9.87463	9.87471	9.87479	9.87487	9.87495	9.87503
	9.87511	9.87519	9.87527	9.87535	9.87543	9.87551	9.87559	9.87566	9.87574	9.87582	9.87589	9.87597	9.87604	9.87611	9.87618	9.87625	9.87632	9.87639	9.87646	9.87653	9.87660	9.87667	9.87673	9.87680	9.87687	9.87693
	9.87700	9.87707	9.87714	9.87721	9.87728	9.87734	9.87741	9.87747	9.87754	9.87760	9.87767	9.87773	9.87779	9.87785	9.87791	9.87797	9.87803	9.87809	9.87815	9.87821	9.87827	9.87833	9.87839	9.87845	9.87851	9.87857
	9.87863	9.87869	9.87875	9.87881	9.87887	9.87893	9.87899	9.87905	9.87911	9.87917	9.87923	9.87929	9.87935	9.87941	9.87947	9.87953	9.87959	9.87965	9.87971	9.87977	9.87983	9.87989	9.87995	9.87999	9.88005	9.88011
	9.88017	9.88023	9.88029	9.88035	9.88041	9.88047	9.88053	9.88059	9.88065	9.88071	9.88077	9.88083	9.88089	9.88095	9.88101	9.88107	9.88113	9.88119	9.88125	9.88131	9.88137	9.88143	9.88149	9.88155	9.88161	9.88167
	9.88173	9.88179	9.88185	9.88191	9.88197	9.88203	9.88209	9.88215	9.88221	9.88227	9.88233	9.88239	9.88245	9.88251	9.88257	9.88263	9.88269	9.88275	9.88281	9.88287	9.88293	9.88299	9.88305	9.88311	9.88317	9.88323
	9.88329	9.88335	9.88341	9.88347	9.88353	9.88359	9.88365	9.88371	9.88377	9.88383	9.88389	9.88395	9.88401	9.88407	9.88413	9.88419	9.88425	9.88431	9.88437	9.88443	9.88449	9.88455	9.88461	9.88467	9.88473	9.88479
	9.88485	9.88491	9.88497	9.88503	9.88509	9.88515	9.88521	9.88527	9.88533	9.88539	9.88545	9.88551	9.88557	9.88563	9.88569	9.88575	9.88581	9.88587	9.88593	9.88599	9.88605	9.88611	9.88617	9.88623	9.88629	9.88635
	9.88641	9.88647	9.88653	9.88659	9.88665	9.88671	9.88677	9.88683	9.88689	9.88695	9.88701	9.88707	9.88713	9.88719	9.88725	9.88731	9.88737	9.88743	9.88749	9.88755	9.88761	9.88767	9.88773	9.88779	9.88785	9.88791
	9.88797	9.88803	9.88809	9.88815	9.88821	9.88827	9.88833	9.88839	9.88845	9.88851	9.88857	9.88863	9.88869	9.88875	9.88881	9.88887	9.88893	9.88899	9.88905	9.88911	9.88917	9.88923	9.88929	9.88935	9.88941	9.88947
	9.88953	9.88959	9.88965	9.88971	9.88977	9.88983	9.88989	9.88995	9.89001	9.89007	9.89013	9.89019	9.89025	9.89031	9.89037	9.89043	9.89049	9.89055	9.89061	9.89067	9.89073	9.89079	9.89085	9.89091	9.89097	9.89103
	9.89109	9.89115	9.89121	9.89127	9.89133	9.89139	9.89145	9.89151	9.89157	9.89163	9.89169	9.89175	9.89181	9.89187	9.89193	9.89199	9.89205	9.89211	9.89217	9.89223	9.89229	9.89235	9.89241	9.89247	9.89253	9.89259
	9.89265	9.89271	9.89277	9.89283	9.89289	9.89295	9.89301	9.89307	9.89313	9.89319	9.89325	9.89331	9.89337	9.89343	9.89349	9.89355	9.89361	9.89367	9.89373	9.89379	9.89385	9.89391	9.89397	9.89403	9.89409	9.89415
	9.89421	9.89427	9.89433	9.89439	9.89445	9.89451	9.89457	9.89463	9.89469	9.89475	9.89481	9.89487	9.89493	9.89499	9.89505	9.89511	9.89517	9.89523	9.89529	9.89535	9.89541	9.89547	9.89553	9.89559	9.89565	9.89571
	9.89577	9.89583	9.89589	9.89595	9.89601	9.89607	9.89613	9.89619	9.89625	9.89631	9.89637	9.89643	9.89649	9.89655	9.89661	9.89667	9.89673	9.89679	9.89685	9.89691	9.89697	9.89703	9.89709	9.89715	9.89721	9.89727
	9.89733	9.89739	9.89745	9.89751	9.89757	9.89763	9.89769	9.89775	9.89781	9.89787	9.89793	9.89799	9.89805	9.89811	9.89817	9.89823	9.89829	9.89835	9.89841	9.89847	9.89853	9.89859	9.89865	9.89871	9.89877	9.89883
	9.89889	9.89895	9.89901	9.89907	9.89913	9.89919	9.89925	9.89931	9.89937	9.89943	9.89949	9.89955	9.89961	9.89967	9.89973	9.89979	9.89985	9.89991	9.89997	9.90003	9.90009	9.90015	9.90021	9.90027	9.90033	9.90039
	9.90045	9.90051	9.90057	9.90063	9.90069	9.90075	9.90081	9.90087	9.90093	9.90099	9.90105	9.90111	9.90117	9.90123	9.90129	9.90135	9.90141	9.90147	9.90153	9.90159	9.90165	9.90171	9.90177	9.90183	9.90189	9.90195
	9.90201	9.90207	9.90213	9.90219	9.90225	9.90231	9.90237	9.90243	9.90249	9.90255	9.90261	9.90267	9.90273	9.90279	9.90285	9.90291	9.90297	9.90303	9.90309	9.90315	9.90321	9.90327	9.90333	9.90339	9.90345	9.90351
	9.90357	9.90363	9.90369	9.90375	9.90381	9.90387	9.90393	9.90399	9.90405	9.90411	9.90417	9.90423	9.90429	9.90435	9.90441	9.90447	9.90453	9.90459	9.90465	9.90471	9.90477	9.90483	9.90489	9.90495	9.90501	9.90507
	9.90513	9.90519	9.90525	9.90531	9.90537	9.90543	9.90549	9.90555	9.90561	9.90567	9.90573	9.90579	9.90585	9.90591	9.90597	9.90603	9.90609	9.90615	9.90621	9.90627	9.90633	9.90639	9.90645	9.90651	9.90657	9.90663
	9.90669	9.90675	9.90681	9.90687	9.90693	9.90699	9.90705	9.90711	9.90717	9.90723	9.90729	9.90735	9.90741	9.90747	9.90753	9.90759	9.90765	9.90771	9.90777	9.90783	9.90789	9.90795	9.90801	9.90807	9.90813	9.90819
	9.90825	9.90831	9.90837	9.90843	9.90849	9.90855	9.90861	9.90867	9.90873	9.90879	9.90885	9.90891	9.90897	9.90903	9.90909	9.90915	9.90921	9.90927	9.90933	9.90939	9.90945	9.90951	9.90957	9.90963	9.90969	9.90975
	9.90981	9.90987	9.90993	9.90999	9.91005	9.91011	9.91017	9.91023	9.91029	9.91035	9.91041	9.91047	9.91053	9.91059	9.91065	9.91071	9.91077	9.91083	9.91089	9.91095	9.91101	9.91107	9.91113	9.91119	9.91125	9.91131
	9.91137	9.91143	9.91149	9.91155	9.91161	9.91167	9.91173	9.91179	9.91185	9.91191	9.91															



## LOGARITHMIC SINES

	0'	10'	20'	30'	40'	50'	60'	Mean Differences							8'	9'
								1'	2'	3'	4'	5'	6'	7'		
45°	9.84249	9.85074	9.85900	9.86724	9.87548	9.88371	9.89193	44°	12	25	37	50	62	74	87	99 112
46°	9.85693	9.86515	9.87336	9.88156	9.88976	9.89795	9.90613	43°	12	24	36	48	60	72	84	96 108
47°	9.87107	9.87921	9.88734	9.89546	9.90357	9.91167	9.91976	42°	12	23	35	46	58	70	81	93 104
48°	9.88578	9.89387	9.90193	9.91000	9.91805	9.92609	9.93412	41°	11	22	34	45	56	67	78	89 100
49°	9.89996	9.90800	9.91602	9.92403	9.93203	9.93999	9.94794	40°	11	22	32	43	54	65	76	86 97
50°	9.91353	9.92153	9.92951	9.93747	9.94541	9.95333	9.96124	39°	10	21	31	42	52	62	73	83 94
51°	9.92650	9.93445	9.94238	9.95029	9.95818	9.96605	9.97390	38°	10	20	30	40	50	60	70	80 90
52°	9.93893	9.94683	9.95470	9.96254	9.97035	9.97813	9.98589	37°	10	19	29	39	49	58	68	78 87
53°	9.95083	9.95868	9.96650	9.97429	9.98205	9.98978	9.99749	36°	9	19	28	37	47	56	65	84 84
54°	9.96229	9.97008	9.97783	9.98554	9.99322	1.00087	1.00849	35°	9	18	27	36	45	54	63	72 81
55°	9.97329	9.98103	9.98873	9.99639	1.00401	1.01159	1.01914	34°	9	17	26	35	44	52	61	70 78
56°	9.98383	9.99152	9.99917	1.00678	1.01435	1.02188	1.02938	33°	8	17	25	34	42	50	59	67 76
57°	9.99391	1.00155	1.00915	1.01671	1.02423	1.03171	1.03916	32°	8	16	24	32	41	49	57	65 73
58°	1.00353	1.01113	1.01869	1.02621	1.03369	1.04113	1.04854	31°	8	16	23	31	39	47	55	62 70
59°	1.01274	1.02029	1.02780	1.03527	1.04270	1.05009	1.05744	30°	8	15	23	30	37	45	52	60 67
60°	1.02153	1.02903	1.03649	1.04391	1.05129	1.05863	1.06593	29°	7	14	22	29	36	43	50	57 64
61°	1.03000	1.03745	1.04486	1.05223	1.05956	1.06685	1.07410	28°	7	14	21	27	34	41	48	55 62
62°	1.03813	1.04554	1.05291	1.06024	1.06753	1.07478	1.08199	27°	7	13	20	26	33	40	46	53 59
63°	1.04603	1.05340	1.06073	1.06802	1.07527	1.08248	1.08965	26°	6	13	19	25	32	38	44	50 57
64°	1.05369	1.06102	1.06831	1.07556	1.08277	1.08994	1.09707	25°	6	12	18	24	30	36	42	48 54





	60'	50'	40'	30'	20'	10'	0'		1'	2'	3'	4'	5'	6'	7'	8'	9'
20°	9.56107	9.56498	9.56887	9.57274	9.57658	9.58039	9.58418	69°	39	77	116	154	198	231	270	308	347
21°	9.56418	9.56794	9.57168	9.57540	9.57909	9.58276	9.58641	68°	37	74	111	148	185	222	259	296	333
22°	9.56735	9.57104	9.57468	9.57827	9.58182	9.58533	9.58880	67°	36	72	107	143	179	214	250	286	322
23°	9.57052	9.57415	9.57773	9.58126	9.58475	9.58820	9.59161	66°	35	69	104	138	178	208	242	277	311
24°	9.57368	9.57725	9.58077	9.58424	9.58766	9.59104	9.59437	65°	34	67	101	134	168	201	235	268	302
25°	9.57683	9.58034	9.58380	9.58721	9.59057	9.59389	9.59716	64°	33	65	98	130	163	195	228	260	293
26°	9.57998	9.58344	9.58685	9.59021	9.59352	9.59679	9.60001	63°	32	63	95	126	158	190	221	253	284
27°	9.58313	9.58654	9.58990	9.59321	9.59648	9.59970	9.60287	62°	31	62	92	123	154	185	216	246	277
28°	9.58628	9.58964	9.59295	9.59621	9.59943	9.60261	9.60574	61°	30	60	90	120	151	181	211	241	271
29°	9.58943	9.59274	9.59600	9.59921	9.60238	9.60551	9.60860	60°	29	59	88	118	147	177	206	236	265
30°	9.59258	9.59584	9.59905	9.60221	9.60534	9.60843	9.61148	59°	29	58	87	116	144	173	202	231	260
31°	9.59573	9.59894	9.60211	9.60524	9.60833	9.61138	9.61439	58°	28	57	85	113	142	170	198	227	255
32°	9.59888	9.60204	9.60517	9.60826	9.61131	9.61432	9.61729	57°	28	56	84	112	139	167	195	223	251
33°	9.60203	9.60514	9.60821	9.61124	9.61423	9.61718	9.62010	56°	28	55	83	110	137	165	193	220	247
34°	9.60518	9.60824	9.61127	9.61426	9.61721	9.62014	9.62304	55°	27	54	81	108	136	162	190	217	244
35°	9.60833	9.61134	9.61431	9.61724	9.62014	9.62301	9.62584	54°	27	54	80	107	134	160	188	214	241
36°	9.61148	9.61444	9.61737	9.62026	9.62312	9.62595	9.62874	53°	26	53	79	106	132	158	185	212	238
37°	9.61463	9.61754	9.62041	9.62324	9.62603	9.62879	9.63152	52°	26	52	78	105	131	157	183	209	236
38°	9.61778	9.62064	9.62347	9.62626	9.62902	9.63175	9.63444	51°	26	52	78	104	130	156	182	208	234
39°	9.62093	9.62374	9.62651	9.62926	9.63198	9.63465	9.63728	50°	26	52	77	103	129	155	180	206	232
40°	9.62408	9.62684	9.62957	9.63226	9.63492	9.63754	9.64011	49°	26	51	77	102	128	154	179	205	230
41°	9.62723	9.62994	9.63261	9.63524	9.63783	9.64038	9.64289	48°	25	51	76	102	127	153	178	204	229
42°	9.63038	9.63304	9.63565	9.63822	9.64075	9.64324	9.64569	47°	25	51	76	101	127	152	177	203	228
43°	9.63353	9.63614	9.63871	9.64124	9.64373	9.64618	9.64859	46°	25	51	76	101	127	152	177	203	228
44°	9.63668	9.63924	9.64175	9.64422	9.64665	9.64904	9.65139	45°	25	51	76	101	127	152	177	203	228

## LOGARITHMIC TANGENTS

	0'	10'	20'	30'	40'	50'	60'	Mean Differences									
								1'	2'	3'	4'	5'	6'	7'	8'	9'	
45°	10.00000	10.00253	10.00505	10.00758	10.01011	10.01263	10.01516	44°	25	51	76	101	127	152	177	202	228
46°	.01516	.01769	.02022	.02275	.02528	.02781	.03034	43°	25	51	76	101	127	152	177	202	228
47°	.03034	.03288	.03541	.03795	.04048	.04302	.04556	42°	25	51	76	101	127	152	177	203	228
48°	.04556	.04810	.05065	.05319	.05574	.05829	.06084	41°	25	51	76	102	127	153	178	204	229
49°	.06084	.06339	.06594	.06850	.07106	.07362	.07619	40°	26	51	77	102	128	154	179	205	230
50°	10.07619	10.07875	10.08132	10.08390	10.08647	10.08905	10.09163	39°	26	52	77	103	129	155	180	206	232
51°	.09163	.09422	.09680	.09939	.10199	.10459	.10719	38°	26	52	78	104	130	156	182	208	234
52°	.10719	.10980	.11241	.11502	.11764	.12026	.12289	37°	26	52	78	105	131	157	183	209	236
53°	.12289	.12552	.12815	.13079	.13344	.13608	.13874	36°	26	53	79	106	132	158	185	212	238
54°	.13874	.14140	.14406	.14673	.14941	.15209	.15477	35°	27	54	80	107	134	160	188	214	241
55°	10.15477	10.15746	10.16016	10.16287	10.16558	10.16829	10.17101	34°	27	54	81	108	136	162	190	217	244
56°	.17101	.17374	.17648	.17922	.18197	.18472	.18748	33°	28	55	83	110	137	165	192	220	247
57°	.18748	.19025	.19303	.19581	.19860	.20140	.20421	32°	28	56	84	112	139	167	195	223	251
58°	.20421	.20703	.20985	.21268	.21552	.21837	.22123	31°	28	57	85	113	142	170	198	227	255
59°	.22123	.22409	.22697	.22985	.23275	.23565	.23856	30°	29	58	87	116	144	173	202	231	260
60°	10.23856	10.24148	10.24442	10.24736	10.25031	10.25327	10.25625	29°	29	59	88	118	147	177	206	236	265
61°	.25625	.25923	.26223	.26524	.26825	.27128	.27433	28°	30	60	90	120	151	181	211	241	271
62°	.27433	.27738	.28045	.28352	.28661	.28972	.29283	27°	31	62	92	123	154	185	216	246	277
63°	.29283	.29596	.29911	.30226	.30543	.30862	.31182	26°	32	63	95	126	158	190	221	253	284
64°	.31182	.31503	.31826	.32150	.32476	.32804	.33133	25°	33	65	98	130	163	195	228	260	293

TABLE V ]

## LOGARITHMIC TANGENTS

XXI

	1'	2'	3'	4'	5'	6'	7'	8'	9'
65°	34	67	101	134	168	201	235	268	302
66°	35	69	104	138	173	208	242	277	311
67°	36	72	107	143	179	214	251	286	322
68°	37	74	111	148	185	222	259	296	333
69°	39	77	116	154	193	231	270	308	347
70°	40	80	121	160	201	241	281	321	362
71°	42	84	126	168	210	252	294	336	378
72°	44	88	132	176	220	264	308	352	396
73°	46	93	139	186	232	278	325	371	418
74°	49	98	147	196	245	291	343	392	442
75°	52	104	156	208	261	313	365	417	469
76°	56	111	167	222	278	334	389	445	500
77°	60	120	179	239	299	359	419	478	538
78°	65	129	194	259	323	388	453	518	582
79°	71	141	212	282	354	420	494	564	635
80°	78	155	233	310	388	466	543	621	698
81°	87	173	260	346	433	519	606	692	779
82°	98	195	293	391	488	586	684	782	879
83°									
84°									
85°									
86°									
87°									
88°									
89°									
90°									

Differences vary so rapidly here that tabulation is impossible.

## SOME USEFUL CONSTANTS

One radian =  $57^{\circ} 17' 45''$  nearly = 206265 ;

$$\log 206265 = 5.3144255.$$

$$\pi = 3.14159265. \qquad \frac{1}{\pi} = 0.31830989.$$

$$\sqrt{2} = 1.4142135... \qquad \sqrt{3} = 1.7320508...$$

$$\sqrt{5} = 2.2360679... \qquad \sqrt{6} = 2.4494897...$$

$$\sqrt{7} = 2.6457513... \qquad \sqrt{8} = 2.8284271...$$

$$\sqrt{10} = 3.1622776...$$

## SOME USEFUL LOGARITHMS

$$\log 2 = .30103 \qquad \log 3 = .47712$$

$$\log 5 = .69897 \qquad \log 7 = .84510$$

---















